

ORBITS OF REAL FORMS IN COMPLEX FLAG MANIFOLDS

ANDREA ALTOMANI
COSTANTINO MEDORI
MAURO NACINOVICH

24/11/2006

ABSTRACT. We study, from the point of view of CR geometry, the orbits M of a real form $\mathbf{G}$ of a complex semisimple Lie group $\hat{\mathbf{G}}$ in a complex flag manifold $\hat{\mathbf{G}}/\mathbf{Q}$. In particular we characterize those that are of finite type and satisfy some Levi nondegeneracy conditions. These properties are also graphically described by attaching to them some cross-marked diagrams that generalize those for minimal orbits that we introduced in a previous paper. By constructing canonical fibrations over real flag manifolds, with simply connected complex fibers, we are also able to compute their fundamental group.

CONTENTS

1. Introduction
2. Homogeneous CR manifolds and CR algebras
3. Parabolic subalgebras and complex flag manifolds
4. Parabolic CR algebras and parabolic CR manifolds
5. Fit Weyl chambers and CR geometry of $M(\mathfrak{g}, \mathfrak{q})$
6. Canonical fibrations over a parabolic CR manifold
7. The isotropy subgroups
8. The fundamental group of parabolic CR manifolds
9. Examples

1. INTRODUCTION

A complex flag manifold is a compact complex manifold $\mathfrak{M}$ that is homogeneous for the action of a semisimple complex Lie group $\hat{\mathbf{G}}$. The orbits M in $\mathfrak{M}$ of a real form $\mathbf{G}$ of $\hat{\mathbf{G}}$ inherit from the complex structure of $\mathfrak{M}$ a $\mathbf{G}$ -homogeneous CR structure. In this way we obtain a large class of CR manifolds, that we call *parabolic CR manifolds*. They are homogeneous for the CR action of a real semisimple Lie group. Special examples are the compact standard homogeneous CR manifolds,

2000 *Mathematics Subject Classification*. Primary: 32V05; Secondary: 14M15, 17B20, 22E15, 32M10, 32V40, 53C30.

Key words and phrases. Complex flag manifold, homogeneous CR manifold, orbit of a real form, parabolic CR algebra.

corresponding to Levi-Tanaka algebras (see e.g. [MN1], [MN2], [MN3]), and the symmetric CR manifolds in [LN].

In [AMN] we investigated the CR structure of all minimal orbits. Here we extend our consideration to *all* orbits M of $\mathbf{G}$ in $\mathfrak{M}$. Several CR and topological invariants of M are encoded in its *cross-marked diagrams*. These are obtained from the Dynkin diagrams of the root system $\mathcal{R}$ of the complex Lie algebra $\hat{\mathfrak{g}}$ of $\hat{\mathbf{G}}$ with respect to the complexification of a Cartan subalgebra $\mathfrak{h}$ of the Lie algebra $\mathfrak{g}$ of $\mathbf{G}$. At variance with the systematic use of cross-marked Satake diagrams in the minimal orbit case, here we need more general descriptions, involving general Cartan subalgebras $\mathfrak{h}$ of the Lie algebra $\mathfrak{g}$ of $\mathbf{G}$ and different choices of simple systems of roots in $\mathcal{R}$.

The orbits of $\mathbf{G}$ in $\mathfrak{M}$ were already considered in [W]. Among the more recent contributions to the study of this subject, we cite [Ka] in the context of infinite dimensional representation theory, [GM], [HW] and [KZ] for applications to the geometry of symmetric spaces.

In our work we stress the point of view of CR geometry. Beyond its independent interest, it also provides a helpful perspective to investigate their topology. The open orbits have been extensively studied (see e.g. [FHW]). In particular, they are all simply connected (see [W]). Also the topology of the real flag manifolds has been thoroughly investigated (see e.g. [CS], [DKV], [Wi]). In our paper we show that every parabolic CR manifold M is the total space of a canonical fibration over a real flag manifold. The fiber may be disconnected and each connected component is a simply connected complex manifold, which can be retracted onto an open orbit. This essentially reduces the computation of the fundamental group of M to counting the connected components of the fiber.

The paper is organized as follows.

In §2 we review the notion of CR algebra from [MN4], [AMN], that was also recently utilized in [Fe] and [FeK]. We collect here the main results and fix the notation that will be employed in the rest of the paper, also correcting a mistake, related to the notion of *ideal nondegeneracy* in [MN4], that was pointed out to us by G.Fels.

In §3 we quickly rehearse parabolic complex Lie subalgebras and complex flag manifolds.

In §4 we begin the study of the CR algebras that are associated to the real orbits M in the complex flag manifolds $\mathfrak{M}$, also investigating the canonical $\mathbf{G}$ -equivariant maps of [MN4] in this special situation.

§5 is the core of our investigation of the CR properties of M . Through the introduction of adapted Cartan subalgebras and S- and V-fit Weyl chambers, we associate to M special systems of simple roots. Weak (i.e. holomorphic according to [BER]) nondegeneracy and fundamentality (i.e. finite type according to [BIG]) are proved to be equivalent to properties of these systems of simple roots. These can be checked from the pattern of some *cross marked diagrams* associated to M , that generalize those of [AMN], [MN2], [LN].

In §6 we turn to the construction of homogeneous CR manifolds that fiber over our orbit M and that are useful both for finding the S- and V-fit Weyl chambers and for investigating the topological properties of M in the following sections. In particular, we construct the *weakest CR model* of M , that is a step to build a chain of fibrations, with simply connected fibers, that in some instances coincides with, and in general can be considered as a substitute of, the *holomorphic arc components*

of [W].

In §7 we investigate the connectivity of the isotropy subgroup of M . This is needed to study the connectivity of the fibers of a fibration of M over a real flag manifold M' , that we utilize to compute the fundamental group of M . This is a somehow delicate point: the simply connected fibers of our construction may be not connected (the statement of [W, Theorem 8.15(1b)], that would imply connectedness, is not correct). We use classical results from [BT1], [BT2] to characterize Cartan subgroups and isotropy subgroups of connected semisimple real linear groups in terms of characters.

Finally, in §8 we discuss the fundamental group of M .

In the last section (§9) of our paper we provide several examples which show how effective our results are for the study of CR and topological properties of the orbits.

2. HOMOGENEOUS CR MANIFOLDS AND CR ALGEBRAS

A CR manifold of type (n, k) is a triple (M, HM, J) , consisting of a smooth paracompact manifold M of real dimension $(2n+k)$, a smooth real vector subbundle HM of rank $2n$ of its real tangent bundle TM , and a smooth complex structure $J: HM \rightarrow HM$ on the fibers of HM . It is also required that J satisfies the *formal integrability conditions*: $[\mathcal{C}^\infty(M, T^{0,1}M), \mathcal{C}^\infty(M, T^{0,1}M)] \subset \mathcal{C}^\infty(M, T^{0,1}M)$. Here $T^{0,1}M = \{X + iJX \mid X \in HM\}$ is the complex subbundle of the complexification $\mathbb{C}HM$ of HM corresponding to the eigenvalue $-i$ of J ; we have $T^{1,0}M \cap T^{0,1}M = 0$ and $T^{1,0}M \oplus T^{0,1}M = \mathbb{C}HM$, where $T^{1,0}M = \overline{T^{0,1}M}$. When $k = 0$, we recover the abstract definition of a *complex* manifold, via the Newlander-Nirenberg theorem.

If (M, HM, J) and (M', HM', J') are CR manifolds, a smooth $f: M \rightarrow M'$ is a CR map if: (i) $df(HM) \subset HM'$, (ii) $df \circ J = J' \circ df$ on HM . Assume that (M', HM', J') is a CR manifold and $f: M \rightarrow M'$ a smooth immersion. For $x \in M$, we set :

$$H_x M = [df(x)]^{-1} ([df(T_x M) \cap H_{f(x)} M'] \cap [J'(df(T_x M) \cap H_{f(x)} M')]) \quad \forall x \in M$$

and

$$J_x(v) = [df(x)]^{-1} (J'([df(x)](v))) \quad \forall v \in H_x M, \forall x \in M.$$

If the dimension $H_x M$ is a constant integer, independent of $x \in M$, then the disjoint union HM of the $H_x M$'s, and the map $J: HM \rightarrow HM$, equal to J_x on the fiber $H_x M$, define a CR manifold (M, HM, J) . This is the CR structure on M with the maximal CR dimension among those for which f is a CR map. In this case the map $f: M \rightarrow M'$ is called a CR immersion. If (M', HM', J') is of type (n', k') and (M, HM, J) of type (n, k) , we always have $n+k \leq n'+k'$. The immersion is *generic* when the equality $n+k = n'+k'$ holds. When $f: M \rightarrow M'$ is also an embedding, we say that f is a CR embedding or a *generic CR embedding*, respectively.

A CR map $f: (M, HM, J) \rightarrow (M', HM', J')$ is a CR submersion if $f: M \rightarrow M'$ is a smooth submersion and moreover $df(x)(H_x M) = H_{f(x)} M'$ for all $x \in M$. If (M, HM, J) is of type (n, k) and (M', HM', J') of type (n', k') , the existence of a CR submersion implies that $n \geq n'$ and $k \geq k'$.

When $f: M \rightarrow M'$ is also a smooth fiber bundle, we say that $f: (M, HM, J) \rightarrow (M', HM', J')$ is a CR fibration. The fibers are CR manifolds of type $(n-n', k-k')$.

A *CR diffeomorphism* of (M, HM, J) onto (M', HM', J') is a diffeomorphism $f: M \rightarrow M'$ such that both f and f^{-1} are smooth *CR* maps. The set of all *CR* diffeomorphisms of (M, HM, J) onto itself (*CR automorphisms*) is a group with the composition operation.

We say that (M, HM, J) is a *homogeneous CR manifold* if there is a Lie group of *CR* automorphisms that acts transitively on M .

Let (M, HM, J) be a *CR* manifold. A vector field $X \in \mathcal{C}^\infty(U, TM)$, defined on an open subset U of M , is an *infinitesimal CR automorphism* if the maps $\varphi_X(t)$ of the local 1-parameter group of local transformations generated by X are *CR*. This is equivalent to the fact that $[X, \mathcal{C}^\infty(U, T^{0,1}M)] \subset \mathcal{C}^\infty(U, T^{0,1}M)$. We say that $(M, HM, J; \mathbf{o})$ is a *locally homogeneous CR manifold* at a point $\mathbf{o} \in M$ if, for each $v \in T_{\mathbf{o}}M$, there is an infinitesimal *CR* automorphism X , defined in an open neighborhood U of $\mathbf{o}$ in M , with $v = X(\mathbf{o})$.

Homogeneous *CR* manifolds are locally *CR* homogeneous: a homogeneous *CR* manifold (M, HM, J) has a *real analytic CR* structure and therefore (see e.g. [AF]) admits a generic embedding $M \hookrightarrow \hat{M}$ into a complex manifold $\hat{M}$. Then the Lie algebra $\mathfrak{g}$ of the Lie group $\mathbf{G}$ that acts transitively by *CR* automorphisms on M can be identified to a Lie algebra of infinitesimal analytic *CR* automorphisms defined on $U = M$. Each $X^* \in \mathcal{C}^\infty(M, TM)$, corresponding to an $X \in \mathfrak{g}$, is the restriction of the real part of a holomorphic vector field Z^* , defined on an open complex neighborhood $\hat{U}$ of M in $\hat{M}$ (i.e. $X^* = [\operatorname{Re} Z^*]|_M$; see e.g. [BER §12.4]).

The germs of infinitesimal *CR* automorphisms of (M, HM, J) at a point $\mathbf{o} \in M$, with the Lie bracket, form a real Lie algebra $\mathfrak{G} = \mathfrak{G}(M, HM, J; \mathbf{o})$. We consider its complexification $\hat{\mathfrak{G}} = \mathfrak{G} \oplus i\mathfrak{G}$ and denote by $\mathfrak{Q} = \mathfrak{Q}(M, HM, J; \mathbf{o})$ the complex Lie subalgebra of $\hat{\mathfrak{G}}$ consisting of all $Z \in \hat{\mathfrak{G}}$ with $Z(\mathbf{o}) \in T_{\mathbf{o}}^{0,1}M$. The fact that $\mathfrak{Q}$ is actually a complex Lie subalgebra of $\hat{\mathfrak{G}}$ is a consequence of the formal integrability of the partial complex structure J .

When (M, HM, J) is locally *CR* homogeneous at $\mathbf{o} \in M$, the pair $(\mathfrak{G}, \mathfrak{Q}) = (\mathfrak{G}(M, HM, J; \mathbf{o}), \mathfrak{Q}(M, HM, J; \mathbf{o}))$ completely determines the *germ* of the *CR* manifold (M, HM, J) at $\mathbf{o}$. Vice versa, if $\mathfrak{g}$ is a *finite dimensional* real Lie algebra and $\mathfrak{q}$ a complex Lie subalgebra of its complexification $\hat{\mathfrak{g}}$, the general construction¹ of a germ $(M, \mathbf{o})$ of homogeneous manifold associated to the Lie algebra $\mathfrak{g}$ and to its Lie subalgebra $\mathfrak{g}_+ = \mathfrak{g} \cap \mathfrak{q}$ (cf. e.g. [Po, Ch.VII]) yields a unique, modulo local *CR* diffeomorphisms, germ of locally homogeneous *CR* manifold $(M, HM, J; \mathbf{o})$ at $\mathbf{o} \in M$, such that the complexification $\hat{\iota}$ of the correspondence $\iota: \mathfrak{g} \rightarrow \mathcal{C}_{\mathbf{o}}^\infty(M, TM)$ yields a homomorphism of complex Lie algebras: $\hat{\iota}: \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{G}}(M, HM, J; \mathbf{o})$ with

$$(2.1) \quad \iota(\mathfrak{g}) \subset \mathfrak{G}(M, HM, J; \mathbf{o}), \quad \hat{\iota}(\mathfrak{q}) \subset \mathfrak{Q}(M, HM, J; \mathbf{o})$$

and for which the induced map on the quotients

$$\mathfrak{g}/(\mathfrak{g} \cap \mathfrak{q}) \rightarrow \mathfrak{G}(M, HM, J; \mathbf{o})/(\mathfrak{G}(M, HM, J; \mathbf{o}) \cap \mathfrak{Q}(M, HM, J; \mathbf{o}))$$

is an isomorphism. In this case we say that the germ of (M, HM, J) at $\mathbf{o}$ is *associated* to the pair $(\mathfrak{g}, \mathfrak{q})$.

¹If $\mathfrak{g}$ is a finite dimensional real Lie algebra and $\mathfrak{g}_+$ a real Lie subalgebra of $\mathfrak{g}$, we can find a germ $(M, \mathbf{o})$ of analytic real manifold, unique modulo germs of analytic diffeomorphisms, for which there is a real Lie algebras homomorphism $\iota: \mathfrak{g} \rightarrow \mathcal{C}_{\mathbf{o}}^\infty(M, TM)$ with $\{\iota(X)(\mathbf{o}) \mid X \in \mathfrak{g}\} = T_{\mathbf{o}}M$ and $\mathfrak{g}_+ = \{X \in \mathfrak{g} \mid \iota(X)(\mathbf{o}) = 0\}$.

These remarks led to the introduction in [MN4] of the abstract notion of a *CR algebra*. A *CR algebra* $(\mathfrak{g}, \mathfrak{q})$ is the pair consisting of a *real* Lie algebra $\mathfrak{g}$ and of a *complex* Lie subalgebra $\mathfrak{q}$ of its complexification $\hat{\mathfrak{g}}$, such that the quotient $\mathfrak{g}/(\mathfrak{g} \cap \mathfrak{q})$ is a finite dimensional real linear space. Note that we do not require that $\mathfrak{g}$ is finite dimensional. The intersection $\mathfrak{g}_+ = \mathfrak{g} \cap \mathfrak{q}$ is the *isotropy* of $(\mathfrak{g}, \mathfrak{q})$. Let $\mathcal{H}_+ = \{\operatorname{Re} Z \mid Z \in \mathfrak{q}\}$ and denote by $\bar{Z}$ the conjugate of $Z \in \hat{\mathfrak{g}}$ with respect to the real form $\mathfrak{g}$.

A *CR algebra* $(\mathfrak{g}, \mathfrak{q})$ is :

totally real if $\mathcal{H}_+ = \mathfrak{g}_+$,

totally complex if $\mathcal{H}_+ = \mathfrak{g}$,

fundamental if $\mathcal{H}_+$ generates $\mathfrak{g}$ as a real Lie algebra,

transitive, or effective if $\mathfrak{g}_+$ does not contain any nonzero ideal of $\mathfrak{g}$,

ideal nondegenerate if all ideals of $\mathfrak{g}$ contained in $\mathcal{H}_+$ are contained in $\mathfrak{g}_+$,

weakly nondegenerate if there is no complex Lie subalgebra $\mathfrak{q}'$ of $\hat{\mathfrak{g}}$ with :

$$\mathfrak{q} \subsetneq \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}},$$

strictly nondegenerate if $\mathfrak{g}_+ = \{X \in \mathcal{H}_+ \mid [X, \mathcal{H}_+] \subset \mathcal{H}_+\}$.

Clearly :

$$\text{strictly nondegenerate} \implies \text{weakly nondegenerate} \implies \text{ideal nondegenerate}.$$

Fundamentality of $(\mathfrak{g}, \mathfrak{q})$ is equivalent to the fact that the associated germ of homogeneous *CR manifold* $(M, HM, J; \mathbf{o})$ is of *finite type in the sense of* [BlG], i.e. that the smallest involutive distribution of tangent vectors containing HM also contains $T_{\mathbf{o}}M$.

Strict and weak nondegeneracy hold, or do not hold, for all *CR algebras* that are associated to the same germ $(M, HM, J; \mathbf{o})$ of locally homogeneous *CR manifold*. They correspond indeed to the nondegeneracy of the (vector valued) Levi form and of its higher order analog, respectively (see e.g. [MN4, §13]). In particular, *weak nondegeneracy* at a point $\mathbf{o} \in M$ of a (germ of) *CR manifold* (M, HM, J) means that, for every $L \in \mathcal{C}^\infty(M, T^{1,0}M)$ with $L(\mathbf{o}) \neq 0$, there exist finitely many vector fields $\bar{Z}_1, \dots, \bar{Z}_m \in \mathcal{C}^\infty(M, T^{0,1}M)$ such that $[L, \bar{Z}_1, \dots, \bar{Z}_m](\mathbf{o}) \notin T_{\mathbf{o}}^{1,0}M \oplus T_{\mathbf{o}}^{0,1}M$.

When $(\mathfrak{g}, \mathfrak{q})$ defines at $\mathbf{o}$ a germ of homogeneous *CR manifold*, the two notions of weak nondegeneracy, the one for *CR algebras* and the one above for *CR manifolds*, coincide and also coincide with the *holomorphic nondegeneracy* of [BER].

We have the following (see [AMN, Proposition 4.1]) :

Proposition 2.1. *Let (M, HM, J) and (M', HM', J') be CR manifolds. Assume that M' is locally embeddable and that there exists a CR fibration $(M, HM, J) \xrightarrow{\pi} (M', HM', J')$, with totally complex fibers of positive dimension. Then M is weakly degenerate. $\square$*

Differently, both ideal degenerate and ideal nondegenerate *CR algebras* may correspond to the same (weakly degenerate) germ of locally homogeneous *CR manifold*.

From [MN4, Theorem 9.1] we know that *if $(\mathfrak{g}, \mathfrak{q})$ is fundamental, effective, and ideal nondegenerate, then $\mathfrak{g}$ is finite dimensional*.

From this result we deduce the following :

Theorem 2.2. *Let $(\mathfrak{g}, \mathfrak{q})$ be a fundamental effective CR algebra. Then there exist a germ of homogeneous complex manifold $(\hat{M}, \mathfrak{o})$ at a point $\mathfrak{o}$, and a germ of homogeneous generic CR submanifold $(M, HM, J; \mathfrak{o})$ of $(\hat{M}, \mathfrak{o})$ at $\mathfrak{o}$, with associated CR algebra $(\mathfrak{g}, \mathfrak{q})$.*

Proof. First we note that the statement holds true when $\mathfrak{g}$ is finite dimensional: by [Po, Ch.VII] there is a germ of homogeneous complex manifold $(\hat{M}, \mathfrak{o})$ at $\mathfrak{o}$ corresponding to the complex Lie algebra $\hat{\mathfrak{g}}$ and to its complex Lie subalgebra $\mathfrak{q}$; the inclusion $\mathfrak{g} \hookrightarrow \hat{\mathfrak{g}}$ yields the embedding into $(\hat{M}, \mathfrak{o})$ of a germ of homogeneous CR manifold $(M, HM, J; \mathfrak{o})$ at $\mathfrak{o}$, corresponding to the pair $(\mathfrak{g}, \mathfrak{q})$.

Consider now the general case. We keep the notation introduced above. By [MN4, Lemma 7.2] there is a largest ideal $\mathfrak{a}$ of $\mathfrak{g}$ contained in $\mathcal{H}_+$. By [MN4, Theorem 9.1], $\mathfrak{g}/\mathfrak{a}$ is finite dimensional and by the first part of the proof there is a germ of complex homogeneous manifold $(\hat{N}, \mathfrak{o})$ at $\mathfrak{o}$, and a germ of generic CR submanifold $(N, HN, J_N; \mathfrak{o})$ of $\hat{N}$ at $\mathfrak{o}$, associated to the pair $(\mathfrak{g}/\mathfrak{a}, \mathfrak{q}/(\mathfrak{q} \cap \hat{\mathfrak{a}}))$ (here $\hat{\mathfrak{a}} = \mathbb{C} \otimes_{\mathbb{R}} \mathfrak{a}$ is the complexification of $\mathfrak{a}$ in $\hat{\mathfrak{g}}$). If $2d$ is the real dimension of $\mathfrak{a}/(\mathfrak{a} \cap \mathfrak{g}_+)$, we can take $\hat{M} = \hat{N} \times \mathbb{C}^d$ and, likewise, $M = N \times \mathbb{C}^d$, with $HM = HN \times T(\mathbb{C}^d)$ and $J = J_N \times J_{\mathbb{C}^d}$. Then $(M, HM, J; (\mathfrak{o}, 0))$ is associated to $(\mathfrak{g}, \mathfrak{q})$. $\square$

Note that the ideal nondegeneracy of $(\mathfrak{g}, \mathfrak{q})$ implies that *all ideals $\mathfrak{r}$ of the complex Lie algebra $\hat{\mathfrak{g}}$ that are contained in $\mathfrak{q}$ are contained in $\mathfrak{q} \cap \bar{\mathfrak{q}}$* . Indeed, if $\mathfrak{r}$ is a (complex) ideal of $\hat{\mathfrak{g}}$ contained in $\mathfrak{q}$, then $\mathfrak{a} = (\mathfrak{r} + \bar{\mathfrak{r}}) \cap \mathfrak{g}$ is an ideal of $\mathfrak{g}$ contained in $\mathcal{H}_+ = (\mathfrak{q} + \bar{\mathfrak{q}}) \cap \mathfrak{g}$, and $\mathfrak{a} \subset \mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$ implies that $\mathfrak{r} + \bar{\mathfrak{r}} \subset \mathfrak{q} \cap \bar{\mathfrak{q}}$.

Let $(\mathfrak{g}, \mathfrak{q})$ be a CR algebra with a finite dimensional $\mathfrak{g}$. We denote by $\hat{\mathbf{G}}$ the connected and simply connected complex Lie group with Lie algebra $\hat{\mathfrak{g}}$ and by $\mathbf{Q}$ its analytic Lie subgroup, generated by $\mathfrak{q}$. Let $\mathbf{G}$ be the analytic subgroup of $\hat{\mathbf{G}}$ with Lie algebra $\mathfrak{g}$ and set $\mathbf{G}_+ = \mathbf{G}_+(\mathfrak{g}, \mathfrak{q}) := \mathbf{Q} \cap \mathbf{G}$. This is a Lie subgroup of $\mathbf{G}$ with Lie subalgebra $\mathfrak{g}_+$. Denote by $\tilde{\mathbf{G}}$ a connected and simply connected Lie group with Lie algebra $\mathfrak{g}$ and by $\tilde{\mathbf{G}}_+ = \tilde{\mathbf{G}}_+(\mathfrak{g}, \mathfrak{q})$ its analytic Lie subgroup with Lie subalgebra $\mathfrak{g}_+$. Since $\hat{\mathbf{G}}$ is simply connected, the conjugation σ of $\hat{\mathfrak{g}}$ with respect its real form $\mathfrak{g}$ defines an antiholomorphic involution, still denoted by σ , in $\hat{\mathbf{G}}$. Thus $\mathbf{G}$, being the connected component of the identity in the set of fixed points of σ , is closed in $\hat{\mathbf{G}}$. We have the implications:

$$\mathbf{Q} \text{ closed in } \hat{\mathbf{G}} \implies \mathbf{G}_+ \text{ closed in } \mathbf{G}, \quad \mathbf{G}_+ \text{ closed in } \mathbf{G} \implies \tilde{\mathbf{G}}_+ \text{ closed in } \tilde{\mathbf{G}}.$$

When $\tilde{\mathbf{G}}_+$ is closed in $\tilde{\mathbf{G}}$, we can uniquely define a $\tilde{\mathbf{G}}$ -homogeneous CR manifold $\tilde{M}(\mathfrak{g}, \mathfrak{q}) = (\tilde{M}, H\tilde{M}, \tilde{J})$, where the underlying smooth manifold $\tilde{M}$ is the $\tilde{\mathbf{G}}$ -homogeneous space $\tilde{\mathbf{G}}/\tilde{\mathbf{G}}_+$, and $(\mathfrak{g}, \mathfrak{q})$ is associated to the germ $(\tilde{M}, H\tilde{M}, \tilde{J}; \mathfrak{o})$ at the base point $\mathfrak{o} = e \tilde{\mathbf{G}}_+$.

Likewise, for a closed $\mathbf{G}_+ \subset \mathbf{G}$, we define the $\mathbf{G}$ -homogeneous CR manifold $M(\mathfrak{g}, \mathfrak{q}) = (M, HM, J)$ with $M = \mathbf{G}/\mathbf{G}_+$ and $(\mathfrak{g}, \mathfrak{q})$ associated to $(M, HM, J; \mathfrak{o})$ for $\mathfrak{o} = e \mathbf{G}_+$.

If $\mathbf{Q}$ is closed, $\hat{M} = \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q}) := \hat{\mathbf{G}}/\mathbf{Q}$ is a $\hat{\mathbf{G}}$ -homogeneous complex manifold and $M(\mathfrak{g}, \mathfrak{q})$ can be identified, its partial complex structure being that of a generic CR submanifold of $\hat{M}$, to the orbit of $\mathbf{G}$ through the base point $\mathfrak{o} = e \mathbf{Q}$ of $\hat{M}$.

Our canonical choice of $M(\mathfrak{g}, \mathfrak{q})$ aims to obtain a homogeneous CR manifold with a *good* embedding in a homogeneous complex manifold $\hat{M} = \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q})$.

A morphism of CR algebras $(\mathfrak{g}, \mathfrak{q}) \xrightarrow{\varphi} (\mathfrak{g}', \mathfrak{q}')$ is a homomorphism of real Lie

algebras $\varphi : \mathfrak{g} \rightarrow \mathfrak{g}'$, with $\hat{\varphi}(\mathfrak{q}) \subset \mathfrak{q}'$. (We shall consistently use "hat" to indicate complexification: e.g. $\hat{\varphi} : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}'$ is the complexification of $\varphi : \mathfrak{g} \rightarrow \mathfrak{g}'$). It is called:

- a *CR immersion* if the quotient map $[\varphi] : \mathfrak{g}/\mathfrak{g}_+ \rightarrow \mathfrak{g}'/\mathfrak{g}'_+$ is injective and $\hat{\varphi}^{-1}(\mathfrak{q}') = \mathfrak{q}$;
- a *CR submersion* if both $[\varphi] : \mathfrak{g}/\mathfrak{g}_+ \rightarrow \mathfrak{g}'/\mathfrak{g}'_+$ and $[\hat{\varphi}] : \mathfrak{q}/\hat{\mathfrak{g}}_+ \rightarrow \mathfrak{q}'/\hat{\mathfrak{g}}'_+$ are onto;
- a *local CR isomorphism* if both $[\varphi] : \mathfrak{g}/\mathfrak{g}_+ \rightarrow \mathfrak{g}'/\mathfrak{g}'_+$ and $[\hat{\varphi}] : \mathfrak{q}/\hat{\mathfrak{g}}_+ \rightarrow \mathfrak{q}'/\hat{\mathfrak{g}}'_+$ are isomorphisms;
- a *CR isomorphism* if φ is an isomorphism of CR algebras with $\hat{\varphi}(\mathfrak{q}) = \mathfrak{q}'$.

We quote from [MN4]:

Proposition 2.3. *Let $(\mathfrak{g}, \mathfrak{q}) \xrightarrow{\varphi} (\mathfrak{g}', \mathfrak{q}')$ be a morphism of CR algebras, with $\mathfrak{g}$ and $\mathfrak{g}'$ finite dimensional. Let (M, HM, J) and (M', HM', J') be the germs of homogeneous CR manifolds at $\mathfrak{o} \in M$, $\mathfrak{o}' \in M'$, associated to $(\mathfrak{g}, \mathfrak{q})$, $(\mathfrak{g}', \mathfrak{q}')$, respectively. Then there is a unique germ of smooth CR map $\Phi : (M, HM, J) \rightarrow (M', HM', J')$ with $\Phi(\mathfrak{o}) = \mathfrak{o}'$ such that $d\Phi_{\mathfrak{o}}(\iota(X)) = \iota'(\varphi(X))$. Here ι, ι' are the homomorphisms of Lie algebras $\iota : \mathfrak{g} \rightarrow \mathfrak{G}(M, HM, J)$ and $\iota' : \mathfrak{g}' \rightarrow \mathfrak{G}(M', HM', J')$ of (2.1).*

The germ Φ of smooth CR map is a CR immersion, submersion, diffeomorphism if and only if the corresponding morphism φ of CR algebras is a CR immersion, a CR submersion, a local CR isomorphism, respectively.

Let $\tilde{\mathbf{G}}$ and $\tilde{\mathbf{G}}'$ be the connected and simply connected real Lie groups with Lie algebras $\mathfrak{g}$ and $\mathfrak{g}'$, respectively. If the analytic subgroup $\tilde{\mathbf{G}}_+$ of $\tilde{\mathbf{G}}$ with Lie algebra $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$ and the analytic subgroup $\tilde{\mathbf{G}}'_+$ of $\tilde{\mathbf{G}}'$ with Lie algebra $\mathfrak{g}'_+ = \mathfrak{q}' \cap \mathfrak{g}'$ are both closed, then there is a unique smooth CR map $\tilde{\Phi} : \tilde{M}(\mathfrak{g}, \mathfrak{q}) \rightarrow \tilde{M}(\mathfrak{g}', \mathfrak{q}')$ that makes the following diagram commutes:

$$\begin{array}{ccccc} \mathfrak{g} & \xrightarrow{\exp} & \tilde{\mathbf{G}} & \longrightarrow & \tilde{M}(\mathfrak{g}, \mathfrak{q}) \\ \varphi \downarrow & & \downarrow & & \downarrow \tilde{\Phi} \\ \mathfrak{g}' & \xrightarrow{\exp} & \tilde{\mathbf{G}}' & \longrightarrow & \tilde{M}(\mathfrak{g}', \mathfrak{q}') \end{array}$$

The map $\tilde{\Phi}$ is a CR immersion, a CR submersion or a local CR diffeomorphism if and only if the corresponding CR morphism of CR algebras φ is a CR immersion, a CR submersion or a local CR isomorphism, respectively.

Let $\hat{\mathbf{G}}$ and $\hat{\mathbf{G}}'$ be the connected and simply connected complex Lie groups with Lie algebras $\hat{\mathfrak{g}}$ and $\hat{\mathfrak{g}}'$, respectively. Let $\mathbf{G}, \mathbf{Q} \subset \hat{\mathbf{G}}$ and $\mathbf{G}', \mathbf{Q}' \subset \hat{\mathbf{G}}'$ be the analytic subgroups with Lie algebras $\mathfrak{g}, \mathfrak{q}$ and $\mathfrak{g}', \mathfrak{q}'$, respectively. If $\mathbf{G}_+ = \mathbf{Q} \cap \mathbf{G}$ and $\mathbf{G}' = \mathbf{Q}' \cap \mathbf{G}'$ are closed, then there is a unique smooth CR map $\Phi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}', \mathfrak{q}')$ such that the diagram:

$$\begin{array}{ccc} \tilde{M}(\mathfrak{g}, \mathfrak{q}) & \xrightarrow{\tilde{\Phi}} & \tilde{M}(\mathfrak{g}', \mathfrak{q}') \\ \downarrow & & \downarrow \\ M(\mathfrak{g}, \mathfrak{q}) & \xrightarrow{\Phi} & M(\mathfrak{g}', \mathfrak{q}') \end{array}$$

where the vertical arrows are the natural projections from the universal coverings, commutes.

The map Φ is a CR immersion, a CR submersion or a local CR diffeomorphism if and only if the corresponding morphism of CR algebras φ is a CR immersion, a CR submersion or a local CR isomorphism, respectively.

If $\mathbf{Q} \subset \hat{\mathbf{G}}$ and $\mathbf{Q}' \subset \hat{\mathbf{G}}'$ are closed, the map $M(\mathfrak{g}, \mathfrak{q}) \xrightarrow{\Phi} M(\mathfrak{g}', \mathfrak{q}')$ is the restriction of the holomorphic map $\hat{\Phi} : \hat{M} = \hat{\mathbf{G}}/\mathbf{Q} \rightarrow \hat{M}' = \hat{\mathbf{G}}'/\mathbf{Q}'$ defined by the commutative diagram :

$$\begin{array}{ccccc} \hat{\mathfrak{g}} & \xrightarrow{\exp} & \hat{\mathbf{G}} & \longrightarrow & \hat{M} \\ \hat{\varphi} \downarrow & & \downarrow & & \downarrow \hat{\Phi} \\ \hat{\mathfrak{g}}' & \xrightarrow{\exp} & \hat{\mathbf{G}}' & \longrightarrow & \hat{M}' \end{array}$$

where the central vertical arrow is the homomorphism of complex connected simply connected Lie algebras defined by the homomorphism $\hat{\varphi} : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}'$ of their Lie algebras. $\square$

To discuss, later on, the structure of the fibers of some *CR* fibrations, we briefly introduce the notion of *semidirect sum* of *CR* algebras.

Let $(\mathfrak{g}_1, \mathfrak{q}_1)$, $(\mathfrak{g}_2, \mathfrak{q}_2)$ be *CR* algebras, and assume that $\mathfrak{g}_2$ has a structure of $\mathfrak{g}_1$ -module and that $\mathfrak{q}_2$ is a $\mathfrak{q}_1$ -module for the restriction of the complexification of the action of $\mathfrak{g}_1$ on $\mathfrak{g}_2$. Then $\mathfrak{q} = \mathfrak{q}_1 \rtimes \mathfrak{q}_2$ (semidirect sum) is a complex Lie subalgebra of the complexification of the semidirect sum $\mathfrak{g} = \mathfrak{g}_1 \rtimes \mathfrak{g}_2$, and the *CR* algebra $(\mathfrak{g}, \mathfrak{q}) = (\mathfrak{g}_1 \rtimes \mathfrak{g}_2, \mathfrak{q}_1 \rtimes \mathfrak{q}_2)$ is called the *semidirect sum* of the *CR* algebras $(\mathfrak{g}_1, \mathfrak{q}_1)$ and $(\mathfrak{g}_2, \mathfrak{q}_2)$:

$$(2.2) \quad (\mathfrak{g}, \mathfrak{q}) = (\mathfrak{g}_1 \rtimes \mathfrak{g}_2, \mathfrak{q}_1 \rtimes \mathfrak{q}_2) = (\mathfrak{g}_1, \mathfrak{q}_1) \rtimes (\mathfrak{g}_2, \mathfrak{q}_2) \quad (\text{semidirect sum}).$$

We shall assume that $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are finite dimensional. Denote by :

$\hat{\mathbf{G}}, \hat{\mathbf{G}}_1, \hat{\mathbf{G}}_2$ the connected and simply connected complex Lie groups with Lie algebras $\hat{\mathfrak{g}}, \hat{\mathfrak{g}}_1, \hat{\mathfrak{g}}_2$, respectively;

$\mathbf{G}, \mathbf{G}_1, \mathbf{G}_2$ the analytic real subgroups of the corresponding complex connected Lie groups $\hat{\mathbf{G}}, \hat{\mathbf{G}}_1, \hat{\mathbf{G}}_2$, with Lie algebras $\mathfrak{g}, \mathfrak{g}_1, \mathfrak{g}_2$, respectively;

$\mathbf{Q} \subset \hat{\mathbf{G}}, \mathbf{Q}_1 \subset \hat{\mathbf{G}}_1, \mathbf{Q}_2 \subset \hat{\mathbf{G}}_2$ the Lie subgroups corresponding to the Lie subalgebras $\mathfrak{q} \subset \hat{\mathfrak{g}}, \mathfrak{q}_1 \subset \hat{\mathfrak{g}}_1, \mathfrak{q}_2 \subset \hat{\mathfrak{g}}_2$, respectively;

$\tilde{\mathbf{G}}, \tilde{\mathbf{G}}_1, \tilde{\mathbf{G}}_2$ connected and simply connected real Lie groups with Lie algebras $\mathfrak{g}, \mathfrak{g}_1, \mathfrak{g}_2$, respectively;

$$\mathbf{G}_+ = \mathbf{Q} \cap \mathbf{G}, \mathbf{G}_{1+} = \mathbf{Q} \cap \mathbf{G}_1, \mathbf{G}_{2+} = \mathbf{Q} \cap \mathbf{G}_2;$$

$\tilde{\mathbf{G}}_+ \subset \tilde{\mathbf{G}}, \tilde{\mathbf{G}}_{1+} \subset \tilde{\mathbf{G}}_1, \tilde{\mathbf{G}}_{2+} \subset \tilde{\mathbf{G}}_2$ the analytic subgroups corresponding to the Lie subalgebras $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}, \mathfrak{g}_{1+} = \mathfrak{q}_1 \cap \mathfrak{g}_1, \mathfrak{g}_{2+} = \mathfrak{q}_2 \cap \mathfrak{g}_2$, respectively.

Let $(M, HM, J; \mathbf{o}), (M_1, HM_1, J_1; \mathbf{o}_1), (M_2, HM_2, J_2; \mathbf{o}_2)$ be the germs of locally homogeneous *CR* manifolds associated to the *CR* algebras $(\mathfrak{g}, \mathfrak{q}), (\mathfrak{g}_1, \mathfrak{q}_1), (\mathfrak{g}_2, \mathfrak{q}_2)$, respectively.

We obtain :

Theorem 2.4. *The diffeomorphism $\mathbf{G}_1 \times \mathbf{G}_2 \ni (g_1, g_2) \rightarrow g_1 g_2 \in \mathbf{G}_1 \rtimes \mathbf{G}_2$ defines a germ of *CR* diffeomorphism :*

$$(M_1, HM_1, J_1; \mathbf{o}_1) \times (M_2, HM_2, J_2; \mathbf{o}_2) \rightarrow (M, HM, J; \mathbf{o}).$$

If $\tilde{\mathbf{G}}_{1+}$ and $\tilde{\mathbf{G}}_{2+}$ are closed, then $\tilde{\mathbf{G}}_+$ is closed and we obtain a global *CR* diffeomorphism :

$$\tilde{M}(\mathfrak{g}_1, \mathfrak{q}_1) \times \tilde{M}(\mathfrak{g}_2, \mathfrak{q}_2) \rightarrow \tilde{M}(\mathfrak{g}, \mathfrak{q}).$$

If $\mathbf{G}_{1+}$ and $\mathbf{G}_{2+}$ are closed, then $\mathbf{G}_+$ is closed and we obtain a global CR diffeomorphism :

$$(*) \quad M(\mathfrak{g}_1, \mathfrak{q}_1) \times M(\mathfrak{g}_2, \mathfrak{q}_2) \rightarrow M(\mathfrak{g}, \mathfrak{q}) .$$

When $\mathbf{Q}_1$ and $\mathbf{Q}_2$ are closed, also $\mathbf{Q}$ is closed and the map $(*)$ is the restriction of a biholomorphic map

$$\left(\hat{\mathbf{G}}_1 / \mathbf{Q}_1 \right) \times \left(\hat{\mathbf{G}}_2 / \mathbf{Q}_2 \right) \rightarrow \hat{\mathbf{G}} / \mathbf{Q} . \quad \square$$

Let $\mathfrak{g}$ be a real Lie algebra and $\mathfrak{q}, \mathfrak{q}'$ complex subalgebras of its complexification $\hat{\mathfrak{g}}$, with $\mathfrak{q} \subset \mathfrak{q}'$. Then the identity map in $\mathfrak{g}$ and the inclusion $\mathfrak{q} \hookrightarrow \mathfrak{q}'$ define a $\mathfrak{g}$ -equivariant CR submersion of CR algebras

$$(2.3) \quad (\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}') .$$

If $(M, HM, J; \mathbf{o})$ and $(M', HM', J'; \mathbf{o}')$ are germs of locally homogeneous CR manifolds with associated CR algebras $(\mathfrak{g}, \mathfrak{q})$ and $(\mathfrak{g}, \mathfrak{q}')$, respectively, then the identity map $\mathfrak{g} \rightarrow \mathfrak{g}$ defines, by passing to the quotients, the differential of a CR map $\pi_{(\mathbf{o})} : (M, HM, J; \mathbf{o}) \rightarrow (M', HM', J'; \mathbf{o}')$ that is locally $\mathbf{G}$ -equivariant for a (connected) real Lie group with Lie algebra $\mathfrak{g}$.

Let $\mathbf{G}$ be a connected Lie group with Lie algebra $\mathfrak{g}$ and assume that there are two $\mathbf{G}$ -homogeneous CR manifolds (M, HM, J) , (M', HM', J') that are associated to $(\mathfrak{g}, \mathfrak{q})$ at some $\mathbf{o} \in M$ and to $(\mathfrak{g}, \mathfrak{q}')$ at some $\mathbf{o}' \in M'$, respectively. Then there is a unique $\mathbf{G}$ -equivariant CR map $\pi : (M, HM, J) \rightarrow (M', HM', J')$ with $\pi(\mathbf{o}) = \mathbf{o}'$.

In general $\pi_{(\mathbf{o})}$ (and π , when defined) are smooth, but not CR , $\mathbf{G}$ -equivariant fibrations: a necessary and sufficient condition for (2.3) to be a $\mathfrak{g}$ -equivariant CR fibration, and hence for $\pi_{(\mathbf{o})}$ (and π , when defined) to be $\mathbf{G}$ -equivariant local (resp. global) fibrations is that (see [MN4, Lemma 5.1])

$$(2.4) \quad \mathfrak{q}' = \mathfrak{q} + \hat{\mathfrak{g}}_+ .$$

We call $(\mathfrak{g}, \mathfrak{q}')$ the *basis* of the fibration (2.3). The *fiber* of (2.3) is the CR algebra $(\mathfrak{g}'_+, \mathfrak{q}'')$ where $\mathfrak{q}'' = \hat{\mathfrak{g}}'_+ \cap \mathfrak{q}$. It is a CR algebra associated to the germ $(F, HF, J|_{HF}; \mathbf{o})$, where $(F; \mathbf{o}) = \left(\pi_{(\mathbf{o})}^{-1}(\mathbf{o}'); \mathbf{o} \right)$, and the germ of partial complex structure $(HF, J|_{HF}; \mathbf{o})$ that is characterized by requiring that the smooth embedding $(\pi^{-1}(\mathbf{o}'); \mathbf{o}) \hookrightarrow (M; \mathbf{o})$ is a CR immersion.

We know that (2.3) is always a CR fibration, with a totally complex fiber, when $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$: indeed in this case $\mathfrak{q}' = \mathfrak{q} + \hat{\mathfrak{g}}_+$.

From [MN4, §5,] we have :

Proposition 2.5. *Let $(\mathfrak{g}, \mathfrak{q})$ be a CR algebra. Then there exist :*

- a largest ideal $\mathfrak{i}$ of $\mathfrak{g}$ with $\mathfrak{i} \subset \mathfrak{g}_+$*
- a largest ideal $\mathfrak{a}$ of $\mathfrak{g}$ with $\mathfrak{a} \subset \mathcal{H}_+$*
- a largest complex Lie subalgebra $\mathfrak{q}'$ of $\hat{\mathfrak{g}}$ with $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$*
- a smallest complex Lie subalgebra $\mathfrak{q}''$ of $\hat{\mathfrak{g}}$ with $\mathfrak{q} + \bar{\mathfrak{q}} \subset \mathfrak{q}''$.*

We have $\mathfrak{i} \subset \mathfrak{a} \subset \mathfrak{q}' \subset \mathfrak{q}''$ and $\mathfrak{q}'' = \bar{\mathfrak{q}}'' = \hat{\mathfrak{g}}''$ for a real Lie subalgebra $\mathfrak{g}''$ of $\mathfrak{g}$.

The identity in $\mathfrak{g}$ defines $\mathfrak{g}$ -equivariant CR fibrations $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}') \rightarrow (\mathfrak{g}, \mathfrak{q}'')$, where $(\mathfrak{g}, \mathfrak{q}')$ is weakly nondegenerate and $(\mathfrak{g}, \mathfrak{q}'')$ is totally real. For all complex Lie subalgebras $\mathfrak{f}$ of $\hat{\mathfrak{g}}$ with $\mathfrak{q} \subset \mathfrak{f} \subsetneq \mathfrak{q}'$, the CR algebra $(\mathfrak{g}, \mathfrak{f})$ is weakly degenerate. For all complex Lie subalgebras $\mathfrak{f}$ with $\mathfrak{q} \subset \mathfrak{f} \subset \mathfrak{q}'$, the $\mathfrak{g}$ -equivariant map $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{f})$ is a CR fibration with a totally complex fiber.

The CR algebra $(\mathfrak{g}'', \mathfrak{q})$ is fundamental and, for all real Lie subalgebras $\mathfrak{l}$ of $\mathfrak{g}$ with $\mathfrak{g}'' \subsetneq \mathfrak{l} \subset \mathfrak{g}$ the CR algebra $(\mathfrak{l}, \mathfrak{q})$ is not fundamental. $\square$

3. PARABOLIC SUBALGEBRAS AND COMPLEX FLAG MANIFOLDS

In this section we collect the notions on complex parabolic subalgebras and fix that notation that will be utilized throughout the paper. This is mostly a review of classical results, for which general references are [B1, Ch.IV, §2.6, Ch.VI, §1], [B2, Ch.VIII, §3], [Kn, Ch.VII] [Wa, Ch.1], [W].

Let $\hat{\mathfrak{g}}$ be a complex Lie algebra. A maximal solvable complex Lie subalgebra $\mathfrak{b}$ of $\hat{\mathfrak{g}}$ is called a *Borel*, or *minimal parabolic* complex Lie subalgebra of $\hat{\mathfrak{g}}$. A complex Lie subalgebra $\mathfrak{q}$ of $\hat{\mathfrak{g}}$ is *parabolic* if it contains a complex Borel subalgebra $\mathfrak{b}$ of $\hat{\mathfrak{g}}$.

For our purposes, it will be sufficient to consider the case of a semisimple $\hat{\mathfrak{g}}$. Thus from now on we shall assume that $\hat{\mathfrak{g}}$ is a semisimple complex Lie algebra.

A parabolic subalgebra $\mathfrak{q}$ of $\hat{\mathfrak{g}}$ contains a complex Cartan subalgebra $\hat{\mathfrak{h}}$ of $\hat{\mathfrak{g}}$. Let $\mathcal{R} = \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}})$ be the root system of $\hat{\mathfrak{g}}$ with respect to $\hat{\mathfrak{h}}$. We denote by $\mathfrak{h}_{\mathbb{R}}$ the real form of $\hat{\mathfrak{h}}$ on which all roots are real valued. Thus $\mathcal{R}$ is a subset of the real dual space $\mathfrak{h}_{\mathbb{R}}^*$. The Killing form $\kappa_{\hat{\mathfrak{g}}}$ of $\hat{\mathfrak{g}}$ restricts to a real positive scalar product in $\mathfrak{h}_{\mathbb{R}}$. We shall write $(A|B) = \kappa_{\hat{\mathfrak{g}}}(A, B)$ for $A, B \in \mathfrak{h}_{\mathbb{R}}$. We set also $(\xi|\eta) = (T_{\xi}|T_{\eta})$ for $\xi, \eta \in \mathfrak{h}_{\mathbb{R}}^*$ and $(T_{\xi}|A) = \xi(A)$, $(T_{\eta}|A) = \eta(A)$ for all $A \in \mathfrak{h}_{\mathbb{R}}$ (dual scalar product in $\mathfrak{h}_{\mathbb{R}}^*$). Roots $\alpha, \beta \in \mathcal{R}$ for which $\alpha \pm \beta \notin \mathcal{R}$ are called *strongly orthogonal*. Note that strongly orthogonal roots are also orthogonal for the scalar product in $\mathfrak{h}_{\mathbb{R}}^*$.

An element $H \in \hat{\mathfrak{h}}$ is *regular* if $\alpha(H) \neq 0$ for all $\alpha \in \mathcal{R}$. Denote by $\mathfrak{C}(\mathcal{R})$ the set of the Weyl chambers of $\mathcal{R}$. They are the connected components of the set of regular elements of $\mathfrak{h}_{\mathbb{R}}$. For $C \in \mathfrak{C}(\mathcal{R})$, and $H \in C$, the set $\mathcal{R}^+(C) = \{\alpha \in \mathcal{R} \mid \alpha(H) > 0\}$ is independent of the choice of $H \in C$: it is called the set of *positive roots* with respect to C . The set $\mathcal{R}^-(C) = \mathcal{R}^+(C^{\text{opp}})$, for $C^{\text{opp}} = \{-H \mid H \in C\}$, is the complement of $\mathcal{R}^+(C)$ in $\mathcal{R}$ and is called the set of *negative roots* with respect to C . A Weyl chamber C also defines a partial order relation " $\prec_C$ " in $\mathfrak{h}_{\mathbb{R}}^*$, by:

$$\eta \prec_C \xi \text{ if } \eta(A) < \xi(A) \text{ for all } A \in C.$$

Thus $\mathcal{R}^+(C) = \{\alpha \in \mathcal{R} \mid \alpha \succ_C 0\}$.

With $\hat{\mathfrak{g}}^{\alpha} = \{X \in \hat{\mathfrak{g}} \mid [H, X] = \alpha(H) X \forall H \in \hat{\mathfrak{h}}\}$, we set

$$(3.1) \quad \mathcal{Q} = \{\alpha \in \mathcal{R} \mid \hat{\mathfrak{g}}^{\alpha} \subset \mathfrak{q}\}.$$

This is the *parabolic set* associated to $\mathfrak{q}$ and $\hat{\mathfrak{h}}$. Parabolic sets of roots are abstractly defined by the two conditions:

- (i) $\alpha, \beta \in \mathcal{Q}, \alpha + \beta \in \mathcal{R} \implies \alpha + \beta \in \mathcal{Q}$ (closedness)
- (ii) $\mathcal{Q} \cup \mathcal{Q}^{\text{opp}} = \mathcal{R}$ where $\mathcal{Q}^{\text{opp}} = \{-\alpha \mid \alpha \in \mathcal{Q}\}$.

Condition (ii) is equivalent to the fact that $\mathcal{Q} \supset \mathcal{R}^+(C)$ for some Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$. We have

$$(3.2) \quad \mathfrak{q} = \mathfrak{q}_{\mathcal{Q}} = \hat{\mathfrak{h}} \oplus \sum_{\alpha \in \mathcal{Q}} \hat{\mathfrak{g}}^{\alpha}$$

and the correspondence $\mathcal{Q} \longleftrightarrow \mathfrak{q}_{\mathcal{Q}}$ is one-to-one between parabolic subsets of $\mathcal{R}$ and parabolic subalgebras of $\hat{\mathfrak{g}}$ containing $\hat{\mathfrak{h}}$.

Given a parabolic set $\mathcal{Q} \subset \mathcal{R}$ we set :

$$(3.3) \quad \mathcal{Q}^r = \{\alpha \in \mathcal{Q} \mid -\alpha \in \mathcal{Q}\} \quad \text{and} \quad \mathcal{Q}^n = \mathcal{Q} \setminus \mathcal{Q}^r = \{\alpha \in \mathcal{Q} \mid -\alpha \notin \mathcal{Q}\}.$$

Then

$$(3.4) \quad \mathfrak{q}^n = \sum_{\alpha \in \mathcal{Q}^n} \hat{\mathfrak{g}}^\alpha$$

is the *nilradical* of $\mathfrak{q}$, i.e. the set of the elements Z of its radical $\mathfrak{r}(\mathfrak{q})$ for which $\text{ad}_{\hat{\mathfrak{g}}}(Z)$ is nilpotent, and

$$(3.5) \quad \mathfrak{q}^r = \hat{\mathfrak{h}} \oplus \sum_{\alpha \in \mathcal{Q}^r} \hat{\mathfrak{g}}^\alpha$$

a reductive complement of $\mathfrak{q}^n$ in $\mathfrak{q}$. The complex parabolic Lie subalgebra $\mathfrak{q}$ of $\hat{\mathfrak{g}}$ is its own normalizer and the normalizer of its nilradical $\mathfrak{q}^n$:

$$(3.6) \quad \mathfrak{q} = \{Z \in \hat{\mathfrak{g}} \mid [Z, \mathfrak{q}] \subset \mathfrak{q}\} = \{Z \in \hat{\mathfrak{g}} \mid [Z, \mathfrak{q}^n] \subset \mathfrak{q}^n\}.$$

If $A \in \mathfrak{h}_{\mathbb{R}}$, then the set

$$(3.7) \quad \mathcal{Q}_A = \{\alpha \in \mathcal{R} \mid \alpha(A) \geq 0\}$$

is parabolic, with $\mathcal{Q}_A^r = \{\alpha \in \mathcal{R} \mid \alpha(A) = 0\}$ and $\mathcal{Q}_A^n = \{\alpha \in \mathcal{R} \mid \alpha(A) > 0\}$. Vice versa, if $\mathcal{Q}$ is parabolic, set $\delta = \sum \{\alpha \mid \alpha \in \mathcal{Q}^n\}$, and define $T_\delta \in \mathfrak{h}_{\mathbb{R}}$ by $(T_\delta | A) = \delta(A)$ for all $A \in \mathfrak{h}_{\mathbb{R}}$. Then $\mathcal{Q} = \mathcal{Q}_{T_\delta} = \{\alpha \in \mathcal{R} \mid (\alpha | \delta) \geq 0\}$. The set of $A \in \mathfrak{h}_{\mathbb{R}}$ for which $\mathcal{Q} = \mathcal{Q}_A$ is in fact a relatively open convex cone in $\mathfrak{h}_{\mathbb{R}}$.

When $\mathcal{Q} = \mathcal{Q}_A$ for some $A \in \mathfrak{h}_{\mathbb{R}}$, we shall also write $\mathfrak{q}_A$ for $\mathfrak{q}_{\mathcal{Q}_A}$.

The sets $\mathcal{Q}^n$ associated to parabolic $\mathcal{Q}$'s are called *horocyclic* (see [Wa, §1.1]). The correspondence $\mathcal{Q}^n \longleftrightarrow \mathfrak{q}^n = \sum_{\alpha \in \mathcal{Q}^n} \hat{\mathfrak{g}}^\alpha$ is one-to-one between horocyclic sets of roots in $\mathcal{R}$ and nilradicals of complex parabolic Lie subalgebras containing $\hat{\mathfrak{h}}$.

Given a parabolic subset $\mathcal{Q} \subset \mathcal{R}$, we use the notation $\mathcal{Q}^{-n}$ for its *opposite horocyclic set* $[\mathcal{Q}^n]^{\text{opp}} = \mathcal{R} \setminus \mathcal{Q}$: the corresponding nilpotent ideal $\mathfrak{q}^{-n} = \sum_{\alpha \in \mathcal{Q}^{-n}} \hat{\mathfrak{g}}^\alpha$ is a complement of $\mathfrak{q}$ in $\hat{\mathfrak{g}}$.

To a parabolic $\mathcal{Q} \subset \mathcal{R}$ we associate the set of Weyl chambers :

$$(3.8) \quad \mathfrak{C}(\mathcal{R}, \mathcal{Q}) = \{C \in \mathfrak{C}(\mathcal{R}) \mid \mathcal{R}^+(C) \subset \mathcal{Q}\} = \{C \in \mathfrak{C}(\mathcal{R}) \mid \mathcal{R}^+(C) \supset \mathcal{Q}^n\}.$$

We denote by $\mathcal{B}(C)$ the simple roots of $\mathcal{R}^+(C)$, for $C \in \mathfrak{C}(\mathcal{R})$. Every $\alpha \in \mathcal{R}$ can be written in a unique way as a linear combination, with integral coefficients (either all ≥ 0 or all ≤ 0), of the simple roots in $\mathcal{B}(C)$:

$$(3.9) \quad \alpha = \sum_{\beta \in \mathcal{B}(C)} k_\alpha^\beta(C) \beta.$$

We set

$$(3.10) \quad \text{supp}_C(\alpha) = \{\beta \in \mathcal{B}(C) \mid k_\alpha^\beta(C) \neq 0\}.$$

If $\mathcal{Q}$ is parabolic, $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, and $\Phi_C(\mathcal{Q}) = \mathcal{B}(C) \cap \mathcal{Q}^n$, then

$$(3.11) \quad \mathcal{Q}^n = \{\alpha \in \mathcal{R}^+(C) \mid \text{supp}_C(\alpha) \cap \Phi_C(\mathcal{Q}) \neq \emptyset\}.$$

The correspondence

$$(3.12) \quad \mathcal{B}(C) \supset \Phi_C \longleftrightarrow \mathfrak{q} = \hat{\mathfrak{h}} \oplus \sum_{\alpha \in \mathcal{R}^+(C)} \hat{\mathfrak{g}}^\alpha \oplus \sum_{\substack{\alpha \in \mathcal{R}^-(C) \\ \text{supp}_C(\alpha) \cap \Phi_C = \emptyset}} \hat{\mathfrak{g}}^\alpha$$

is one-to-one between subsets Φ_C of $\mathcal{B}(C)$ and complex parabolic Lie subalgebra of $\hat{\mathfrak{g}}$ that contain $\hat{\mathfrak{h}}$ and have an associated parabolic set $\mathcal{Q}$ with $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$.

Having fixed a Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$ and $\Phi_C \subset \mathcal{B}(C)$, we shall denote by $\mathfrak{q}_{\Phi_C}$ the complex parabolic Lie subalgebra of $\hat{\mathfrak{g}}$ defined by the right hand side of (3.12) and by $\mathcal{Q}_{\Phi_C}$ the corresponding parabolic set.

We denote by $\mathbf{W}(\mathcal{R})$ the Weyl group of $\mathcal{R}$, (i.e. the group of isometries of $\mathfrak{h}_{\mathbb{R}}^*$ generated by the symmetries $\xi \rightarrow s_\alpha(\xi) = \xi - 2[(\xi|\alpha)/\|\alpha\|^2]\alpha$ for $\alpha \in \mathcal{R}$) and by $\mathbf{A}(\mathcal{R})$ the group of all isometries of $\mathfrak{h}_{\mathbb{R}}^*$ (with respect to the scalar product defined above) that transform $\mathcal{R}$ into itself. For $C \in \mathfrak{C}(\mathcal{R})$ we denote by $\mathbf{A}_C(\mathcal{R})$ the subgroup of $w \in \mathbf{A}(\mathcal{R})$ for which $w(\mathcal{R}^+(C)) = \mathcal{R}^+(C)$. Then $\mathbf{A}(\mathcal{R}) = \mathbf{A}_C(\mathcal{R}) \rtimes \mathbf{W}(\mathcal{R})$.

We define $\mathbf{W}(\mathcal{R}, \mathcal{Q})$ and $\mathbf{A}(\mathcal{R}, \mathcal{Q})$ as the subgroups of $\mathbf{W}(\mathcal{R})$ and $\mathbf{A}(\mathcal{R})$, respectively, that transform $\mathcal{Q}$ into itself. Then we have Chevalley's Lemma (see e.g. [Wa, Theorem 1.1.2.8]):

Lemma 3.1. *The group $\mathbf{W}(\mathcal{R}, \mathcal{Q})$ is generated by the symmetries s_α with $\alpha \in \mathcal{Q}^r$. If $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, then the symmetries s_α with $\alpha \in \mathcal{B}(C) \setminus \Phi_C(\mathcal{Q})$ generate $\mathbf{W}(\mathcal{R}, \mathcal{Q})$ and $\mathbf{A}(\mathcal{R}, \mathcal{Q})$ is a semidirect product $\mathbf{A}(\mathcal{R}, \mathcal{Q}) = \mathbf{A}_C(\mathcal{R}, \mathcal{Q}) \rtimes \mathbf{W}(\mathcal{R}, \mathcal{Q})$, with $\mathbf{A}_C(\mathcal{R}, \mathcal{Q}) = \mathbf{A}_C(\mathcal{R}) \cap \mathbf{A}(\mathcal{R}, \mathcal{Q})$. $\square$*

Let $\hat{\mathbf{G}}$ be a connected complex Lie group with Lie algebra $\hat{\mathfrak{g}}$. If $\mathbf{Q}$ is any Lie subgroup of $\hat{\mathbf{G}}$ with complex Lie subalgebra $\mathfrak{q}$ that is parabolic in $\hat{\mathfrak{g}}$, then $\mathbf{Q}$ is closed, connected and coincides with its normalizer in $\hat{\mathbf{G}}$ and is the normalizer of its Lie algebra for the adjoint representation:

$$(3.13) \quad \mathbf{Q} = \{g \in \hat{\mathbf{G}} \mid g \mathbf{Q} g^{-1} = \mathbf{Q}\} = \{g \in \hat{\mathbf{G}} \mid \text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{q}) = \mathfrak{q}\}$$

The homogeneous space $\mathfrak{M} = \hat{\mathbf{G}}/\mathbf{Q}$ is compact and simply connected. Since the center $\mathbf{Z}(\hat{\mathbf{G}})$ of $\hat{\mathbf{G}}$ is contained in all its parabolic subgroups, the choice of different connected complex Lie groups $\hat{\mathbf{G}}$ yields the same $\mathfrak{M}$. Hence we can consider the complex flag manifold $\mathfrak{M} = \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q})$ as an object associated simply to the pair $(\hat{\mathfrak{g}}, \mathfrak{q})$. We also recall (see [W, 2.7]) that the integral cohomology $H^*(\mathfrak{M}, \mathbb{Z})$ is torsion free and 0 in odd degrees.

If $\hat{\mathfrak{h}}$ is a Cartan subalgebra of $\hat{\mathfrak{g}}$ contained in $\mathfrak{q}$ and $\mathcal{Q} \subset \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}})$ the parabolic set of roots associated to $\mathfrak{q}$, the complex dimension of $\mathfrak{M} = \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q})$ equals the number of roots in $\mathcal{Q}^n$.

4. PARABOLIC CR ALGEBRAS AND PARABOLIC CR MANIFOLDS

A CR algebra $(\mathfrak{g}, \mathfrak{q})$ is called *parabolic* if $\mathfrak{g}$ is finite dimensional, and $\mathfrak{q}$ is a *parabolic* subalgebra of its complexification $\hat{\mathfrak{g}}$.

By the results quoted at the end of §3, all the homogeneous spaces $\tilde{M} = \tilde{M}(\mathfrak{g}, \mathfrak{q})$, $M = M(\mathfrak{g}, \mathfrak{q})$, and $\mathfrak{M} = \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q})$ of §2 are well defined and M is an orbit, in the complex flag manifold $\mathfrak{M}$ of a connected complex semisimple Lie group $\hat{\mathbf{G}}$, of a connected real form $\mathbf{G}$ of $\hat{\mathbf{G}}$. We say that M is a *parabolic CR manifold*.

Vice versa, if $\mathbf{G}$ is a connected real form of the complex semisimple Lie group $\hat{\mathbf{G}}$, then all $\mathbf{G}$ -orbits in the complex flag manifolds $\mathfrak{M} = \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q})$ are homogeneous CR manifolds of the form $M = M(\mathfrak{g}, \mathfrak{q}')$, for some parabolic complex Lie subalgebra $\mathfrak{q}'$ of $\hat{\mathfrak{g}}$, conjugated to $\mathfrak{q}$ by an inner automorphism.

It is worth noticing that, in the definition of the homogeneous CR manifold $M(\mathfrak{g}, \mathfrak{q}) = \mathbf{G}/\mathbf{G}_+$, we can define the isotropy $\mathbf{G}_+ = \mathbf{G}_+(\mathfrak{g}, \mathfrak{q})$ by

$$(4.1) \quad \mathbf{G}_+ = \{g \in \mathbf{G} \mid \text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{q}) = \mathfrak{q}\}.$$

Since the center of $\mathbf{G}$ is always contained in $\mathbf{G}_+$, we obtain an equivalent definition of $M(\mathfrak{g}, \mathfrak{q})$ if we substitute to $\hat{\mathbf{G}}$ any connected complex Lie group $\hat{\mathbf{G}}'$ with the same Lie algebra $\hat{\mathfrak{g}}$ and to $\mathbf{G}$ the analytic subgroup $\mathbf{G}'$ of $\hat{\mathbf{G}}'$ with Lie algebra $\mathfrak{g}$. However, it is more convenient to fix a simply connected $\hat{\mathbf{G}}$, since in this case, by [BT2, Corollaire 4.7], we have:

$$(4.2) \quad \mathbf{G} = \hat{\mathbf{G}}^\sigma = \{g \in \hat{\mathbf{G}} \mid \sigma(g) = g\},$$

where $\sigma : \hat{\mathbf{G}} \rightarrow \hat{\mathbf{G}}$ is the anti-holomorphic involution of $\hat{\mathbf{G}}$ corresponding to the conjugation σ of $\hat{\mathfrak{g}}$ defined by the real form $\mathfrak{g}$.

We begin by recalling some general facts about parabolic CR algebras, and their associated CR manifolds, that were explained in [AMN, §5].

Proposition 4.1. *A parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ is effective if and only if: (i) $\mathfrak{g}$ is semisimple, (ii) no simple ideal of $\hat{\mathfrak{g}}$ is contained in $\mathfrak{q} \cap \bar{\mathfrak{q}}$.*

An effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ with $\mathfrak{g}$ simple is either totally complex or ideal nondegenerate. $\square$

Proposition 4.2. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and let $\mathfrak{g} = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_\ell$ be the decomposition of $\mathfrak{g}$ into the direct sum of its simple ideals. Then:*

- (i) $\mathfrak{q} = \mathfrak{q}_1 \oplus \cdots \oplus \mathfrak{q}_\ell$ where $\mathfrak{q}_j = \mathfrak{q} \cap \hat{\mathfrak{g}}_j$ for $j = 1, \dots, \ell$;
- (ii) for each $j = 1, \dots, \ell$, $(\mathfrak{g}_j, \mathfrak{q}_j)$ is an effective parabolic CR algebra;
- (iii) $(\mathfrak{g}, \mathfrak{q})$ is ideal (resp. weakly, strictly) nondegenerate if and only if for each $j = 1, \dots, \ell$, the CR algebra $(\mathfrak{g}_j, \mathfrak{q}_j)$ is ideal (resp. weakly, strictly) nondegenerate;
- (iv) $(\mathfrak{g}, \mathfrak{q})$ is fundamental if and only if for each $j = 1, \dots, \ell$, the CR algebra $(\mathfrak{g}_j, \mathfrak{q}_j)$ is fundamental.
- (v) We have ($\cong$ meaning biholomorphic or CR equivalence):

$$\begin{aligned} \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q}) &\cong \hat{M}(\hat{\mathfrak{g}}_1, \mathfrak{q}_1) \times \cdots \times \hat{M}(\hat{\mathfrak{g}}_\ell, \mathfrak{q}_\ell), \\ \tilde{M}(\mathfrak{g}, \mathfrak{q}) &\cong \tilde{M}(\mathfrak{g}_1, \mathfrak{q}_1) \times \cdots \times \tilde{M}(\mathfrak{g}_\ell, \mathfrak{q}_\ell), \\ M(\mathfrak{g}, \mathfrak{q}) &\cong M(\mathfrak{g}_1, \mathfrak{q}_1) \times \cdots \times M(\mathfrak{g}_\ell, \mathfrak{q}_\ell). \end{aligned} \quad \square$$

When $\mathfrak{q}$ is parabolic in $\hat{\mathfrak{g}}$, its conjugate $\bar{\mathfrak{q}}$ with respect to the real form $\mathfrak{g}$ is also parabolic in $\hat{\mathfrak{g}}$. Therefore the intersection $\mathfrak{q} \cap \bar{\mathfrak{q}}$ contains a Cartan subalgebra $\hat{\mathfrak{h}}$ that is invariant under conjugation. The intersection $\mathfrak{h} = \hat{\mathfrak{h}} \cap \mathfrak{g}$ is a Cartan subalgebra of $\mathfrak{g}$, contained in $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$.

A Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ contained in $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$ is said to be *adapted* to $(\mathfrak{g}, \mathfrak{q})$. We also recall [AMN, Proposition 5.4]:

Proposition 4.3. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra, with isotropy subalgebra $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$. The elements A of the radical $\mathfrak{r}(\mathfrak{g}_+)$ of $\mathfrak{g}_+$ for which $\text{ad}_{\mathfrak{g}}(A) : \mathfrak{g} \rightarrow \mathfrak{g}$ is nilpotent, form a nilpotent ideal $\mathfrak{n}$ of $\mathfrak{g}_+$. It admits a reductive complement $\mathfrak{g}_0$ in $\mathfrak{g}_+$:*

$$(4.3) \quad \mathfrak{g}_+ = \mathfrak{n} \oplus \mathfrak{g}_0.$$

The reductive subalgebra $\mathfrak{g}_0$ is uniquely determined modulo inner automorphisms of $\mathfrak{g}_+$ from the subgroup generated by those of the form $\exp(\text{ad}_{\mathfrak{g}_+}(X))$ with $X \in \mathfrak{n}$. $\square$

Let $\mathfrak{z}_0$ be the center and $\mathfrak{s}_0 = [\mathfrak{g}_0, \mathfrak{g}_0]$ the semisimple ideal of $\mathfrak{g}_0$. Then

$$(4.4) \quad \mathfrak{g}_0 = \mathfrak{z}_0 \oplus \mathfrak{s}_0.$$

Thus, a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}_+$ of $\mathfrak{g}$ can be taken as the direct sum

$$(4.5) \quad \mathfrak{h} = \mathfrak{z}_0 \oplus \mathfrak{h}_0$$

of the center $\mathfrak{z}_0$ of $\mathfrak{g}_0$ and a Cartan subalgebra $\mathfrak{h}_0$ of $\mathfrak{s}_0$. Vice versa, every Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$ has the form (4.5) for some reductive subalgebra $\mathfrak{g}_0$ of $\mathfrak{g}_+$.

It is also convenient to consider a Cartan decomposition (see e.g. [B2]):

$$(4.6) \quad \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$$

of $\mathfrak{g}$, corresponding to a Cartan involution ϑ . The set $\mathfrak{k} = \{X \in \mathfrak{g} \mid \vartheta(X) = X\}$ of fixed points of ϑ is a maximal compact Lie subalgebra of $\mathfrak{g}$ and $\mathfrak{p} = \{X \in \mathfrak{g} \mid \vartheta(X) = -X\}$ its orthogonal for the Killing form $\kappa_{\mathfrak{g}}$ of $\mathfrak{g}$. Any ϑ -invariant Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ decomposes into the direct sum $\mathfrak{h} = \mathfrak{h}^+ \oplus \mathfrak{h}^-$ of its *compact* (or *toroidal*) part $\mathfrak{h}^+ = \mathfrak{h} \cap \mathfrak{k} \subset \mathfrak{k}$ and its noncompact (or *vector part*) $\mathfrak{h}^- = \mathfrak{h} \cap \mathfrak{p} \subset \mathfrak{p}$.

We say that the Cartan decomposition (4.6) is *adapted* to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ if $\mathfrak{k}$ contains a maximal compact Lie subalgebra of $\mathfrak{g}_+$. Then:

Lemma 4.4. *If (4.6) is adapted to the parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, then every Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ that is adapted to $(\mathfrak{g}, \mathfrak{q})$ is conjugate, modulo an inner automorphism of $\mathfrak{g}_+$, to a ϑ -invariant Cartan subalgebra $\mathfrak{h}_0$ of $\mathfrak{g}$ that is adapted to $(\mathfrak{g}, \mathfrak{q})$.*

Vice versa, if $\mathfrak{h}$ is a Cartan subalgebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$, then there exists a Cartan decomposition (4.6), adapted to $(\mathfrak{g}, \mathfrak{q})$, such that $\mathfrak{h} = (\mathfrak{h} \cap \mathfrak{k}) \oplus (\mathfrak{h} \cap \mathfrak{p})$.

In particular, if $\{\mathfrak{q}_i \mid i \in I\}$ is a family of complex parabolic Lie subalgebras of $\hat{\mathfrak{g}}$ such that $\bigcap_{i \in I} \mathfrak{q}_i$ is parabolic in $\hat{\mathfrak{g}}$, then there exist both a Cartan decomposition (4.6) and a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$, compatible with (4.6), that are adapted to all the $(\mathfrak{g}, \mathfrak{q}_i)$'s. $\square$

We say that $(\vartheta, \mathfrak{h})$ is an *adapted Cartan pair* for $(\mathfrak{g}, \mathfrak{q})$ if:

- (i) ϑ is the Cartan involution of a Cartan decomposition (4.6) adapted to $(\mathfrak{g}, \mathfrak{q})$;
- (ii) $\mathfrak{h}$ is a ϑ -invariant Cartan subalgebra of $\mathfrak{g}$ contained in $\mathfrak{g}_+ = \mathfrak{g} \cap \mathfrak{q}$.

Being $\sigma : \hat{\mathfrak{g}} \ni X \rightarrow \bar{X} \in \hat{\mathfrak{g}}$ the conjugation in $\hat{\mathfrak{g}}$ associated to the real form $\mathfrak{g}$, and having fixed (4.6), we also consider the conjugation $\tau : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}$ of $\hat{\mathfrak{g}}$ with respect to its compact real form $\mathfrak{u} = \mathfrak{k} \oplus i\mathfrak{p}$ and use the same symbol ϑ to denote the $\mathbb{C}$ -linear extension to $\hat{\mathfrak{g}}$ of the Cartan involution ϑ of $\mathfrak{g}$. We obtain in this way three commuting involutions σ, τ, ϑ of $\hat{\mathfrak{g}}$, each being the composition product of the other two:

$$(4.7) \quad \tau = \vartheta \circ \sigma = \sigma \circ \vartheta, \quad \sigma = \vartheta \circ \tau = \tau \circ \vartheta, \quad \vartheta = \sigma \circ \tau = \tau \circ \sigma.$$

In particular $\mathfrak{u}$ is invariant under σ : $\sigma(\mathfrak{u}) = \mathfrak{u}$.

Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to $(\mathfrak{g}, \mathfrak{q})$. Then $\mathfrak{h}_{\mathbb{R}}$ decomposes as: $\mathfrak{h}_{\mathbb{R}} = \mathfrak{h}^- \oplus i\mathfrak{h}^+$ and $\mathfrak{h}_{\mathbb{R}}$ is a Cartan subalgebra of a split real form $\mathfrak{g}_{\mathbb{R}}$ of $\hat{\mathfrak{g}}$. The involutions σ, τ and ϑ transform $\mathfrak{h}_{\mathbb{R}}$ into itself. Hence, by transposition, they define involutions on $\mathfrak{h}_{\mathbb{R}}^*$, that we still denote by the same symbols σ, τ and ϑ , and that transform the set of roots $\mathcal{R}$ into itself. We set $\bar{\alpha} = \sigma(\alpha)$ for all $\alpha \in \mathfrak{h}_{\mathbb{R}}^*$. We have:

$$(4.8) \quad \tau(\alpha) = -\alpha, \quad \vartheta(\alpha) = -\bar{\alpha} \quad \forall \alpha \in \mathfrak{h}_{\mathbb{R}}^*.$$

The reductive complement $\mathfrak{q}^r$ of $\mathfrak{q}^n$ in $\mathfrak{q}$ of (3.5) is $\mathfrak{q} \cap \tau(\mathfrak{q})$, while the reductive complement $\mathfrak{g}_0$ of $\mathfrak{n}$ in $\mathfrak{g}_+$ of (4.3) can be taken equal to $\mathfrak{g}_+ \cap \vartheta(\mathfrak{g}_+)$.

A real Lie subalgebra $\mathfrak{t}$ of $\mathfrak{g}$ is *triangular* if all linear maps $\text{ad}_{\mathfrak{g}}(X) \in \mathfrak{gl}_{\mathbb{R}}(\mathfrak{g})$ with $X \in \mathfrak{t}$ can be simultaneously represented by triangular matrices in a suitable basis of $\mathfrak{g}$. All maximal triangular subalgebras of $\mathfrak{g}$ are conjugate by an inner automorphism (cf. [Mo, §5.4]). A real Lie subalgebra of $\mathfrak{g}$ containing a maximal triangular subalgebra of $\mathfrak{g}$ is called a *t-subalgebra*.

An effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ will be called *minimal* if $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$ is a *t-subalgebra* of $\mathfrak{g}$.

We observe that a maximal triangular subalgebra of $\mathfrak{g}$ contains a maximal Abelian subalgebra of semisimple elements of $\mathfrak{g}$ having real eigenvalues. We have (see [AMN, §5]):

Proposition 4.5. *An effective parabolic minimal CR algebra $(\mathfrak{g}, \mathfrak{q})$ admits an adapted Cartan pair $(\vartheta, \mathfrak{h})$ in which $\mathfrak{h}$ is a maximally noncompact Cartan subalgebra of $\mathfrak{g}$.*

Let $\mathfrak{g}$ be a semisimple real Lie algebra and $\mathfrak{q}$ a parabolic subalgebra of its complexification $\hat{\mathfrak{g}}$. Then, up to CR isomorphisms, there is a unique parabolic minimal CR algebra $(\mathfrak{g}', \mathfrak{q}')$ with $\mathfrak{g}'$ isomorphic to $\mathfrak{g}$ and $\mathfrak{q}'$ isomorphic to $\mathfrak{q}$.

A totally real effective parabolic CR algebra is minimal. $\square$

By [AMN, Theorem 8.6 and Corollary 9.2] we have:

Theorem 4.6. *Let $(\mathfrak{g}, \mathfrak{q})$ be a parabolic minimal fundamental CR algebra. Then $\tilde{M}(\mathfrak{g}, \mathfrak{q}) \cong M(\mathfrak{g}, \mathfrak{q})$ is compact and simply connected.* $\square$

Next we consider the CR fibrations of Proposition 2.5 in the special case of a parabolic CR algebra.

Theorem 4.7. *Every effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ admits a unique $\mathfrak{g}$ -equivariant CR fibration $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \hat{\mathfrak{g}}')$, where $\hat{\mathfrak{g}}' \supset \mathfrak{q}$ is the complexification of a real parabolic subalgebra $\mathfrak{g}'$ of $\mathfrak{g}$, and the fiber is fundamental. The basis $(\mathfrak{g}, \hat{\mathfrak{g}}')$ is a totally real parabolic CR algebra and also the fiber $(\mathfrak{g}', \mathfrak{q})$ is parabolic.*

This yields a $\mathbf{G}$ -equivariant CR fibration $\pi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \hat{\mathfrak{g}}')$ with compact basis. Each connected component of the fiber is CR diffeomorphic to $M(\mathfrak{g}', \mathfrak{q})$.

Proof. Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra. The complex subalgebra $\mathfrak{q}''$ generated by $\mathfrak{q} + \bar{\mathfrak{q}}$ is parabolic in $\hat{\mathfrak{g}}$ because contains $\mathfrak{q}$, and is the complexification of a real parabolic subalgebra $\mathfrak{g}'$ of $\mathfrak{g}$ because $\bar{\mathfrak{q}}'' = \mathfrak{q}''$. Then (2.3) yields a $\mathfrak{g}$ -equivariant CR fibration with a totally real basis. The fiber is $(\mathfrak{g}', \mathfrak{q})$. This is parabolic because $\mathfrak{q}$, being parabolic in $\hat{\mathfrak{g}}$, is also parabolic in $\hat{\mathfrak{g}}' \subset \hat{\mathfrak{g}}$.

The final statement follows from the commutative diagram :

$$\begin{array}{ccc} M(\mathfrak{g}, \mathfrak{q}) & \longrightarrow & \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q}) \\ \pi \downarrow & & \hat{\pi} \downarrow \\ M(\mathfrak{g}, \hat{\mathfrak{g}}') & \longrightarrow & \hat{M}(\hat{\mathfrak{g}}, \hat{\mathfrak{g}}') \end{array}$$

that yields an embedding of each fiber of $\pi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \hat{\mathfrak{g}}')$ into a fiber of $\hat{\pi} : \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q}) \rightarrow \hat{M}(\hat{\mathfrak{g}}, \hat{\mathfrak{g}}')$. The basis $M(\mathfrak{g}, \hat{\mathfrak{g}}')$ is compact because $(\mathfrak{g}, \hat{\mathfrak{g}}')$ is totally real and hence minimal. $\square$

Example 4.1. The fiber is not necessarily connected. An example of a $\mathbf{G}$ -equivariant fibration $M(\mathfrak{g}, \mathfrak{q}) \xrightarrow{\pi} M(\mathfrak{g}, \mathfrak{q}')$ of parabolic CR manifolds with a disconnected fiber is the following.

Let $\mathfrak{g} = \mathfrak{sl}(3, \mathbb{R}) \subset \hat{\mathfrak{g}} = \mathfrak{sl}(3, \mathbb{C})$, and

$$\begin{aligned} \mathfrak{q} &= \left\{ \begin{pmatrix} z_{1,1} & z_{1,2} & z_{1,3} \\ 0 & z_{2,2} & z_{2,3} \\ 0 & 0 & z_{3,3} \end{pmatrix} \in \mathfrak{sl}(3, \mathbb{C}) \right\}, \\ \mathfrak{q}' &= \left\{ \begin{pmatrix} z_{1,1} & z_{1,2} & z_{1,3} \\ 0 & z_{2,2} & z_{2,3} \\ 0 & z_{3,2} & z_{3,3} \end{pmatrix} \in \mathfrak{sl}(3, \mathbb{C}) \right\}. \end{aligned}$$

Let e_1, e_2, e_3 be the canonical basis of $\mathbb{C}^3$. We have $\mathbf{G} = \mathbf{SL}(3, \mathbb{R}) \subset \hat{\mathbf{G}} = \mathbf{SL}(3, \mathbb{C})$. The isotropy subgroup $\mathbf{G}_+ = \{g \in \mathbf{SL}(3, \mathbb{R}) \mid \text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{q}) = \mathfrak{q}\}$ of the point $\mathfrak{o} = e\mathbf{Q} \in M(\hat{\mathfrak{g}}, \mathfrak{q})$ is the stabilizer in $\mathbf{SL}(3, \mathbb{R})$ of the complete flag $\langle e_1 \rangle \subset \langle e_1, e_2 \rangle$. It has 4 connected components. Likewise, the isotropy subgroup $\mathbf{G}' = \{g \in \mathbf{SL}(3, \mathbb{R}) \mid \text{ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{q}') = \mathfrak{q}'\}$, at $\mathfrak{o}' = e\mathbf{Q}' \in M(\mathfrak{g}, \mathfrak{q}')$, is the stabilizer in $\mathbf{SL}(3, \mathbb{R})$ of the line $\langle e_1 \rangle$, and has 2 connected components. The fiber $\mathbf{G}'/\mathbf{G}_+$ is isomorphic to the stabilizer of $\langle e_1 \rangle$ in $\mathbf{SL}_{\mathbb{R}}(\mathbb{R}e_1 + \mathbb{R}e_2)$ and has 2 connected components.

Theorem 4.8. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra. Then there is a unique $\mathfrak{g}$ -equivariant CR fibration $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}')$ with a weakly nondegenerate basis $(\mathfrak{g}, \mathfrak{q}')$ and a totally complex fiber. The basis $(\mathfrak{g}, \mathfrak{q}')$ is a parabolic CR algebra.*

Proof. We recall from [MN4, §5], that $\mathfrak{q}'$ is the unique maximal subalgebra of $\hat{\mathfrak{g}}$ that contains $\mathfrak{q}$ and is contained in $\mathfrak{q} + \bar{\mathfrak{q}}$. Clearly $\mathfrak{q}'$ is parabolic because it contains the parabolic subalgebra $\mathfrak{q}$. $\square$

Next we investigate the general structure of the fiber of a $\mathbf{G}$ -equivariant submersion $M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}')$ for a pair of complex parabolic subalgebras $\mathfrak{q} \subset \mathfrak{q}'$ of $\hat{\mathfrak{g}}$.

Theorem 4.9. *Let $\mathfrak{q} \subset \mathfrak{q}'$ be complex parabolic Lie subalgebras of $\hat{\mathfrak{g}}$. With $\mathfrak{g}' = \mathfrak{g} \cap \mathfrak{q}'$, the CR algebra $(\mathfrak{g}', \bar{\mathfrak{q}}' \cap \mathfrak{q})$ is the fiber of the $\mathfrak{g}$ -equivariant fibration $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}')$ (see §2).*

The nilradical $\mathfrak{n}'$ of $\mathfrak{g}'$, consisting of the $\text{ad}_{\mathfrak{g}}$ -nilpotent elements of the radical $\mathfrak{r}(\mathfrak{g}')$ of $\mathfrak{g}'$, has a reductive complement $\mathfrak{g}'_0$ in $\mathfrak{g}'$ such that :

- (i) *The CR algebra $(\mathfrak{g}'_0, \hat{\mathfrak{g}}'_0 \cap \mathfrak{q})$ is parabolic.*
- (ii) *The fiber $(\mathfrak{g}', \hat{\mathfrak{g}}' \cap \mathfrak{q})$ is the semidirect sum of the parabolic CR algebra $(\mathfrak{g}'_0, \hat{\mathfrak{g}}'_0 \cap \mathfrak{q})$ and of the nilpotent CR algebra $(\mathfrak{n}', \hat{\mathfrak{n}}' \cap \mathfrak{q})$.*
- (iii) *The nilpotent CR algebra $(\mathfrak{n}', \hat{\mathfrak{n}}' \cap \mathfrak{q})$ is totally complex.*
- (iv) *The connected components of the fibers of the $\mathbf{G}$ -equivariant fibration $\pi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}')$ are CR diffeomorphic to the Cartesian product of a parabolic CR manifold $M(\mathfrak{g}'_0, \mathfrak{q} \cap \hat{\mathfrak{g}}'_0)$ and of a Euclidean complex manifold $(\cong \mathbb{C}^\ell)$.*

Proof. Fix a Cartan pair $(\vartheta, \mathfrak{h})$, that is adapted for both $(\mathfrak{g}, \mathfrak{q})$ and $(\mathfrak{g}, \mathfrak{q}')$. With the notation introduced above, set $\mathfrak{q}^r = \mathfrak{q} \cap \tau(\mathfrak{q})$ and $\mathfrak{q}'^r = \mathfrak{q}' \cap \tau(\mathfrak{q}')$. Since $\mathfrak{q} \subset \mathfrak{q}'$ and $\mathfrak{q}$ is parabolic, we have the inclusions: $\mathfrak{q}^r \subset \mathfrak{q}'^r$ and $\mathfrak{q}^n \supset \mathfrak{q}'^n$. The complexification of the fiber $\mathfrak{g}'$ is:

$$\hat{\mathfrak{g}}' = \mathfrak{q}' \cap \bar{\mathfrak{q}}' = \hat{\mathfrak{g}}'_0 \rtimes \hat{\mathfrak{n}}',$$

where :

$$\begin{cases} \hat{\mathfrak{g}}'_0 = \mathfrak{q}'^r \cap \bar{\mathfrak{q}}'^r, \\ \hat{\mathfrak{n}}' = (\mathfrak{q}'^r \cap \bar{\mathfrak{q}}'^n) \oplus (\mathfrak{q}'^n \cap \bar{\mathfrak{q}}'^r) \oplus (\mathfrak{q}'^n \cap \bar{\mathfrak{q}}'^n) = (\mathfrak{q}' \cap \bar{\mathfrak{q}}'^n) + (\mathfrak{q}'^n \cap \bar{\mathfrak{q}}'). \end{cases}$$

Thus $\mathfrak{g}' = \mathfrak{g}'_0 \rtimes \mathfrak{n}'$, where $\mathfrak{n}' = \hat{\mathfrak{n}}' \cap \mathfrak{g}$ is a real form of the nilradical $\hat{\mathfrak{n}}'$ of $\mathfrak{q}' \cap \bar{\mathfrak{q}}'$ and $\mathfrak{g}'_0 := \hat{\mathfrak{g}}'_0 \cap \mathfrak{g}$ a reductive complement of $\mathfrak{n}'$ in $\mathfrak{g}'$.

We have:

$$\hat{\mathfrak{g}}' \cap \mathfrak{q} = (\hat{\mathfrak{g}}'_0 \rtimes \hat{\mathfrak{n}}') \cap \mathfrak{q} = (\hat{\mathfrak{g}}'_0 \cap \mathfrak{q}) \rtimes (\hat{\mathfrak{n}}' \cap \mathfrak{q}),$$

so that :

$$(\mathfrak{g}', \hat{\mathfrak{g}}' \cap \mathfrak{q}) = (\mathfrak{g}'_0, \hat{\mathfrak{g}}'_0 \cap \mathfrak{q}) \rtimes (\mathfrak{n}', \hat{\mathfrak{n}}' \cap \mathfrak{q}).$$

The complex Lie subalgebra $\hat{\mathfrak{g}}'_0 \cap \mathfrak{q}$ is parabolic in $\hat{\mathfrak{g}}'_0$, because $\hat{\mathfrak{g}}'_0$ is reductive, $\mathfrak{q}$ is parabolic in $\hat{\mathfrak{g}}$, and $\hat{\mathfrak{g}}'_0 \cap \mathfrak{q}$ contains a Cartan subalgebra of $\hat{\mathfrak{g}}'_0$ and of $\hat{\mathfrak{g}}$.

Note that $\hat{\mathfrak{n}}' \cap \mathfrak{q}$ is contained in, but in general not equal to, the nilradical $\hat{\mathfrak{n}}$ of $\mathfrak{q} \cap \bar{\mathfrak{q}}$. We have:

$$\hat{\mathfrak{n}}' \cap \mathfrak{q} \supset \mathfrak{q}'^n \cap \bar{\mathfrak{q}}',$$

so that :

$$\hat{\mathfrak{n}}' \cap \mathfrak{q} + \overline{\hat{\mathfrak{n}}' \cap \mathfrak{q}} \supset \mathfrak{q}'^n \cap \bar{\mathfrak{q}}' + \mathfrak{q}' \cap \bar{\mathfrak{q}}'^n = \hat{\mathfrak{n}}'$$

shows that actually :

$$\hat{\mathfrak{n}}' \cap \mathfrak{q} + \overline{\hat{\mathfrak{n}}' \cap \mathfrak{q}} = \hat{\mathfrak{n}}'$$

and the nilpotent CR algebra $(\mathfrak{n}', \hat{\mathfrak{n}}' \cap \mathfrak{q})$ is totally complex.

The $\mathbf{G}$ -equivariant fibration $\pi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}')$ is the restriction of the $\hat{\mathbf{G}}$ equivariant fibration $\hat{\pi} : \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q}) \rightarrow \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q}')$. The typical fiber $\mathfrak{F}$ of $\hat{\pi}$ is $\mathbf{Q}'/\mathbf{Q}$. Since $\mathfrak{q}$ is a parabolic complex Lie subalgebra of $\mathfrak{q}'$, the fiber $\mathfrak{F}$ is a complex flag manifold and, in particular, is compact, connected and simply connected. Thus the typical fiber F of $\pi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}')$ is a submanifold of a complex flag manifold.

Denote still by $\tau : \hat{\mathbf{G}} \rightarrow \hat{\mathbf{G}}$ the involution of $\hat{\mathbf{G}}$ associated to the conjugation $\tau : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}$ with respect to the compact real form $\mathfrak{u} = \mathfrak{k} \oplus i\mathfrak{p}$. Since $\mathfrak{q}'^n \subset \mathfrak{q}^n$, the fiber $\mathfrak{F}$ can be viewed also as a flag manifold of the reductive complex closed connected Lie subgroup $\mathbf{Q}'^r = \mathbf{Q}' \cap \tau(\mathbf{Q}')$ of $\hat{\mathbf{G}}$. The fiber F is contained in the orbit $\mathfrak{F}_0 \subset \mathfrak{F}$ of the closed complex Lie subgroup $\hat{\mathbf{G}}' := \mathbf{Q}' \cap \sigma(\mathbf{Q}')$ of $\mathbf{Q}'$. The group $\hat{\mathbf{G}}'$ is connected because it contains a Cartan subgroup of $\hat{\mathbf{G}}$, and decomposes into the semidirect product

$$(4.9) \quad \hat{\mathbf{G}}' = \hat{\mathbf{G}}'_0 \rtimes \hat{\mathbf{N}}',$$

where $\hat{\mathbf{G}}'_0$ and $\hat{\mathbf{N}}'$ are the analytic complex Lie subgroups of $\hat{\mathbf{G}}$ generated by the Lie subalgebras $\hat{\mathfrak{g}}'_0$ and $\hat{\mathfrak{n}}'$, respectively. We have $\hat{\mathbf{G}}'_0 = \mathbf{Q}'^r \cap \sigma(\mathbf{Q}'^r)$, so that $\hat{\mathbf{G}}'_0$ is closed in $\hat{\mathbf{G}}$. Moreover, since $\text{ad}_{\hat{\mathfrak{g}}}(Z)$ is nilpotent for all $Z \in \hat{\mathfrak{n}}$, by Engel's theorem and the semisimplicity of $\hat{\mathfrak{g}}$, we obtain that $\exp : \hat{\mathfrak{n}}' \rightarrow \hat{\mathbf{N}}'$ is an analytic diffeomorphism, and $\hat{\mathbf{N}}'$ is Euclidean.

The validity of (iv) is then a consequence of the next Proposition. $\square$

Proposition 4.10. *Let $\hat{\mathbf{N}}'$ be a connected nilpotent complex Lie group with complex Lie algebra $\hat{\mathfrak{n}}'$ and $\mathfrak{n}'$ a real form of $\hat{\mathfrak{n}}'$. Let $\mathbf{N}'$ be the real analytic Lie subgroup of $\hat{\mathbf{N}}'$ with Lie algebra $\mathfrak{n}'$, and $\mathbf{Q}_0$ a closed connected complex Lie subgroup of $\hat{\mathbf{N}}'$, with Lie algebra $\mathfrak{q}_0 \subset \hat{\mathfrak{n}}'$, and set $\mathbf{N} = \mathbf{Q}_0 \cap \mathbf{N}'$. Assume that the CR algebra $(\mathfrak{n}', \mathfrak{q}_0)$ is totally complex. With $E = \mathbf{N}'/\mathbf{N}$ and $\hat{E} = \hat{\mathbf{N}}'/\mathbf{Q}_0$, the natural map $E \rightarrow \hat{E}$ obtained from the inclusion $\mathbf{N}' \hookrightarrow \hat{\mathbf{N}}'$ is a diffeomorphism.*

Proof. The condition that $(\mathfrak{n}', \mathfrak{q}_0)$ is totally complex is equivalent to the equality $\mathfrak{n}' + \mathfrak{q}_0 = \hat{\mathfrak{n}}'$. Since $\hat{\mathfrak{n}}'$ is nilpotent, this equality implies (see the proof below) that the map $\mathbf{N}' \times \mathbf{Q}_0 \ni (n, q) \rightarrow n \cdot q \in \hat{\mathbf{N}}'$ is onto, and hence the inclusion $\mathbf{N}' \hookrightarrow \hat{\mathbf{N}}'$ yields, by passing to the quotients, a smooth one-to-one map $f : E \rightarrow \hat{E}$. We note that $E = \mathbf{N}'/\mathbf{N}$ is a complex manifold with the homogeneous CR structure defined by the CR algebra $(\mathfrak{n}', \mathfrak{q}_0)$. With this complex structure on E , and with the complex structure that $\hat{E}$ inherits from $\hat{\mathbf{N}}'$, the map f is holomorphic. Being one-to-one, f is a biholomorphism.

We give here a simple argument to prove that $\hat{\mathbf{N}}' = \mathbf{N}'\mathbf{Q}_0$.

Consider the lower central series

$$\begin{aligned} \hat{\mathfrak{n}} &= C^{(0)}(\hat{\mathfrak{n}}') \supset C^{(1)}(\hat{\mathfrak{n}}') = [\hat{\mathfrak{n}}', \hat{\mathfrak{n}}'] \supset \dots \supset C^{(h)}(\hat{\mathfrak{n}}') = [C^{(h-1)}(\hat{\mathfrak{n}}'), \hat{\mathfrak{n}}'] \supset \dots \\ &\dots \supset C^{(m)}(\hat{\mathfrak{n}}') = [C^{(m-1)}(\hat{\mathfrak{n}}'), \hat{\mathfrak{n}}'] = \{0\}. \end{aligned}$$

Since $\hat{\mathfrak{n}}'$ is nilpotent and $\hat{\mathbf{N}}'$ is connected, the exponential map $\exp : \hat{\mathfrak{n}}' \rightarrow \hat{\mathbf{N}}'$ is surjective. Let $Z \in \hat{\mathfrak{n}}'$. We want to prove that there is $g \in \mathbf{N}'$ such that $g^{-1} \cdot \exp(Z) \in \mathbf{Q}_0$. To this aim, let $X \in \mathfrak{n}'$ and $W \in \mathfrak{q}_0$ be such that $Z = X + W$. Let $Z_1 \in \hat{\mathfrak{n}}'$ be such that $\exp(Z_1) = \exp(-X) \exp(Z) \exp(-W)$. We claim that, if $Z \in C^h(\hat{\mathfrak{n}}')$, then $Z_1 \in C^{h+1}(\hat{\mathfrak{n}}')$.

While proving this claim, we can assume that $\hat{\mathbf{N}}'$ is also simply connected, so that all Lie subgroups $\hat{\mathbf{N}}'_h = \exp(C^{(h)}(\hat{\mathfrak{n}}'))$ are closed and simply connected. For each integer $h \geq 0$ we have a commutative diagram :

$$\begin{array}{ccc} \hat{\mathfrak{n}}' & \xrightarrow{\exp} & \hat{\mathbf{N}}' \\ \pi_h \downarrow & & \downarrow p_h \\ \hat{\mathfrak{n}}'/C^{h+1}(\hat{\mathfrak{n}}') & \xrightarrow{[\exp]} & \hat{\mathbf{N}}'/\hat{\mathbf{N}}'_{h+1} \end{array}$$

where $[\exp]$ denotes the exponential map on the quotient. If $Z \in C^h(\hat{\mathfrak{n}}')$, then $\pi_h(Z)$ belongs to the center of the quotient Lie algebra $\hat{\mathfrak{n}}'/C^{h+1}(\hat{\mathfrak{n}}')$. Hence we obtain :

$$\begin{aligned} [\exp](\pi_h(Z_1)) &= [\exp](-\pi_h(X)) \cdot [\exp](\pi_h(Z)) \cdot [\exp](\pi_h(X - Z)) \\ &= [\exp](-\pi_h(X)) \cdot [\exp](\pi_h(Z) + \pi_h(X - Z)) \\ &= [\exp](-\pi_h(X)) \cdot [\exp](\pi_h(X)) = \mathbf{1}_{\hat{\mathfrak{n}}'/\hat{\mathfrak{n}}'_{h+1}}. \end{aligned}$$

Since $[\exp]$ is a diffeomorphism, we obtain that $Z \in C^{h+1}(\hat{\mathfrak{n}}')$.

We show by recurrence that for every $Z \in C^{(m-i)}(\hat{\mathfrak{n}}')$ there is some $X \in \mathfrak{n}$ such that $\exp(-X) \cdot \exp(Z) \in \mathbf{Q}_0$. This is trivially true when $m = 0$, as $Z = 0$ in this case. If $Z \in C^{(m-i)}(\hat{\mathfrak{n}}')$ for some $i > 0$, and $X \in \mathfrak{n}'$ is such that $X - Z \in \mathfrak{q}_0$, then $\exp(-X) \cdot \exp(Z) \cdot \exp(X - Z) = \exp(Z_1)$ for some $Z_1 \in C^{(m-i+1)}(\hat{\mathfrak{n}}')$. By the recursive assumption, there is $X_1 \in \mathfrak{n}'$ such that $\exp(-X_1)\exp(Z_1) \in \mathbf{Q}_0$. Then $g = \exp(X_1) \cdot \exp(X) \in \mathbf{N}'$ and $g^{-1} \cdot \exp(Z) \in \mathbf{Q}_0$. For $i = m$ we obtain our contention. $\square$

From Theorem 4.9 we obtain :

Theorem 4.11. *Let $M = M(\mathfrak{g}, \mathfrak{q})$ and $M' = M(\mathfrak{g}, \mathfrak{q}')$ be parabolic CR manifolds. If $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$, then the $\mathbf{G}$ -equivariant fibration $M \rightarrow M'$ is CR and has a totally complex simply connected fiber.*

Proof. We already noted in §2 that the CR algebra associated to the fiber F of the fibration $M \rightarrow M'$, and hence F itself, is totally complex when $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$. By Theorem 4.9 the connected components of the fiber are the product of a Euclidean complex nilmanifold and of a manifold $M(\mathfrak{g}'_0, \hat{\mathfrak{g}}'_0 \cap \mathfrak{q})$, for a totally complex parabolic CR algebra $(\mathfrak{g}'_0, \hat{\mathfrak{g}}'_0 \cap \mathfrak{q})$. This $M(\mathfrak{g}'_0, \hat{\mathfrak{g}}'_0 \cap \mathfrak{q})$ is an open orbit of a connected real form $\mathbf{G}'_0$ of a connected complex Lie group $\hat{\mathbf{G}}'_0$ with Lie algebra $\hat{\mathfrak{g}}'_0$, and thus is simply connected by [W, Theorem 5.4]. $\square$

Corollary 4.12. *Let $(\mathfrak{g}, \mathfrak{q}')$ be the weakly nondegenerate reduction of the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$. Then*

$$(4.10) \quad f : M = M(\mathfrak{g}, \mathfrak{q}) \rightarrow M' = M(\mathfrak{g}, \mathfrak{q}')$$

is a $\mathbf{G}$ -equivariant CR fibration with complex simply connected fibers. $\square$

We give here a simple general criterion that ensures the existence and the connectedness of the fiber of some $\mathbf{G}$ -equivariant fibration.

Proposition 4.13. *Let $(\mathfrak{g}, \mathfrak{q})$, $(\mathfrak{g}, \mathfrak{q}')$ be two effective parabolic CR algebras. Assume that $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g} \subset \mathfrak{g}'_+ = \mathfrak{q}' \cap \mathfrak{g}$ and that $\mathfrak{g}_+$ contains a Cartan subalgebra $\mathfrak{h}$ that is maximally noncompact among the Cartan subalgebras of $\mathfrak{g}$ that are contained in $\mathfrak{g}'_+$. Then the germ of local $\mathbf{G}$ -equivariant submersion $(M(\mathfrak{g}, \mathfrak{q}), \mathfrak{o}) \rightarrow (M(\mathfrak{g}, \mathfrak{q}'), \mathfrak{o}')$, defined by the projection $\mathfrak{g}/\mathfrak{g}_+ \rightarrow \mathfrak{g}/\mathfrak{g}'_+$, extends to a $\mathbf{G}$ -equivariant fibration $\pi : M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}')$ with connected fibers.*

Proof. Let $\mathbf{H} = \mathbf{Z}_{\mathbf{G}}(\mathfrak{h}) = \{h \in \mathbf{G} \mid \text{Ad}_{\mathfrak{g}}(h)(H) = H, \forall H \in \mathfrak{h}\}$ be the Cartan subgroup of $\mathbf{G}$ corresponding to $\mathfrak{h}$. We have $\text{Ad}_{\hat{\mathfrak{g}}}(h)(\mathfrak{q}) = \mathfrak{q}$ and $\text{Ad}_{\hat{\mathfrak{g}}}(h)(\mathfrak{q}') = \mathfrak{q}'$ for all $h \in \mathbf{H}$. Hence $\mathbf{H} \subset \mathbf{G}_+ \cap \mathbf{G}'_+$. Since $\mathfrak{h}$ is maximally noncompact in $\mathfrak{g}'_+$ and a fortiori in $\mathfrak{g}_+$, by [Kn Prop.7.90], all connected components of $\mathbf{G}'_+$ and $\mathbf{G}_+$ intersect $\mathbf{H}$. The connected component of the identity $\mathbf{G}^0_+$ of $\mathbf{G}_+$ is contained in

the connected component $[\mathbf{G}'_+]^0$ of the identity in $\mathbf{G}'_+$. Since $\mathbf{G}_+$ is generated by $\mathbf{G}_+^0$ and $\mathbf{H}$, and likewise $\mathbf{G}'_+$ is generated by $[\mathbf{G}'_+]^0$ and $\mathbf{H}$, we obtain at the same time that $\mathbf{G}_+ \subset \mathbf{G}'_+$ and that the fiber $\mathbf{G}'_+/\mathbf{G}_+$ is connected. $\square$

Using Proposition 4.13, we can prove:

Theorem 4.14. *Let $(\mathfrak{g}, \mathfrak{q})$ and $(\mathfrak{g}, \mathfrak{q}')$ be two effective parabolic CR algebras. If*

$$\mathfrak{q} \cap \bar{\mathfrak{q}} = \mathfrak{q}' \cap \bar{\mathfrak{q}}',$$

then the CR manifolds $M = M(\mathfrak{g}, \mathfrak{q})$ and $M' = M(\mathfrak{g}, \mathfrak{q}')$ are diffeomorphic, by a $\mathbf{G}$ -equivariant diffeomorphism.

Proof. Let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}$, contained in $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g} = \mathfrak{g}'_+ = \mathfrak{q}' \cap \mathfrak{g}$ and maximally noncompact as a Cartan subalgebra of $\mathfrak{g}_+$.

Let $A, A' \in \mathfrak{h}_{\mathbb{R}}$ be such that $\mathfrak{q} = \mathfrak{q}_A$, $\mathfrak{q}' = \mathfrak{q}_{A'}$. We can assume that $\mathfrak{q} \neq \mathfrak{q}'$, so that A and A' are linearly independent. Then we set $A_t = A + t(A' - A)$, for $0 \leq t \leq 1$, so that $A_0 = A$ and $A_1 = A'$. Let us take a partition $t_0 = 0 < t_1 < \dots < t_{m-1} < t_m = 1$ such that the rank of $\text{ad}_{\mathfrak{g}}(A_t)$ is constant for t in the open intervals $t_{j-1} < t < t_j$, $1 \leq j \leq m$, so that $\mathfrak{q}_{A_t} = \mathfrak{q}_{A_{t'}}$ for $t_{j-1} < t, t' < t_j$. Let $M_j = M(\mathfrak{g}, \mathfrak{q}_{A_{t_j}})$, for $0 \leq j \leq m$, and $N_j = M(\mathfrak{g}, \mathfrak{q}_{A_{(t_{j-1}+t_j)/2}})$, for $1 \leq j \leq m$. Since: $\mathfrak{q}_{A_{(t_{j-1}+t_j)/2}} \subset \mathfrak{q}_{A_{t_{j-1}}} \cap \mathfrak{q}_{A_{t_j}}$, there are $\mathbf{G}$ -equivariant maps:

$$M_{j-1} \xleftarrow{f_j} N_j \xrightarrow{F_j} M_j$$

for all $1 \leq j \leq m$. By Proposition 4.13, all these maps, being covering maps with connected fibers, are diffeomorphisms. Thus:

$$(F_m \circ f_m^{-1}) \circ (F_{m-1} \circ f_{m-1}^{-1}) \circ \dots \circ (F_1 \circ f_1^{-1}) : M \longrightarrow M'$$

is a $\mathbf{G}$ -equivariant diffeomorphism. $\square$

5. FIT WEYL CHAMBERS AND CR GEOMETRY OF $M(\mathfrak{g}, \mathfrak{q})$

Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and $(\vartheta, \mathfrak{h})$ a Cartan pair adapted to $(\mathfrak{g}, \mathfrak{q})$. We keep the notation of the preceding sections, for roots, parabolic sets, Cartan decomposition, etc. In particular, we denote by $\sigma : \mathfrak{h}_{\mathbb{R}}^* \ni \alpha \rightarrow \bar{\alpha} \in \mathfrak{h}_{\mathbb{R}}^*$ the adjoint map of the restriction to $\mathfrak{h}_{\mathbb{R}} = \mathfrak{h}^- \oplus i\mathfrak{h}^+$ of the conjugation in $\hat{\mathfrak{g}}$ defined by the real form $\mathfrak{g}$. We say that a root α is *real* if $\bar{\alpha} = \alpha$, *imaginary* if $\bar{\alpha} = -\alpha$, *complex* if $\bar{\alpha} \neq \pm\alpha$ and denote by $\mathcal{R}_{\text{re}}$, $\mathcal{R}_{\text{im}}$ and $\mathcal{R}_{\text{cp}}$ the sets of real, imaginary and complex roots, respectively. When α is imaginary, the eigenspace $\hat{\mathfrak{g}}^\alpha$ is contained either in $\hat{\mathfrak{k}} = \mathbb{C} \otimes_{\mathbb{R}} \mathfrak{k}$, or in $\hat{\mathfrak{p}} = \mathbb{C} \otimes_{\mathbb{R}} \mathfrak{p}$. In the first case we say that α is *compact*, in the second that α is *noncompact*. Thus $\mathcal{R}_{\text{im}}$ is the disjoint union of the set $\mathcal{R}_{\bullet}$ of the compact and of the set $\mathcal{R}_*$ of the noncompact imaginary roots: $\mathcal{R}_{\text{im}} = \mathcal{R}_{\bullet} \cup \mathcal{R}_*$.

5.1. S-fit and V-fit Weyl chambers.

The conjugation σ defines an involution in $\mathfrak{h}_{\mathbb{R}}^*$ that belongs to $\mathbf{A}(\mathcal{R})$. Vice versa, every involution σ in $\mathbf{A}(\mathcal{R})$ can be obtained from a conjugation with respect to a real form $\mathfrak{g}$ of $\hat{\mathfrak{g}}$. Note that, in general, σ does not uniquely determine the isomorphism class of $\mathfrak{g}$. Let us describe the structure of an arbitrary involution σ in $\mathbf{A}(\mathcal{R})$:

Theorem 5.1. *Let σ be an involution in $\mathbf{A}(\mathcal{R})$. Then there exist: a set of pairwise strongly orthogonal roots $\alpha_1, \dots, \alpha_m$ in $\mathcal{R}$, with $\sigma(\alpha_j) = -\alpha_j$ for $j = 1, \dots, m$, a Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$, and an involution $j \in \mathbf{A}_C(\mathcal{R})$, with $j(\alpha_i) = \alpha_i$, and hence commuting with s_{α_i} , for all $i = 1, \dots, m$, such that :*

$$(5.1) \quad \sigma = j \circ s_{\alpha_1} \circ \dots \circ s_{\alpha_m} ;$$

$$(5.2) \quad \alpha \in \mathcal{R}^+(C) \implies \begin{cases} \text{either } \sigma(\alpha) = -\alpha \\ \text{or } \sigma(\alpha) \in \mathcal{R}^+(C) \end{cases}$$

Proof. Let $F^-(\sigma) = \{\alpha \in \mathfrak{h}_{\mathbb{R}}^* \mid \sigma(\alpha) = -\alpha\}$. Take a maximal subset $\alpha_1, \dots, \alpha_m$ of pairwise orthogonal roots in $F^-(\sigma) \cap \mathcal{R}$ and consider $j = \sigma \circ s_{\alpha_1} \circ \dots \circ s_{\alpha_m}$. We have $j(\alpha_i) = \sigma(-\alpha_i) = \alpha_i$ for all $i = 1, \dots, m$. We claim that $j(\alpha) \neq -\alpha$ for all $\alpha \in \mathcal{R}$. Indeed, if there was $\alpha \in \mathcal{R}$ with $j(\alpha) = -\alpha$, from $(\alpha|\alpha_i) = (j(\alpha)|j(\alpha_i)) = -(\alpha|\alpha_i)$ we obtain that $(\alpha|\alpha_i) = 0$ for all $i = 1, \dots, m$. Hence $s_{\alpha_i}(\alpha) = \alpha$ for all α and therefore $\sigma(\alpha) = j(\alpha) = -\alpha$, contradicting the fact that $\alpha_1, \dots, \alpha_m$ was a maximal system of pairwise orthogonal roots in $\mathcal{R} \cap F^-(\sigma)$.

To obtain that $\alpha_1, \dots, \alpha_m$ are strongly orthogonal it suffices to choose the sequence $\alpha_1, \dots, \alpha_m$ with a maximal sum $\sum_{i=1}^m \|\alpha_i\|^2$. Indeed, if α_j and α_h are orthogonal, but not strongly orthogonal, then both $\alpha_j + \alpha_h$ and $\alpha_j - \alpha_h$ are roots. Setting $\alpha'_i = \alpha_i$ for $i \neq j, h$, and $\alpha'_j = \alpha_j + \alpha_h$, $\alpha'_h = \alpha_j - \alpha_h$, we obtain a new sequence $\alpha'_1, \dots, \alpha'_m$ of pairwise orthogonal roots in $F^-(\sigma) \cap \mathcal{R}$. It is contained in a maximal one and the inequality $\sum_{i=1}^m \|\alpha'_i\|^2 = (\sum_{i=1}^m \|\alpha_i\|^2) + \|\alpha_j\|^2 + \|\alpha_h\|^2 > \sum_{i=1}^m \|\alpha_i\|^2$, contradicts the maximality of $\sum_{i=1}^m \|\alpha_i\|^2$.

We claim that there exists a Weyl chamber C such that :

$$(*) \quad j(\mathcal{R}^+(C)) = \mathcal{R}^+(C) .$$

Indeed $(*)$ is equivalent to $j(\mathcal{B}(C)) \subset \mathcal{R}^+(C)$. For a Weyl chamber C , denote by n_C the number of the elements in $\mathcal{R}^+(C) \cap j(\mathcal{R}^+(C))$. Fix C with n_C maximum. If C does not satisfy $(*)$, take $\alpha \in \mathcal{B}(C)$ with $j(\alpha) \notin \mathcal{R}^+(C)$ and consider the chamber $C' = s_{\alpha}(C)$. From $\mathcal{R}^+(C') = (\mathcal{R}^+(C) \setminus \{\alpha\}) \cup \{-\alpha\}$ and $j(-\alpha) \in \mathcal{R}^+(C) \setminus \{\alpha\} \subset \mathcal{R}^+(C')$, we obtain $n_{C'} = n_C + 1$, contradicting our choice of C . Hence C satisfies $(*)$ and therefore also (5.2). This completes the proof. $\square$

Using Theorem 5.1, we obtain the formula :

$$(5.3) \quad \sigma(\beta) = j(\beta) - \sum_{j=1}^m (\beta|\alpha_j^{\vee}) \alpha_j, \quad \text{with} \quad \alpha_j^{\vee} = 2\alpha_j/\|\alpha_j\|^2, \quad \forall \beta \in \mathfrak{h}_{\mathbb{R}}^* .$$

Likewise, we have the following :

Theorem 5.2. *Let σ be an involution in $\mathbf{A}(\mathcal{R})$. Then there exists a set of pairwise strongly orthogonal roots $\delta_1, \dots, \delta_m \in \mathcal{R}$, with $\sigma(\delta_j) = \delta_j$ for $j = 1, \dots, m$, a Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$ and an involution ϖ , that commutes with s_{δ_j} , satisfies $\varpi(\delta_j) = -\delta_j$ for all $j = 1, \dots, m$, and transforms C into C^{opp} , such that :*

$$(5.4) \quad \sigma = \varpi \circ s_{\delta_1} \circ \dots \circ s_{\delta_m} ,$$

$$(5.5) \quad \alpha \in \mathcal{R}^+(C) \implies \begin{cases} \text{either } \sigma(\alpha) = \alpha \\ \text{or } \sigma(\alpha) \in \mathcal{R}^-(C). \end{cases}$$

Proof. We take $\sigma' = s_0 \circ \sigma$, where s_0 is the symmetry with respect to the origin of $\mathfrak{h}_{\mathbb{R}}^*$. By the preceding Theorem, $\sigma' = j \circ s_{\delta_1} \circ \cdots \circ s_{\delta_m}$, where $j \in \mathbf{A}_C(\mathcal{R})$ for some $C \in \mathfrak{C}(\mathcal{R})$, and $\delta_1, \dots, \delta_m$ is a maximal system of strongly orthogonal roots in $F^-(\sigma') \cap \mathcal{R} = \{\alpha \in \mathcal{R} \mid \sigma(\alpha) = \alpha\}$, with $j(\delta_j) = \delta_j$. The statement follows by taking $\varpi = s_0 \circ j$. $\square$

With ϖ and $\delta_1, \dots, \delta_m$ as in Theorem 5.2, we obtain the formula :

$$(5.6) \quad \sigma(\beta) = \varpi(\beta) + \sum_{j=1}^m (\beta | \delta_j^\vee) \delta_j, \quad \text{with} \quad \delta_j^\vee = 2\delta_j / \|\delta_j\|^2, \quad \forall \beta \in \mathfrak{h}_{\mathbb{R}}^*.$$

A Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$ that satisfies condition (5.2) (resp. (5.5)) is said to be² *S-adapted* (resp. *V-adapted*) to the conjugation σ .

Our description of the minimal orbits in [AMN] in terms of *cross-marked Satake diagrams* used the fact that, for a parabolic minimal $(\mathfrak{g}, \mathfrak{q})$, there are a maximally noncompact Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$, adapted to $(\mathfrak{g}, \mathfrak{q})$, and an S-adapted Weyl chamber C in $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$. For general *CR* algebras $(\mathfrak{g}, \mathfrak{q})$, there could be no adapted Cartan subalgebras $\mathfrak{h}$ that are either maximally compact or maximally noncompact in $\mathfrak{g}$. This is a major drawback in the classification of the orbits of $\mathbf{G}$ in $\mathfrak{M}$ (see e.g. the references in [BL]), but, while discussing fundamentality, weak nondegeneracy and some topological properties, it turns out that the choice of $\mathfrak{h}$ is not as crucial as that of special Weyl chambers C in $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$. In general $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$ may not contain any Weyl chamber that is either S- or V-adapted to σ . In the following lemmas we describe chambers in $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that are as close as possible to being S- or V-adapted.

Lemma 5.3. *Let $(\mathfrak{g}, \mathfrak{h})$ be a Cartan pair adapted to $(\mathfrak{g}, \mathfrak{q})$. Then there exists a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that satisfies the equivalent conditions :*

- (i) *If $\alpha \succ_C 0$ and $\bar{\alpha} \prec_C 0$, then both α and $-\bar{\alpha}$ belong to $\mathcal{Q}^n$.*
- (ii) *$\bar{\alpha} \succ_C 0$ for all $\alpha \in \mathcal{B}(C) \setminus (\Phi_C \cup \mathcal{R}_{\text{im}})$.*

Assume that C satisfies the equivalent conditions (i) and (ii). Then :

- (iii) *If moreover $\mathfrak{h}$ is maximally noncompact among the Cartan subalgebras of $\mathfrak{g}$ contained in $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$, then $\mathcal{B}(C) \cap \mathcal{R}_* \subset \Phi_C$.*

Proof. Choose $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ with a maximal $\mathcal{R}^+(C) \cap \sigma(\mathcal{R}^+(C))$. Then (ii) is satisfied. Indeed, if there was $\alpha \in \mathcal{B}(C) \setminus (\Phi_C \cup \mathcal{R}_{\text{im}})$ with $\bar{\alpha} \prec_C 0$, we would take $C' = s_\alpha(C)$. Then $\mathcal{R}^+(C') = (\mathcal{R}^+(C) \setminus \{\alpha\}) \cup \{-\alpha\} \subset \mathcal{Q}$, so that $C' \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, and $\mathcal{R}^+(C') \cap \sigma(\mathcal{R}^+(C')) \supsetneq \mathcal{R}^+(C) \cap \sigma(\mathcal{R}^+(C))$, yielding a contradiction. Clearly (i) $\implies$ (ii). Vice versa, if $\alpha = \sum_{\beta \in \mathcal{B}(C)} k_\alpha^\beta \beta \in \mathcal{R}^+(C)$ and $\bar{\alpha} \prec_C 0$, then either $\alpha \in \mathcal{R}_{\text{im}}$, or else there is some $\beta \in \text{supp}_C(\alpha) \cap \mathcal{R}_{\text{cp}}$ with $\bar{\beta} \prec_C 0$; by (ii), we have $\beta \in \Phi_C$ and hence $\alpha \in \mathcal{Q}^n$. The same argument, applied to $-\bar{\alpha}$, shows that also $-\bar{\alpha} \in \mathcal{Q}^n$. This completes the proof of the equivalence (i) $\Leftrightarrow$ (ii).

²If we choose $\mathfrak{h}$ maximally noncompact, then in an S-adapted Weyl chamber C the conjugation can be described by a Satake diagram; if instead we take $\mathfrak{h}$ with a maximal compact part, in a V-adapted Weyl chamber the conjugation is described by a Vogan diagram (see e.g. [Ar], [Kn]).

Finally, if $\alpha \in (\mathcal{B}(C) \cap \mathcal{R}_*) \setminus \Phi_C$, both α and $\bar{\alpha} = -\alpha$ belong to $\mathcal{Q}$. Let $\Gamma = \{(X_\alpha, H_\alpha)_{\alpha \in \mathcal{R}}\}$ be a Chevalley system, as in [B2]. Then $T_\alpha = X_\alpha - X_{-\alpha} \in \mathfrak{p} \cap \mathfrak{g}_+$ (for this construction cf. [S1], [S2]) is a semisimple element of $\mathfrak{g}$ that commutes with all elements of $\mathfrak{h}^-$ and of $\mathfrak{j}^+ = \{H \in \mathfrak{h}^+ \mid \alpha(H) = 0\}$. Hence $\mathfrak{j} = \mathfrak{h}^- \oplus \mathbb{R} T_\alpha \oplus \mathfrak{j}^+$ is a Cartan subalgebra of $\mathfrak{g}$, contained in $\mathfrak{g}^+$, with $\mathfrak{j}^- = \mathfrak{h}^- \oplus \mathbb{R} T_\alpha \not\subseteq \mathfrak{h}^-$. Thus, if $\mathfrak{h}^-$ is maximal, we have $\mathcal{B}(C_0) \cap \mathcal{R}_* \subset \Phi_C$. $\square$

An alternative construction of a Weyl chamber C satisfying (i) and (ii) of Lemma 5.3 is the following (which is a particular application of the general construction that will be described at the beginning of §6). Fix a Weyl chamber $C_0 \in \mathfrak{C}(\mathcal{R})$ that is S -adapted to σ (this means that $\sigma(\mathcal{R}^+(C_0) \setminus \mathcal{R}_{\text{im}}) \subset \mathcal{R}^+(C_0)$), and consider the Borel subalgebra :

$$\mathfrak{b}_{C_0} = \hat{\mathfrak{h}} \oplus \bigoplus_{\alpha \in \mathcal{R}^+(C_0)} \hat{\mathfrak{g}}^\alpha \subset \hat{\mathfrak{g}}.$$

Then $\mathfrak{b} = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \mathfrak{b}_{C_0})$ is a Borel subalgebra of $\hat{\mathfrak{g}}$, corresponding to a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that satisfies (i) and (ii) of Lemma 5.3.

A Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that satisfies the equivalent conditions (i) and (ii) of Lemma 5.3 is called *S-fit* to $(\mathfrak{g}, \mathfrak{q})$.

With arguments similar to those employed to prove Lemma 5.3, we obtain :

Lemma 5.4. *Let $(\mathfrak{v}, \mathfrak{h})$ be a Cartan pair adapted to $(\mathfrak{g}, \mathfrak{q})$. Then there exists a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that satisfies the equivalent conditions :*

- (iv) *If $\alpha \succ_C 0$ and $\bar{\alpha} \succ_C 0$, then both α and $\bar{\alpha}$ belong to $\mathcal{Q}^n$.*
- (v) *$\bar{\alpha} \prec_C 0$ for all $\alpha \in \mathcal{B}(C) \setminus (\Phi_C \cup \mathcal{R}_{\text{re}})$.*

Assume that C satisfies the equivalent conditions (iv) and (v). Then :

- (vi) *If moreover $\mathfrak{h}$ is maximally compact among the Cartan subalgebras of $\mathfrak{g}$ contained in $\mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g}$, then $\mathcal{B}(C) \cap \mathcal{R}_{\text{re}} \subset \Phi_C$. $\square$*

A Weyl chamber $C \in \mathfrak{C}(\mathcal{Q}, \mathcal{R})$ that satisfies the equivalent conditions (iv) and (v) of Lemma 5.4 is called *V-fit* to $(\mathfrak{g}, \mathfrak{q})$.

V-fit Weyl chambers can be obtained as follows : if $\mathfrak{b}_{C_0} = \hat{\mathfrak{h}} \oplus \sum_{\alpha \in \mathcal{R}^+(C_0)} \hat{\mathfrak{g}}^\alpha$ is the Borel subalgebra associated to a Weyl chamber C_0 that is V-adapted to the conjugation σ , then $\mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \mathfrak{b}_{C_0})$ is the Borel subalgebra associated to a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that is V-fit to $(\mathfrak{g}, \mathfrak{q})$.

5.2. Fundamental parabolic CR algebras.

Theorem 5.5. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra. Fix a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$ and fix an S-fit Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ for $(\mathfrak{g}, \mathfrak{q})$. Then $(\mathfrak{g}, \mathfrak{q})$ is fundamental if and only if :*

$$(5.7) \quad \forall \alpha_0 \in \Phi_C \cap \bar{\mathcal{Q}}^n \begin{cases} \text{either } \exists \beta \in \mathcal{B}(C) \setminus \Phi_C \text{ with } \alpha_0 \in \text{supp}_C(\bar{\beta}), \\ \text{or } \exists \beta \in \Phi_C \text{ with } \bar{\beta} \in \mathcal{R}^-(C) \text{ and } \alpha_0 \in \text{supp}_C(\bar{\beta}). \end{cases}$$

Proof. The parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ is fundamental if, and only if, there is no complex parabolic subalgebra $\mathfrak{q}'$ of $\hat{\mathfrak{g}}$ with $\mathfrak{q} + \bar{\mathfrak{q}} \subset \mathfrak{q}' \subsetneq \hat{\mathfrak{g}}$. All complex parabolic $\mathfrak{q}'$ that contain $\mathfrak{q}$ are of the form $\mathfrak{q}' = \mathfrak{q}_{\Psi_C}$ for some set of simple roots $\Psi_C \subset \Phi_C$.

We can limit ourselves to consider the cases where $\Psi_C = \{\alpha_0\}$ for some $\alpha_0 \in \Phi_C$. Thus we obtain :

$$(5.8) \quad (\mathfrak{g}, \mathfrak{q}_{\Phi_C}) \text{ is not fundamental} \iff \begin{cases} \exists \alpha_0 \in \Phi_C \text{ such that} \\ \{\beta \in \mathcal{R} \mid \beta \succeq_C \alpha_0\} \subset \bar{\mathcal{Q}}^n. \end{cases}$$

Indeed, $\mathfrak{q} + \bar{\mathfrak{q}} \subset \mathfrak{q}'$ if and only if $\mathfrak{q}'^n \subset \mathfrak{q}^n \cap \bar{\mathfrak{q}}^n$. Thus it suffices to further restrict our consideration to simple roots $\alpha_0 \in \Phi_C \cap \bar{\mathcal{Q}}^n$ and check whether $\mathcal{F} = \{\beta \in \mathcal{R} \mid \beta \succeq_C \alpha_0\}$ is contained or not in $\bar{\mathcal{Q}}^n$.

First we show that condition (5.7) is sufficient. Let $\alpha_0 \in \Phi_C \cap \bar{\mathcal{Q}}^n$. If $\beta \in \mathcal{B}(C) \setminus \Phi_C$ and $\alpha_0 \in \text{supp}_C(\beta)$, then $\bar{\beta} \succeq_C \alpha_0$ by the assumption that C is S-fit to $(\mathfrak{g}, \mathfrak{q})$; hence $\bar{\beta} \in \mathcal{F} \setminus \bar{\mathcal{Q}}^n$. Likewise, if $\beta \in \Phi_C$, $\bar{\beta} \prec_C 0$ and $\alpha_0 \in \text{supp}_C(\bar{\beta})$, then $-\bar{\beta} \in \mathcal{F} \setminus \bar{\mathcal{Q}}^n$. This completes the proof of sufficiency.

To prove that (5.7) is also necessary, fix again $\alpha_0 \in \Phi_C \cap \bar{\mathcal{Q}}^n$. If $(\mathfrak{g}, \mathfrak{q})$ is fundamental, then $\mathcal{F} \not\subset \bar{\mathcal{Q}}^n$, and hence there is a root α with $\alpha \succeq_C \alpha_0$ and $\bar{\alpha} \notin \bar{\mathcal{Q}}^n$. If $\bar{\alpha} \succ_C 0$, then $\text{supp}_C(\bar{\alpha}) \cap \Phi_C = \emptyset$. Since $\alpha_0 \in \text{supp}_C(\alpha) \subset \bigcup_{\beta \in \text{supp}_C(\bar{\alpha})} \text{supp}_C(\bar{\beta})$, there is at least a $\beta \in \mathcal{B}(C) \setminus \Phi_C$ with $\alpha_0 \in \text{supp}_C(\bar{\beta})$. Assume now that α_0 does not belong to $\text{supp}_C(\bar{\beta})$ for any $\beta \in \mathcal{B} \setminus \Phi_C$. Then, for a root $\alpha \succ_C \alpha_0$ that does not belong to $\bar{\mathcal{Q}}^n$, we have $\bar{\alpha} \prec_C 0$. From $\alpha_0 \in \bigcup_{\beta \in \text{supp}_C(\bar{\alpha})} \text{supp}_C(\bar{\beta})$ we obtain that $\alpha_0 \in \text{supp}_C(\bar{\beta})$ for some $\beta \in \mathcal{B}(C)$ with $\bar{\beta} \prec_C 0$, and hence in Φ_C . The proof is complete. $\square$

Theorem 5.5 provides a criterion, only involving the conjugation of the simple roots of an S-fit Weyl chamber, for an effective parabolic CR algebra to be totally real. We found convenient to formulate the criterion for an arbitrary Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$.

Lemma 5.6. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra, and $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$. A necessary and sufficient condition in order that $(\mathfrak{g}, \mathfrak{q})$ be totally real is that :*

$$(5.9) \quad \begin{cases} \alpha \in \mathcal{B}(C) \setminus \Phi_C & \implies & \text{supp}_C(\bar{\alpha}) \cap \Phi_C = \emptyset \\ \alpha \in \Phi_C & \implies & \bar{\alpha} \in \mathcal{R}^+(C). \end{cases}$$

Proof. The case where $\mathcal{Q} = \mathcal{R}$ is trivial. Assume that $\mathcal{Q} \neq \mathcal{R}$. The first condition in (5.9) implies that $\mathcal{B}(C) \setminus \Phi_C \subset \bar{\mathcal{Q}}^r$. Hence $\mathcal{Q}^r = \bar{\mathcal{Q}}^r$. In particular, if $\alpha \in \mathcal{Q}^n$, then $\bar{\alpha} \in \mathcal{R} \setminus \mathcal{Q}^r$. Then, since $\mathcal{Q}^n = \mathcal{R}^+(C) \setminus \mathcal{Q}^r$, the second condition implies that $\bar{\alpha} \in \mathcal{Q}^n$ for all $\alpha \in \Phi_C$. Hence $\Phi_C \subset \bar{\mathcal{Q}}^n$. Therefore $\mathcal{B}(C) \subset \bar{\mathcal{Q}}$, so that we also have $\bar{\mathcal{Q}}^n = \mathcal{Q}^n$ and hence $\mathcal{Q} = \bar{\mathcal{Q}}$. The condition is obviously also necessary. $\square$

Lemma 5.6 prompts a recursive method to construct the totally real basis of the canonical $\mathfrak{g}$ -equivariant fibration $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}')$, with totally real basis and fundamental fiber.

After taking any $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, we define recursively :

$$(5.10) \quad \begin{cases} \Upsilon_C^{(0)} &= \{\alpha \in \Phi_C \mid \bar{\alpha} \in \mathcal{R}^+(C)\} \\ \Upsilon_C^{(1)} &= \Upsilon_C^{(0)} \setminus \bigcup_{\alpha \in \mathcal{B}(C) \setminus \Upsilon_C^{(0)}} \text{supp}_C(\bar{\alpha}) \\ \Upsilon_C^{(h+1)} &= \Upsilon_C^{(h)} \setminus \bigcup_{\alpha \in \mathcal{B}(C) \setminus \Upsilon_C^{(h)}} \text{supp}_C(\bar{\alpha}) \quad \text{for } h \geq 1 \\ \Upsilon_C &= \bigcap_{h \geq 0} \Upsilon_C^{(h)} \quad (\text{finite intersection}). \end{cases}$$

One easily verifies, using the previous results, that :

Proposition 5.7. *The natural $\mathfrak{g}$ -equivariant fibration $(\mathfrak{g}, \mathfrak{q}_{\Phi_C}) \rightarrow (\mathfrak{g}, \mathfrak{q}_{\Upsilon_C})$ is the fundamental reduction of $(\mathfrak{g}, \mathfrak{q}_{\Phi_C})$. In particular, $(\mathfrak{g}, \mathfrak{q}_{\Phi_C})$ is fundamental if and only if $\Upsilon_C = \emptyset$. $\square$*

5.3. Weakly nondegenerate parabolic CR algebras.

We turn now to weak nondegeneracy for an effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$. We shall see that this property can be better examined in terms of V-fit Weyl chambers.

Lemma 5.8. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra, $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$, and $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ a V-fit Weyl chamber for $(\mathfrak{g}, \mathfrak{q})$. Let $\Phi_C = \mathcal{B}(C) \cap \mathcal{Q}^n$ and $\mathfrak{q}' = \mathfrak{q}_{\Psi_C} \supset \mathfrak{q}$ (cf. (3.12) for the notation), with $\Psi_C \subset \Phi_C$. Then the $\mathfrak{g}$ -equivariant homomorphism of CR algebras $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}')$ is a CR fibration with a totally complex fiber if and only if*

$$(5.11) \quad \bar{\alpha} \prec_C 0 \quad \forall \alpha \in \Phi_C \setminus \Psi_C.$$

Proof. The $\mathfrak{g}$ -equivariant CR homomorphism $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}')$ is a CR fibration with a totally complex fiber if, and only if, $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$. Thus, we need to show that these inclusions are equivalent to (5.11) when $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ satisfies (iv) and (v) of Lemma 5.4.

Clearly, it suffices to consider the case where the difference $\Phi_C \setminus \Psi_C$ consists of a single simple root α_0 . We shall assume in the following that $\Phi_C \setminus \Psi_C = \{\alpha_0\}$.

First we prove that, if $\bar{\alpha}_0 \prec_C 0$, then $\mathcal{Q}' = \mathcal{Q}_{\Psi_C} \subset \mathcal{Q} \cup \bar{\mathcal{Q}}$. Assume by contradiction that there is $\beta \in \mathcal{Q}' \setminus (\mathcal{Q} \cup \bar{\mathcal{Q}})$. Then $-\beta, -\bar{\beta} \in \mathcal{Q}^n \subset \mathcal{R}^+(C)$. Moreover $\beta \in \mathcal{Q}'^r$, because $\mathcal{Q}'^n \subset \mathcal{Q}^n \subset (\mathcal{Q} \cup \bar{\mathcal{Q}})$. Thus $\text{supp}_C(\beta) \cap \Psi_C = \emptyset$, and hence $\text{supp}_C(\beta) \cap \Phi_C = \{\alpha_0\}$, because $-\beta \in \mathcal{Q}^n$. As C is V-fit, $\bar{\alpha} \prec_C 0$ for all $\alpha \in \text{supp}_C(\beta) \setminus \mathcal{R}_{\text{re}}$. Since β is not real, this implies that $\bar{\beta} \succ_C 0$, giving a contradiction. Hence $\mathcal{Q}' \subset (\mathcal{Q} \cup \bar{\mathcal{Q}})$, and this proves that $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$ when $\bar{\alpha}_0 \in \mathcal{R}^-(C)$.

Vice versa, assume that $\mathfrak{q} \subset \mathfrak{q}' \subset \mathfrak{q} + \bar{\mathfrak{q}}$. In particular, $-\alpha_0 \in \bar{\mathcal{Q}}$. If $\bar{\alpha}_0 \in \mathcal{Q}^r$, then $\bar{\alpha}_0$ and $\alpha_0 = \bar{\bar{\alpha}}_0$ belong to opposite cones $\mathcal{R}^\pm(C)$ and thus $\bar{\alpha}_0 \prec_C 0$. When $\bar{\alpha}_0 \notin \mathcal{Q}^r$, we have $-\bar{\alpha}_0 \in \mathcal{Q}^n \subset \mathcal{R}^+(C)$, and thus $\bar{\alpha}_0 \prec_C 0$. The proof is complete. $\square$

Theorem 5.9. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra, and let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$. Let $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ be V-fit to $(\mathfrak{g}, \mathfrak{q})$. Then $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate if and only if:*

$$(5.12) \quad \bar{\alpha} \succ_C 0 \quad \forall \alpha \in \Phi_C.$$

Proof. Fix a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ for which (iv) and (v) of Lemma 5.4 are valid. By Lemma 5.8, the necessary and sufficient condition for $(\mathfrak{g}, \mathfrak{q})$ to be weakly degenerate is that there exists $\alpha_0 \in \Phi_C$ contradicting (5.12). $\square$

Corollary 5.10. *Let $(\mathfrak{g}, \mathfrak{q})$ be a weakly nondegenerate parabolic CR algebra. Let $\mathcal{Q}$ be the parabolic set associated to $\mathfrak{q}$ in $\mathcal{R} = \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}})$, for an admissible Cartan subalgebra $\mathfrak{h}$ of $(\mathfrak{g}, \mathfrak{q})$. Then $(\mathfrak{g}, \mathfrak{q})$ is totally real if and only if $\bar{\mathcal{Q}}^n \cap \mathcal{Q}^r = \emptyset$.*

Proof. Choose a V-fit Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ for $(\mathfrak{g}, \mathfrak{q})$. The second condition in (5.9) being automatically satisfied because C is V-fit and $(\mathfrak{g}, \mathfrak{q})$ weakly nondegenerate, we observe that the first line in (5.9) is equivalent to the condition that $\bar{\mathcal{Q}}^n \cap \mathcal{Q}^r = \emptyset$. $\square$

5.4. A criterion of minimality.

In [AMN] we only used maximally noncompact adapted Cartan subalgebras. Here we give a characterization of effective parabolic minimal CR algebras $(\mathfrak{g}, \mathfrak{q})$ in terms of the set of roots $\mathcal{Q}$ associated to $\mathfrak{q}$ by *any* choice of an adapted Cartan pair $(\vartheta, \mathfrak{h})$.

Proposition 5.11. *A necessary and sufficient condition in order that a weakly nondegenerate effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ be parabolic minimal³, is that*

$$(5.13) \quad \mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n} \subset \mathcal{R}_\bullet.$$

Proof. A necessary and sufficient condition for $(\mathfrak{g}, \mathfrak{q})$ to be minimal is that $\mathfrak{g}_+ + \mathfrak{k} = \mathfrak{g}$. By complexification this condition can be rewritten as:

$$(5.14) \quad \hat{\mathfrak{g}} = \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}.$$

Since $\hat{\mathfrak{h}} \subset \mathfrak{q} \cap \bar{\mathfrak{q}}$, it suffices to show that (5.13) is equivalent to:

$$(5.15) \quad \hat{\mathfrak{g}}^\alpha \subset \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}} \quad \text{for all } \alpha \in \mathcal{R}.$$

To make the equivalence more clear, we first prove:

Lemma 5.12. *For all $\alpha \notin (\mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n}) \cup (\mathcal{Q}^{-n} \cap \bar{\mathcal{Q}}^n) = (\mathcal{Q}^n \setminus \bar{\mathcal{Q}}) \cup (\bar{\mathcal{Q}}^n \setminus \mathcal{Q})$ we have*

$$\hat{\mathfrak{g}}^\alpha \subset \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}.$$

Proof. We get: $\mathcal{R} \setminus [(\mathcal{Q}^n \setminus \bar{\mathcal{Q}}) \cup (\bar{\mathcal{Q}}^n \setminus \mathcal{Q})] = \mathcal{Q}^r \cup \bar{\mathcal{Q}}^r \cup (\mathcal{Q}^n \cap \bar{\mathcal{Q}}) \cup (\bar{\mathcal{Q}}^n \cap \mathcal{Q})$.

Clearly $\hat{\mathfrak{g}}^\alpha \subset \mathfrak{q} \cap \bar{\mathfrak{q}} \subset \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}$ when $\alpha \in (\mathcal{Q}^n \cap \bar{\mathcal{Q}}) \cup (\bar{\mathcal{Q}}^n \cap \mathcal{Q})$.

Since the statement is invariant if we interchange $\mathfrak{q}$ and $\bar{\mathfrak{q}}$, to complete the proof of the lemma it suffices to show that $\hat{\mathfrak{g}}^\alpha \subset \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}$ for all $\alpha \in \mathcal{Q}^r$.

Let $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ be V-fit, so that $\sigma((\mathcal{R}^+(C) \cap \mathcal{Q}^r) \setminus \mathcal{R}_{\text{re}}) \subset \mathcal{R}^-(C)$. We also use the notation $\hat{\mathfrak{k}}^{\alpha, -\bar{\alpha}} = \hat{\mathfrak{k}} \cap (\hat{\mathfrak{g}}^\alpha + \hat{\mathfrak{g}}^{-\bar{\alpha}})$. This is a one-dimensional complex subspace of $\hat{\mathfrak{g}}$ when $\alpha \in \mathcal{R} \setminus \mathcal{R}_*$.

For $\alpha \in \mathcal{Q}^r \cap (\mathcal{R}_{\text{re}} \cup \mathcal{R}_{\text{im}})$, we have $\alpha \in \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r \subset \mathcal{Q} \cap \bar{\mathcal{Q}}$ and hence $\hat{\mathfrak{g}}^\alpha \subset \mathfrak{q} \cap \bar{\mathfrak{q}}$.

If $\alpha \in \mathcal{Q}^r \cap \mathcal{R}^+(C) \cap \mathcal{R}_{\text{cp}}$, then $-\bar{\alpha} \in \mathcal{R}^+(C) \subset \mathcal{Q}$. Hence $-\alpha, -\bar{\alpha} \in \mathcal{Q} \cap \bar{\mathcal{Q}}$, so that: $\hat{\mathfrak{g}}^{-\alpha} \subset \mathfrak{q} \cap \bar{\mathfrak{q}} \subset \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}$, $\hat{\mathfrak{g}}^\alpha \subset \hat{\mathfrak{g}}^{-\bar{\alpha}} + \hat{\mathfrak{k}}^{(\alpha, -\bar{\alpha})} \subset \mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}$. $\square$

We conclude now the proof of Proposition 5.11. If condition (5.13) is satisfied, then $(\mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n}) \cup (\mathcal{Q}^{-n} \cap \bar{\mathcal{Q}}^n) \subset \mathcal{R}_\bullet$, hence $\hat{\mathfrak{g}}^\alpha \subset \hat{\mathfrak{k}}$ for all $\alpha \in (\mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n}) \cup (\mathcal{Q}^{-n} \cap \bar{\mathcal{Q}}^n)$. In view of Lemma 5.12, we obtain (5.15) and hence (5.14).

Vice versa, assume that there is $\alpha \in \mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n} \setminus \mathcal{R}_\bullet$. If $\alpha \in \mathcal{R}_*$, then $\hat{\mathfrak{g}}^\alpha \in \hat{\mathfrak{p}}$, and cannot be contained in $\mathfrak{q} \cap \bar{\mathfrak{q}} + \hat{\mathfrak{k}}$. Otherwise, $\alpha \in \mathcal{R}_{\text{cp}}$ and $\alpha, -\bar{\alpha} \in \mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n}$. This implies that $\hat{\mathfrak{g}}^\alpha \oplus \hat{\mathfrak{g}}^{\bar{\alpha}} \oplus \hat{\mathfrak{g}}^{-\alpha} \oplus \hat{\mathfrak{g}}^{-\bar{\alpha}} = \hat{\mathfrak{k}}^{(\alpha, -\bar{\alpha})} \oplus \hat{\mathfrak{k}}^{(-\alpha, \bar{\alpha})} \oplus \hat{\mathfrak{p}}^{(\alpha, -\bar{\alpha})} \oplus \hat{\mathfrak{p}}^{(\alpha, -\bar{\alpha})}$ has intersection $\{0\}$ with $\mathfrak{q} \cap \bar{\mathfrak{q}}$ and therefore (5.14) cannot possibly hold true. $\square$

³cf. [AMN]. Recall that the condition is equivalent to the fact that the corresponding CR manifold $M(\mathfrak{g}, \mathfrak{q})$ is the minimal orbit.

5.5. Cross-marked diagrams and examples.

In the examples that will follow, here and in the next sections, to describe specific parabolic CR algebras, we shall utilize *cross-marked diagrams*. They are Dynkin diagrams, where the simple roots in $\mathcal{B}(C)$ for a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, are indicated by :

- if the root is real;
- if the root is compact imaginary;
- ⊗ if the root is noncompact imaginary;
- ⊕ if the root is complex and its conjugate belongs to $\mathcal{R}^+(C)$;
- ⊖ if the root is complex and its conjugate belongs to $\mathcal{R}^-(C)$

and we cross-mark the roots in Φ_C . Some extra information about the action of σ on the simple roots in $\mathcal{B}(C)$ is provided by some arrows and dotted arrows joining pairs of simple roots that have the same, or opposite, restriction to $\mathfrak{h}^- \subset \mathfrak{h}_{\mathbb{R}}$, or that are the edges of segments whose nodes are the support of real or imaginary roots. However, as we shall see, the most important information is carried by the *colors* of the nodes.

Example 5.1. Consider the CR manifold M consisting of 3-planes ℓ_3 of $\mathbb{C}^6$ with $\dim_{\mathbb{C}}(\ell_3 \cap \bar{\ell}_3) = 1$. This is an orbit of $\mathbf{SL}(6, \mathbb{R})$ in the Grassmannian of 3-planes of $\mathbb{C}^6$. Let $\varepsilon_1, \dots, \varepsilon_6$ be the canonical basis of $\mathbb{C}^6$. It is convenient to represent the Lie algebra $\mathfrak{sl}(6, \mathbb{R})$ in the basis


$$e_1 = \varepsilon_1 + i\varepsilon_6, \quad e_2 = \varepsilon_2 + i\varepsilon_5, \quad e_3 = \varepsilon_3, \quad e_4 = \varepsilon_4, \quad e_5 = \varepsilon_2 - i\varepsilon_5, \quad e_6 = \varepsilon_1 - i\varepsilon_6.$$

Then M is the orbit of the 3 plane generated by e_1, e_2, e_3 .

We can take $\mathfrak{h}_{\mathbb{R}}$ to be the set of real diagonal matrices. The parabolic $\mathfrak{q}$ is $\mathfrak{q}_A$ for $A = \text{diag}(1, 1, 1, -1, -1, -1)$. The Weyl chamber C corresponding to the canonical basis $\alpha_i = e_i - e_{i+1}$ ($i = 1, \dots, 5$) belongs to $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$ and is V-fit. We have indeed

$$\begin{cases} \bar{\alpha}_1 = e_6 - e_5 = -\alpha_5 \\ \bar{\alpha}_2 = e_5 - e_3 = -(\alpha_3 + \alpha_4) \\ \bar{\alpha}_3 = e_3 - e_4 = \alpha_3 \\ \bar{\alpha}_4 = e_4 - e_2 = -(\alpha_2 + \alpha_3) \\ \bar{\alpha}_5 = e_2 - e_1 = -\alpha_1 \end{cases}$$

so that the associated diagram is:




We have $\Phi_C = \{\alpha_3\}$, $\sigma([\mathcal{R}^+(C) \cap \mathcal{Q}^*] \setminus \mathcal{R}_{\text{re}}) \subset \mathcal{R}^-(C)$, $\bar{\Phi}_C = \Phi_C \subset \mathcal{R}^+(C)$. Hence, by Lemma 5.4, $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate. Our M is a CR manifold of hypersurface type $(8, 1)$. Its Levi form has two positive, two negative and four zero eigenvalues. Hence M is fundamental and weakly, but not strictly nondegenerate.

We obtain an S-fit chamber by describing $\mathfrak{sl}(6, \mathbb{R})$ in the basis:

$$e_1 = \varepsilon_1, \quad e_2 = \varepsilon_2 + i\varepsilon_5, \quad e_3 = \varepsilon_3 + i\varepsilon_4, \quad e_4 = \varepsilon_3 - i\varepsilon_4, \quad e_5 = \varepsilon_2 - i\varepsilon_5, \quad e_6 = \varepsilon_6.$$

One verifies that, with the basis of simple roots $\alpha_i = e_i - e_{i+1}$ ($1 \leq i \leq 5$) the corresponding diagram is :



We have $\bar{\alpha}_1 = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 \succ_C \alpha_3$, in accordance with the fact that $(\mathfrak{g}, \mathfrak{q})$ is fundamental.

Example 5.2. Consider the CR manifold M^* consisting of all flags $\ell_1 \subset \ell_2 \subset \ell_4 \subset \ell_5 \subset \mathbb{C}^6$ (the subscript is the dimension of the linear subspace), with

$$\dim_{\mathbb{C}}(\ell_1 \cap \bar{\ell}_1) = 0, \quad \ell_2 = \ell_1 + \bar{\ell}_1, \quad \dim_{\mathbb{C}}(\ell_4 \cap \bar{\ell}_4) = 3, \quad \dim_{\mathbb{C}}(\ell_5 \cap \bar{\ell}_5) = 3.$$

We note that M^* is not connected. Thus an orbit of $\mathbf{SL}(6, \mathbb{R})$ in M^* will be a connected component M of M^* . We can better describe such an M in terms of the choice of a suitable basis of $\mathbb{C}^6$. Denoting by $\varepsilon_1, \dots, \varepsilon_6$ the canonical basis of $\mathbb{C}^6$, we introduce the basis :

$$e_1 = \varepsilon_1 + i\varepsilon_2, \quad e_2 = \varepsilon_1 - i\varepsilon_2, \quad e_3 = \varepsilon_3, \quad e_4 = \varepsilon_4 + i\varepsilon_6, \quad e_5 = \varepsilon_5, \quad e_6 = \varepsilon_4 - i\varepsilon_6.$$

Our M is the orbit, under the action of $\mathbf{SL}(6, \mathbb{R})$, of the flag

$$\langle e_1 \rangle \subset \langle e_1, e_2 \rangle \subset \langle e_1, e_2, e_3, e_4 \rangle \subset \langle e_1, e_2, e_3, e_4, e_5 \rangle.$$

We can take the Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{sl}(6, \mathbb{R})$ that is described, in the basis $e_1, \dots, e_6$, by :

$$\mathfrak{h} = \{ \text{diag}(\lambda, \bar{\lambda}, a, \mu, b, \bar{\mu}) \mid \lambda, \mu \in \mathbb{C}, \quad a, b \in \mathbb{R}, \quad 2\text{Re}(\lambda + \mu) + a + b = 0 \}$$

and consider the corresponding root system $\mathcal{R}$ of $\mathfrak{sl}(6, \mathbb{C})$ with respect to $\hat{\mathfrak{h}}$. We identify $\mathfrak{h}_{\mathbb{R}}$ with the space of real diagonal matrices in $\mathfrak{sl}(6, \mathbb{C})$ and the e_h 's to the evaluation of the h -th diagonal entry of $H \in \mathfrak{h}_{\mathbb{R}}$. Let $(\mathfrak{g}, \mathfrak{q})$ be the effective parabolic CR algebra associated to M and $\mathcal{Q}$ the parabolic set of $\mathfrak{q}$. Then the Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$ with basis $\mathcal{B}(C) = \{\alpha_i = e_i - e_{i+1} \mid 1 \leq i \leq 5\}$ belongs to $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$ and $\Phi_C = \{\alpha_1, \alpha_2, \alpha_4, \alpha_5\}$. We have :


$$\begin{cases} \bar{\alpha}_1 = -\alpha_1 \\ \bar{\alpha}_2 = \alpha_1 + \alpha_2 \\ \bar{\alpha}_3 = \alpha_3 + \alpha_4 + \alpha_5 \\ \bar{\alpha}_4 = -\alpha_5 \\ \bar{\alpha}_5 = -\alpha_4. \end{cases}$$

Thus (5.10) yields in this case :


$$\begin{cases} \Upsilon_C^{(0)} = \{\alpha_2\} \\ \Upsilon_C^{(1)} = \Upsilon_C^{(0)} = \Upsilon_C. \end{cases}$$

Thus the basis of the fundamental reduction is the Grassmannian of 2-planes ℓ_2 in $\mathbb{C}^6$ with $\bar{\ell}_2 = \ell_2$. We give below the diagrams for $(\mathfrak{g}, \mathfrak{q})$, for its fundamental


reduction $(\mathfrak{g}, \mathfrak{q}')$, and for the fiber $(\mathfrak{g}, \mathfrak{q}'')$, (recall that the basis of the fundamental reduction of a parabolic is still parabolic).



The fiber M'' is the product of a complex disk (a connected component of the set of $\mathbb{CP}^1 \setminus \mathbb{RP}^1 \simeq S^2 \setminus S^1$) and a connected CR manifold N , consisting of flags in $\mathbb{C}^6/\mathbb{C}^2$: these can be identified to pairs $\ell_2 \subset \ell_3 \subset \mathbb{C}^4$ such that $\dim_{\mathbb{C}}(\ell_2 \cap \bar{\ell}_2) = 1$, $\ell_3 \not\supset \ell_2 + \bar{\ell}_2$ and $\dim_{\mathbb{C}}(\ell_3 \cap \bar{\ell}_3) = 2$. It is convenient to utilize the basis e_4, e_3, e_5, e_6 of $\mathbb{C}^4 \simeq \langle e_3, e_4, e_5, e_6 \rangle \subset \mathbb{C}^6$. Then $\beta_1 = e_4 - e_3, \beta_2 = e_3 - e_5, \beta_3 = e_5 - e_6$ is the basis related to a Weyl chamber C_N in which the diagram associated to the parabolic CR algebra of N is:



Since C_N is V-fit, we see from this diagram that N is weakly degenerate. The basis N' of its weakly nondegenerate reduction, consists of planes ℓ_2 of $\mathbb{C}^4$ with $\dim_{\mathbb{C}}(\ell_2 \cap \bar{\ell}_2) = 1$, and corresponds to the diagram:



The parabolic CR manifold N' is of hypersurface type $(3, 1)$, with a degenerate Levi form of signature $(1, -1, 0)$. Thus it is 1-pseudoconcave (see e.g. [HN]) and weakly, but not strictly, nondegenerate. The fiber F of the $\mathbf{SL}(4, \mathbb{R})$ -equivariant weakly nondegenerate reduction $N \rightarrow N'$, that lies above a given 2-plane ℓ_2 with $\dim_{\mathbb{C}}(\ell_2 + \bar{\ell}_2) = 3$, is isomorphic to the pencil of 3-planes in $\mathbb{C}^4$, that contain ℓ_2 and are distinct from $(\ell_2 + \bar{\ell}_2)$. Thus $F \simeq \mathbb{CP}^1 \setminus \{\text{a point}\} \simeq \mathbb{C}$. Note that the CR algebra that is naturally associated to the fiber fails in this case to be parabolic. In fact, the CR algebra $(\mathfrak{g}^\sharp, \mathfrak{q}^\sharp)$ of the fiber is given by:

$$\mathfrak{g}^\sharp = \left\{ \begin{pmatrix} \lambda & 0 & \bar{\zeta} & 0 \\ z & h & s & \bar{z} \\ 0 & 0 & k & 0 \\ 0 & 0 & \zeta & \bar{\lambda} \end{pmatrix} \middle| \begin{array}{l} \lambda, z, \zeta \in \mathbb{C} \\ h, k, s \in \mathbb{R} \\ h + k + 2\operatorname{Re}\lambda = 0 \end{array} \right\},$$

$$\mathfrak{q}^\sharp = \left\{ \begin{pmatrix} \lambda_1 & 0 & \zeta_2 & 0 \\ 0 & \lambda_2 & \theta & z_2 \\ 0 & 0 & \lambda_3 & 0 \\ 0 & 0 & \zeta_1 & \lambda_4 \end{pmatrix} \middle| \begin{array}{l} \lambda_i, z_i, \zeta_i, \theta \in \mathbb{C} \\ \sum_{i=1}^4 \lambda_i = 0 \end{array} \right\},$$

where both $\mathfrak{g}^\sharp$ and its complexification $\hat{\mathfrak{g}}^\sharp$ are nilpotent.

6. CANONICAL FIBRATIONS OVER A PARABOLIC CR MANIFOLD

Given an effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, we constructed in the previous sections new parabolic complex subalgebras $\mathfrak{q}'$ of $\hat{\mathfrak{g}}$ with $\mathfrak{q} \subset \mathfrak{q}'$, to obtain smooth fibrations $M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}')$, namely with a weakly nondegenerate basis and totally complex fibers, and with totally real basis and fundamental fibers. Here, and in the next section, we consider smooth fibrations $M(\mathfrak{g}, \mathfrak{q}') \rightarrow M(\mathfrak{g}, \mathfrak{q})$, obtained by choosing special parabolic $\mathfrak{q}' \subset \hat{\mathfrak{g}}$ with $\mathfrak{q}' \subset \mathfrak{q}$, and that will be useful to find suitable Weyl chambers in $\mathfrak{C}(\mathcal{R}, \mathcal{Q})$ and to investigate the topology of the general $M(\mathfrak{g}, \mathfrak{q})$.

We keep the notation of the preceding sections. In particular, we fix a Cartan pair $(\vartheta, \mathfrak{h})$, assuming that it is adapted to all the parabolic CR algebras that we shall consider.

We have:

Proposition 6.1. *Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebras $(\mathfrak{g}, \mathfrak{q})$ and $(\mathfrak{g}, \mathfrak{e})$. Then:*

$$(6.1) \quad \mathfrak{l} = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \mathfrak{e})$$

is a parabolic complex Lie subalgebra of $\hat{\mathfrak{g}}$ with:

$$(6.2) \quad \begin{cases} \mathfrak{h} \subset \mathfrak{l} \\ \mathfrak{l}^n = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \mathfrak{e}^n) \supset \mathfrak{q}^n \\ \mathfrak{l}^r = \mathfrak{q}^r \cap \mathfrak{e}^r \subset \mathfrak{q}^r \\ \mathfrak{l} = \mathfrak{l}^n \oplus \mathfrak{l}^r \subset \mathfrak{q}. \end{cases}$$

Proof. Let $\mathcal{Q}, \mathcal{E}$ be the parabolic sets in $\mathcal{R}$ corresponding to the complex parabolic Lie subalgebras $\mathfrak{q}, \mathfrak{e}$, respectively. To prove that $\mathfrak{l}$ is parabolic, we need to prove that

$$(6.3) \quad \mathcal{L} = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \mathcal{E}) = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \mathcal{E}^r) \cup (\mathcal{Q}^r \cap \mathcal{E}^n)$$

is a parabolic subset of $\mathcal{R}$.

Let $A, B \in \mathfrak{h}_{\mathbb{R}}$ be such that $\mathcal{Q} = \mathcal{Q}_A, \mathcal{E} = \mathcal{Q}_B$ and fix $\varepsilon > 0$ so small that $\varepsilon |\alpha(B)| < \alpha(A)$ for all $\alpha \in \mathcal{Q}^n$. Then we claim that

$$\mathcal{L} = \{\alpha \in \mathcal{R} \mid \alpha(A + \varepsilon B) \geq 0\}.$$

Indeed, when $\alpha \in \mathcal{Q}^n$, then $\alpha(A + \varepsilon B) \geq \alpha(A) - \varepsilon |\alpha(B)| > 0$; when $\alpha \notin \mathcal{Q}$, then $-\alpha \in \mathcal{Q}^n$ and hence $\alpha(A + \varepsilon B) < 0$; finally for $\alpha \in \mathcal{Q}^r$, we have $\alpha(A + \varepsilon B) = \varepsilon \alpha(B)$ and hence $\alpha \in \mathcal{L}$ if and only if $\alpha \in \mathcal{E}$.

The proof of (6.2) is straightforward. $\square$

Vice versa, when $\mathfrak{l}$ is a complex parabolic subalgebra of $\hat{\mathfrak{g}}$ with $\mathfrak{h} \subset \mathfrak{l} \subset \mathfrak{q}$, then $\mathfrak{l} = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \mathfrak{l})$, so that (6.1) gives a way to construct all complex parabolic subalgebras of $\hat{\mathfrak{g}}$ with $\mathfrak{h} \subset \mathfrak{l} \subset \mathfrak{q}$.

Proposition 6.2. *Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$. Consider the complex parabolic Lie subalgebra*

$$(6.4) \quad \mathfrak{w} = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \vartheta(\mathfrak{q}))$$

of $\hat{\mathfrak{g}}$. The σ -invariant reductive subalgebra $\mathfrak{q}^r \cap \bar{\mathfrak{q}}^r$ is a complement in $\mathfrak{w}$ of its nilradical $\mathfrak{w}^n$. We have:

$$(6.5) \quad \begin{cases} \mathfrak{w}^r = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r, & \mathfrak{w}^n = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \vartheta(\mathfrak{q}^n)) \supset \mathfrak{q}^n, & \mathfrak{w} = \mathfrak{w}^n \oplus \mathfrak{w}^r \subset \mathfrak{q}, \\ \mathfrak{w}^n \cap \bar{\mathfrak{w}}^n = \mathfrak{q}^n \cap \bar{\mathfrak{q}}^n, & \mathfrak{w}^r = \bar{\mathfrak{w}}^r, & \mathfrak{w} \cap \bar{\mathfrak{w}} = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r \oplus \mathfrak{q}^n \cap \bar{\mathfrak{q}}^n, \end{cases}$$

and $\mathfrak{w}$ is the smallest parabolic subalgebra of $\hat{\mathfrak{g}}$ that satisfies the conditions:

$$(6.6) \quad \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r \subset \mathfrak{w} \subset \mathfrak{q} \quad \text{and} \quad \mathfrak{w} + \bar{\mathfrak{w}} = \mathfrak{q} + \bar{\mathfrak{q}}.$$

Proof. By Proposition 6.1, $\mathfrak{w}$ is complex parabolic in $\hat{\mathfrak{g}}$. Indeed $\vartheta(\mathfrak{q})$ is complex parabolic in $\hat{\mathfrak{g}}$ and contains $\mathfrak{h}$. The parabolic set associated to $\vartheta(\mathfrak{q})$ is $\vartheta(\mathcal{Q}) = \{\alpha \mid -\bar{\alpha} \in \mathcal{Q}\} = \bar{\mathcal{Q}}^r \cup \bar{\mathcal{Q}}^{-n}$. Hence the parabolic set corresponding to $\mathfrak{w}$ is:

$$(6.7) \quad \mathcal{W} = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \vartheta(\mathcal{Q})) = (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r) \cup \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^{-n}).$$

We obtain (6.5) by using Proposition 6.1.

We have $\mathcal{W}^r = \bar{\mathcal{W}}^r = \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r$, and $\mathcal{W} \cup \bar{\mathcal{W}} = \mathcal{W}^r \cup \mathcal{W}^n \cup \bar{\mathcal{W}}^n = \mathcal{Q} \cup \bar{\mathcal{Q}}$. The right hand side of this equality can be written as a disjoint union:

$$\mathcal{Q} \cup \bar{\mathcal{Q}} = (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r) \cup (\mathcal{Q}^n \cup \bar{\mathcal{Q}}^n) \cup (\mathcal{Q}^r \setminus \bar{\mathcal{Q}}) \cup (\bar{\mathcal{Q}}^r \setminus \mathcal{Q}).$$

In particular, $\mathcal{Q}^r \setminus \bar{\mathcal{Q}} \subset \mathcal{W}$ for the parabolic set $\mathcal{W}$ of any complex parabolic $\mathfrak{w}$ that satisfies (6.6), and this shows that the $\mathfrak{w}$ we constructed is the smallest complex parabolic subalgebra of $\hat{\mathfrak{g}}$ that satisfies (6.6). $\square$

Since $\mathfrak{w} \subset \mathfrak{q}$, we have a $\mathfrak{g}$ -equivariant CR fibration $(\mathfrak{g}, \mathfrak{w}) \rightarrow (\mathfrak{g}, \mathfrak{q})$. We call $(\mathfrak{g}, \mathfrak{w})$ the *canonical CR-lift* of $(\mathfrak{g}, \mathfrak{q})$.

Theorem 6.3. *The canonical CR lift $(\mathfrak{g}, \mathfrak{w}) \rightarrow (\mathfrak{g}, \mathfrak{q})$ is a $\mathfrak{g}$ -equivariant CR fibration (in particular a CR submersion) with totally complex fibers. When $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate, then $(\mathfrak{g}, \mathfrak{w}) \rightarrow (\mathfrak{g}, \mathfrak{q})$ is the weakly nondegenerate reduction of $(\mathfrak{g}, \mathfrak{w})$ (cf. [MN4]).*

Proof. The statement is a consequence of Lemma 5.8 and of the next lemma. $\square$

Lemma 6.4. *Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$. Let $(\mathfrak{g}, \mathfrak{w})$ be its canonical CR-lift. Then a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ is V-fit for $(\mathfrak{g}, \mathfrak{q})$ if and only if $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W})$ and is V-fit for $(\mathfrak{g}, \mathfrak{w})$.*

Proof. Assume that $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ is V-fit for $(\mathfrak{g}, \mathfrak{q})$. We want to show that $\mathcal{R}^+(C) \subset \mathcal{W}$. Since $\mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r) \subset \mathcal{W}$, it suffices to prove that $\alpha \in \mathcal{W}$ for $\alpha \in (\mathcal{R}^+(C) \cap \mathcal{Q}^r) \setminus \bar{\mathcal{Q}}^r$. Since C is V-fit for $(\mathfrak{g}, \mathfrak{q})$, for such a root α we have $\bar{\alpha} \prec_C 0$. Hence $\vartheta(\alpha) = -\bar{\alpha} \in \mathcal{R}^+(C) \subset \mathcal{Q}$, i.e. $\alpha \in \mathcal{Q}^r \cap \vartheta(\mathcal{Q}) \subset \mathcal{W}$. A chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W})$ that is V-fit for $(\mathfrak{g}, \mathfrak{q})$ is also V-fit for $(\mathfrak{g}, \mathfrak{w})$, because $\mathfrak{w} \subset \mathfrak{q}$.

Let now $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W}) \subset \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ be V-fit for $(\mathfrak{g}, \mathfrak{w})$. Note that $\mathcal{R}^+(C) \cap \mathcal{Q}^r = (\mathcal{R}^+(C) \cap \mathcal{W}^r) \cup (\mathcal{W}^n \setminus \mathcal{Q}^n)$, and $\mathcal{W}^n \setminus \mathcal{Q}^n = \mathcal{Q}^r \cap \bar{\mathcal{Q}}^{-n}$. Since $\bar{\alpha} \prec_C 0$ for all $\alpha \in \bar{\mathcal{Q}}^{-n}$, we obtain that $\sigma(\mathcal{R}^+(C) \cap \mathcal{Q}^r \setminus \mathcal{R}_{\text{re}}) \subset \mathcal{R}^-(C)$ and therefore C is also V-fit for $(\mathfrak{g}, \mathfrak{q})$. $\square$

Proposition 6.5. *Let $(\mathfrak{g}, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, and let $(\mathfrak{g}, \mathfrak{w})$ be its canonical lift. Let $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ be V-fit for $(\mathfrak{g}, \mathfrak{q})$, and $\Phi_C = \mathcal{B}(C) \cap \mathcal{Q}^n$, $\tilde{\Phi}_C = \mathcal{B}(C) \cap \mathcal{W}^n$. Then :*

$$(6.8) \quad \tilde{\Phi}_C = \Phi_C \cup \{\alpha \in \mathcal{B}(C) \mid \text{supp}_C(\bar{\alpha}) \cap \Phi_C \neq \emptyset\}.$$

Proof. By Lemma 6.4, $\mathfrak{w} = \mathfrak{q}_{\tilde{\Phi}_C}$ for some set of simple roots $\tilde{\Phi}_C$ with $\Phi_C \subset \tilde{\Phi}_C \subset \mathcal{B}(C)$. Let $\mathcal{Q} = \mathcal{Q}_A = \{\alpha \in \mathcal{R} \mid \alpha(A) \geq 0\}$, with $A \in \mathfrak{h}_{\mathbb{R}}$. Fix a real $\varepsilon > 0$ with $\varepsilon |\bar{\alpha}(A)| < \alpha(A)$ for all $\alpha \in \mathcal{Q}^n$. Then :

$$(6.9) \quad \mathcal{W} = \{\alpha \in \mathcal{R} \mid \alpha(A - \varepsilon \bar{A}) \geq 0\}.$$

Indeed, $\mathcal{W} \subset \mathcal{Q}$ because $\alpha(A - \varepsilon \bar{A}) < 0$ when $\alpha \in \mathcal{R}$ and $\alpha(A) < 0$; moreover $\mathcal{Q}^n \subset \mathcal{W}^n$, and a root $\alpha \in \mathcal{Q}^r$ belongs to $\mathcal{W}$ if and only if $\bar{\alpha}(A) \leq 0$, i.e. if and only if $\mathfrak{v}(\alpha) \in \mathcal{Q}$.

Thus we have :

$$\begin{aligned} \tilde{\Phi}_C &= \mathcal{B}(C) \cap \mathcal{W}^n = \{\alpha \in \mathcal{B}(C) \mid \alpha(A - \varepsilon \bar{A}) > 0\} \\ &= \Phi_C \cup \{\alpha \in \mathcal{B}(C) \mid \alpha(A) = 0, \bar{\alpha}(A) < 0\} \\ &= \Phi_C \cup \{\alpha \in \mathcal{B}(C) \mid \text{supp}_C(\bar{\alpha}) \cap \Phi_C \neq \emptyset\}, \end{aligned}$$

because $\tilde{\Phi}_C \setminus \Phi_C \subset \mathcal{R}_{\text{cp}}$ and, for a complex α in $\mathcal{B}(C) \setminus \Phi_C$ we have $\bar{\alpha} \in \mathcal{R}^-(C)$: hence $\bar{\alpha}(A) < 0$ whenever $\bar{\alpha}(A) \neq 0$, i.e. $\text{supp}_C(\bar{\alpha}) \cap \Phi_C \neq \emptyset$. $\square$

We can slightly improve the criterion of weak non-degeneracy of Theorem 5.9, by using Weyl chambers adapted to the canonical CR lift. We have indeed :

Proposition 6.6. *Let $(\mathfrak{g}, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, and let $(\mathfrak{g}, \mathfrak{w})$ be the canonical CR-lift of $(\mathfrak{g}, \mathfrak{q})$. If $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W})$, then :*

$$(6.10) \quad (\mathfrak{g}, \mathfrak{q}) \text{ is weakly non-degenerate if and only if } \bar{\alpha} \succ_C 0 \quad \forall \alpha \in \Phi_C.$$

Proof. $(\Rightarrow)$ We argue by contradiction. Assume that $\bar{\alpha}_0 \prec_C 0$ for some $\alpha_0 \in \Phi_C$. We want to prove that $\mathcal{Q}' \cup \bar{\mathcal{Q}}' = \mathcal{Q} \cup \bar{\mathcal{Q}}$ for the parabolic set

$$(6.11) \quad \mathcal{Q}' = \mathcal{Q}_{\Psi_C} = \mathcal{Q} \cup \{\beta \in \mathcal{R} \mid \text{supp}_C(\beta) \cap \Phi_C \subset \{\alpha_0\}\},$$

corresponding to $\Psi_C = \Phi_C \setminus \{\alpha_0\} \subset \mathcal{B}(C)$. It suffices to verify that $\bar{\beta} \in \mathcal{Q}$ if $\beta \prec_C 0$ and $\text{supp}_C(\beta) \cap \Phi_C = \{\alpha_0\}$. Since $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W})$, each $\alpha \in \mathcal{B}(C)$ either belongs to $\mathcal{W}^n = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^{-n})$, or to $\mathcal{W}^r = \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r$. Thus $\mathcal{B}(C) \setminus \Phi_C \subset \bar{\mathcal{Q}}^r \cup \bar{\mathcal{Q}}^{-n}$, and, being $\bar{\alpha}_0 \prec_C 0$, we get $-\bar{\alpha} \in \mathcal{Q}$ for all $\alpha \in \text{supp}_C(\beta)$, yielding $\beta \in \bar{\mathcal{Q}}$.

$(\Leftarrow)$ Assume that $(\mathfrak{g}, \mathfrak{q})$ is weakly degenerate. Then, for some $\alpha_0 \in \Phi_C$, (6.11) defines a parabolic set $\mathcal{Q}'$ with $\mathcal{Q}' \cup \bar{\mathcal{Q}}' = \mathcal{Q} \cup \bar{\mathcal{Q}}$. In particular, $-\alpha_0 \in \bar{\mathcal{Q}}$. If $-\alpha_0 \in \bar{\mathcal{Q}}^n$, then $-\bar{\alpha}_0 \in \mathcal{Q}^n \subset \mathcal{R}^+(C)$ and $\bar{\alpha}_0 \prec_C 0$. Otherwise, $\bar{\alpha}_0 \in \mathcal{Q}^r \cap \bar{\mathcal{Q}}^n$ and, because $\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n \subset \mathcal{W}^{-n}$, the condition that $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W})$ implies that $\bar{\alpha}_0 \prec_C 0$. $\square$

Proposition 6.6 gives a way to construct the weakly non-degenerate reduction of a parabolic CR algebra :

Corollary 6.7. *Let $(\mathfrak{g}, \mathfrak{w})$ be the canonical CR-lift of the parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ and $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$. If $C \in \mathfrak{C}(\mathcal{R}, \mathcal{W}) \subset \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, $\Phi_C = \mathcal{B}(C) \cap \mathcal{Q}^n$, and*

$$(6.12) \quad \Psi_C = \{\alpha \in \Phi(C) \mid \bar{\alpha} \in \mathcal{R}^+(C)\},$$

then the $\mathfrak{g}$ -equivariant CR fibration $(\mathfrak{g}, \mathfrak{q}) \rightarrow (\mathfrak{g}, \mathfrak{q}_{\Psi_C})$ is the weakly nondegenerate reduction of $(\mathfrak{g}, \mathfrak{q})$. $\square$

Proposition 6.8. *Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, and let $(\mathfrak{g}, \mathfrak{w})$ be the canonical lift of $(\mathfrak{g}, \mathfrak{q})$. Then the natural $\mathbf{G}$ -equivariant projection $M(\mathfrak{g}, \mathfrak{w}) \rightarrow M(\mathfrak{g}, \mathfrak{q})$ is a CR fibration with totally complex connected fibers.*

Proof. Our CR manifolds are described as homogeneous spaces by the quotients $M(\mathfrak{g}, \mathfrak{q}) = \mathbf{G}/\mathbf{G}_+$ with $\mathbf{G}_+ = \mathbf{N}_{\mathbf{G}}(\mathfrak{q})$ and $M(\mathfrak{g}, \mathfrak{w}) = \mathbf{G}/\mathbf{W}_+$, where $\mathbf{W}_+ = \mathbf{N}_{\mathbf{G}}(\mathfrak{w})$.

We want to prove that every connected component of $\mathbf{G}_+$ contains an element of $\mathbf{W}_+$.

Let $\mathfrak{g}_+ = \mathfrak{g} \cap \mathfrak{q}$ be the Lie algebra of $\mathbf{G}_+$ and consider its decomposition in (4.3) of Proposition 4.3: we have $\mathfrak{g}_+ = \mathfrak{n} \oplus \mathfrak{g}_0$, where $\mathfrak{n}$ is the ideal of the nilpotent elements of the radical of $\mathfrak{g}_+$ and $\mathfrak{g}_0 = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r \cap \mathfrak{g}$ is a ϑ -invariant reductive complement of $\mathfrak{n}$ in $\mathfrak{g}_+$.

Being algebraic, $\mathbf{G}_+$ has a *Chevalley decomposition* (see [Ch, Chap.5, Sect.4]) into the semidirect product $\mathbf{N} \rtimes \mathbf{G}_0$ of the analytic subgroup with Lie algebra $\mathfrak{n}$ and of a closed Lie subgroup $\mathbf{G}_0$ with Lie algebra $\mathfrak{g}_0$.

Let $g \in \mathbf{G}_+$ and denote by Γ_g the connected component of g in $\mathbf{G}_+$. Since $\mathbf{N}$ is connected, we can as well take from the start g in $\mathbf{G}_0$, so that in particular $\text{Ad}_{\mathfrak{g}}(g)(\mathfrak{g}_0) = \mathfrak{g}_0$.

Since $\mathfrak{g}_0$ is a real form of $\mathfrak{q}^r \cap \bar{\mathfrak{q}}^r$, by complexification we obtain that $\text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{q}^r \cap \bar{\mathfrak{q}}^r) = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r$. Thus $\text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{w})$ is a parabolic complex subalgebra of $\hat{\mathfrak{g}}$ with $\mathfrak{q}^r \cap \bar{\mathfrak{q}}^r \subset \text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{w}) \subset \mathfrak{q}$. Since $\dim_{\mathbb{C}}(\text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{w})) = \dim_{\mathbb{C}}(\mathfrak{w})$, it follows from the characterization of $\mathfrak{w}$ in Proposition 6.2, that $\text{Ad}_{\hat{\mathfrak{g}}}(g)(\mathfrak{w}) = \mathfrak{w}$, and hence $g \in \mathbf{W}_+$. $\square$

Next we describe a construction that is similar to the one discussed above. We keep the notation introduced therein.

Proposition 6.9. *Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, and let :*

$$(6.13) \quad \mathfrak{v} = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \bar{\mathfrak{q}}) .$$

Then $\mathfrak{v}$ is a parabolic subalgebra of $\hat{\mathfrak{g}}$, such that :

$$(6.14) \quad \begin{cases} \mathfrak{v}^r = \bar{\mathfrak{v}}^r = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r, & \mathfrak{v}^n = \mathfrak{q}^n \oplus (\mathfrak{q}^r \cap \bar{\mathfrak{q}}^n) \supset \mathfrak{q}^n, & \mathfrak{v} = \mathfrak{v}^r \oplus \mathfrak{v}^n \subset \mathfrak{q} \\ \mathfrak{v}^n \cap \bar{\mathfrak{v}}^n = \mathfrak{q}^n \cap \bar{\mathfrak{q}}^n, & \bar{\mathfrak{v}}^r = \mathfrak{v}^r, & \mathfrak{v} \cap \bar{\mathfrak{v}} = \mathfrak{q} \cap \bar{\mathfrak{q}}. \end{cases}$$

It is uniquely determined by the condition of being the smallest complex parabolic subalgebras $\mathfrak{q}' \subset \hat{\mathfrak{g}}$ with :

$$(6.15) \quad \mathfrak{q} \cap \bar{\mathfrak{q}} \subset \mathfrak{q}' \subset \mathfrak{q} .$$

We note that the construction of $\mathfrak{v}$ is independent from the choice of the Cartan pair $(\mathfrak{g}, \mathfrak{h})$.

Proof. All Cartan subalgebras $\mathfrak{h}$ adapted to $(\mathfrak{g}, \mathfrak{q})$ are also contained in $\bar{\mathfrak{q}}$. We apply Proposition 6.1. The parabolic set corresponding to $\mathfrak{v}$ is:

$$(6.16) \quad \mathcal{V} = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}) = (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r) \cup \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n).$$

Then it is easy to verify (6.14) by using Proposition 6.1.

All complex parabolic $\mathfrak{q}' \subset \hat{\mathfrak{g}}$ that satisfy (6.15), also satisfy $\mathfrak{q}'^n \supset \mathfrak{q}^n$, because $\mathfrak{q}' \subset \mathfrak{q}$, and $\mathfrak{q}^r \cap \bar{\mathfrak{q}} \subset \mathfrak{q} \cap \bar{\mathfrak{q}}$; hence $\mathfrak{v} \subset \mathfrak{q}'$ if $\mathfrak{q}'$ satisfies (6.15). $\square$

The parabolic subalgebra $(\mathfrak{g}, \mathfrak{v})$ defined in Proposition 6.9 is called *the weakest CR model* of $(\mathfrak{g}, \mathfrak{q})$.

By Theorem 4.14, since $\mathfrak{q} \cap \bar{\mathfrak{q}} = \mathfrak{v} \cap \bar{\mathfrak{v}}$, we have:

Theorem 6.10. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and $(\mathfrak{q}, \mathfrak{v})$ its weakest CR model. Then the holomorphic projection $\hat{M}(\hat{\mathfrak{g}}, \mathfrak{v}) \rightarrow \hat{M}(\hat{\mathfrak{g}}, \mathfrak{q})$ restricts to a smooth diffeomorphism $M(\mathfrak{g}, \mathfrak{v}) \rightarrow M(\mathfrak{g}, \mathfrak{q})$. $\square$*

Lemma 6.11. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$. Let $\mathcal{Q} = \mathcal{Q}_A = \{\alpha \in \mathcal{R} \mid \alpha(A) \geq 0\}$, with $A \in \mathfrak{h}_{\mathbb{R}}$. Then the parabolic set $\mathcal{V}$ of the parabolic complex $\mathfrak{v} \subset \hat{\mathfrak{g}}$ of the weakest CR model $(\mathfrak{g}, \mathfrak{v})$ of $(\mathfrak{g}, \mathfrak{q})$ is given by:*

$$(6.17) \quad \mathcal{V} = \{\alpha \in \mathcal{R} \mid \alpha(A + \varepsilon \bar{A}) \geq 0\}$$

where ε is any positive real number with $\varepsilon|\bar{\alpha}(A)| < \alpha(A)$ for all $\alpha \in \mathcal{Q}^n$.

Proof. Since $\alpha(A + \varepsilon \bar{A}) < 0$ when $\alpha \notin \mathcal{Q}$, we have $\mathcal{Q}^n \subset \mathcal{V}^n$ and hence $\mathcal{V} \subset \mathcal{Q}$. Moreover $\alpha \in \mathcal{Q}^r$ belongs to $\mathcal{V}$ if and only if $\bar{\alpha}(A) \geq 0$, i.e. if and only if $\alpha \in \bar{\mathcal{Q}}$. $\square$

Proposition 6.12. *Let $(\mathfrak{g}, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, and let $(\mathfrak{g}, \mathfrak{w})$ be the canonical lift of $(\mathfrak{g}, \mathfrak{q})$. Then $(\mathfrak{g}, \mathfrak{w})$ coincides with its weakest CR model.*

Proof. We use the notation of Lemma 6.11. By Proposition 6.5, $\mathfrak{w} = \mathfrak{q}_B$ with $B = A - \varepsilon \bar{A} \in \mathfrak{h}_{\mathbb{R}}$ for $0 < \varepsilon < \varepsilon_0$. Then, by Lemma 6.11, the weakest CR model of $(\mathfrak{g}, \mathfrak{w})$ is $(\mathfrak{g}, \mathfrak{v}')$ where $\mathfrak{v}' = \mathfrak{q}_C$ with $C = A - \varepsilon' \bar{A} \in \mathfrak{h}_{\mathbb{R}}$, with $\varepsilon' = \frac{\varepsilon - \delta}{1 - \varepsilon \delta}$ for $0 < \delta < \delta_0$ sufficiently small, and hence $\mathfrak{v}' = \mathfrak{w}$. $\square$

We have:

Lemma 6.13. *Let $(\mathfrak{g}, \mathfrak{v})$ be the weakest CR model of $(\mathfrak{g}, \mathfrak{q})$ and $\mathfrak{h}$ a CR algebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$ (and hence also to $(\mathfrak{g}, \mathfrak{v})$). If $\mathcal{Q}, \mathcal{V}$ are the parabolic sets corresponding to $\mathfrak{q}$ and $\mathfrak{v}$ in $\mathcal{R} = \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}})$, then:*

$$(6.18) \quad \mathcal{Q}^n \cap \bar{\mathcal{Q}}^{-n} = \mathcal{V}^n \cap \bar{\mathcal{V}}^{-n}.$$

Proof. We have: $\mathcal{V}^n = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n)$ and hence $\bar{\mathcal{V}}^{-n} = \bar{\mathcal{Q}}^{-n} \cup (\bar{\mathcal{Q}}^r \cap \mathcal{Q}^{-n})$. Since:

$(\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n) \cap (\bar{\mathcal{Q}}^r \cap \mathcal{Q}^{-n}) = \emptyset, \quad (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n) \cap \bar{\mathcal{Q}}^{-n} = \emptyset, \quad \mathcal{Q}^n \cap (\bar{\mathcal{Q}}^r \cap \mathcal{Q}^{-n}) = \emptyset,$
we obtain (6.18). $\square$

Proposition 6.14. *The effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ is minimal if and only if its CR weakest CR model $(\mathfrak{g}, \mathfrak{v})$ is minimal.*

Proof. The statement follows by (6.18) and the characterization of parabolic minimal given in Lemma 5.11. $\square$

Lemma 6.15. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and $(\mathfrak{g}, \mathfrak{v})$ its weakest CR model. Then a Weyl chamber $C \in \mathcal{C}(\mathcal{R}, \mathcal{Q})$ is S-fit for $(\mathfrak{g}, \mathfrak{q})$ if and only if $C \in \mathcal{C}(\mathcal{R}, \mathcal{V})$ and is S-fit for $(\mathfrak{g}, \mathfrak{v})$.*

Proof. We have $\mathcal{C}(\mathcal{R}, \mathcal{V}) \subset \mathcal{C}(\mathcal{R}, \mathcal{Q})$ because $\mathcal{V} \subset \mathcal{Q}$. Since $\mathcal{V}^n = \mathcal{Q}^n \cup (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n)$, for $C \in \mathcal{C}(\mathcal{R}, \mathcal{V})$ and $\alpha \in \mathcal{V}^n \setminus \mathcal{Q}^n$, we get $\bar{\alpha} \in \mathcal{Q}^n \subset \mathcal{V}^n \subset \mathcal{R}^+(C)$. If moreover $C \in \mathcal{C}(\mathcal{R}, \mathcal{V})$ is S-fit for $(\mathfrak{g}, \mathfrak{v})$, then $\bar{\alpha} \succ_C 0$ also for $\alpha \in (\mathcal{V}^r \cap \mathcal{R}^+(C)) \setminus \mathcal{R}_{\text{im}}$; since $\mathcal{Q}^r \cap \mathcal{R}^+(C) = (\mathcal{V}^r \cap \mathcal{R}^+(C)) \cup (\mathcal{V}^n \setminus \mathcal{Q}^n)$, it follows that C is S-fit also for $(\mathfrak{g}, \mathfrak{q})$.

To complete the proof, it suffices to show that an S-fit Weyl chamber for $(\mathfrak{g}, \mathfrak{q})$ is admissible for $(\mathfrak{g}, \mathfrak{v})$. Let C be S-fit for $(\mathfrak{g}, \mathfrak{q})$. The elements α of $\mathcal{V}^n \setminus \mathcal{Q}^n$ belong to $\mathcal{Q}^r \cap \bar{\mathcal{Q}}^n$: this implies that $\bar{\alpha} \succ_C 0$ and therefore that $\alpha \succ_C 0$, by the assumption that C is S-fit for $(\mathfrak{g}, \mathfrak{q})$. Hence $\mathcal{V}^n \subset \mathcal{R}^+(C)$ and therefore $C \in \mathcal{C}(\mathcal{R}, \mathcal{V})$ when $C \in \mathcal{C}(\mathcal{R}, \mathcal{Q})$ is S-fit for $(\mathfrak{g}, \mathfrak{q})$. $\square$

Proposition 6.16. *Let $(\mathfrak{g}, \mathfrak{q})$ be a parabolic CR algebra, with $\mathfrak{q} = \mathfrak{q}_{\Phi_C}$ for an S-fit Weyl chamber $C \in \mathcal{C}(\mathcal{R}, \mathcal{Q})$. Then its weakest CR model is $(\mathfrak{g}, \mathfrak{q}_{\Phi_C^\#})$, for*

$$(6.19) \quad \Phi_C^\# = \Phi_C \cup \{\beta \in \mathcal{B}(C) \cap \mathcal{R}_{\text{cp}} \mid \text{supp}_C(\bar{\beta}) \cap \Phi_C \neq \emptyset\}.$$

Proof. Let $C \in \mathcal{C}(\mathcal{R}, \mathcal{Q})$ be S-fit for $(\mathfrak{g}, \mathfrak{q})$. If $\text{Cal} Q = \mathcal{Q}_A$ for $A \in \mathfrak{h}_{\mathbb{R}}$, then $\Phi_C = \{\beta \in \mathcal{B}(C) \mid \beta(A) > 0\}$ and, by Lemma 6.11 and Lemma 6.15, $C \in \mathcal{C}(\mathcal{Q}, \mathcal{V})$ and thus $\mathfrak{v} = \mathfrak{q}_{\Phi_C^\#}$ for $\Phi_C^\# = \{\beta \in \mathcal{B}(C) \mid \beta(A) + \varepsilon \bar{\beta}(A) > 0\}$ with $\varepsilon > 0$ and sufficiently small, yielding the characterization in the statement of the Proposition. $\square$

From Corollary 5.10 we obtain:

Corollary 6.17. *Let $(\mathfrak{g}, \mathfrak{v})$ be the weakest CR model of a parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$. Then:*

- (i) $(\mathfrak{g}, \mathfrak{v})$ is either totally real, or weakly degenerate.
- (ii) If $(\mathfrak{g}, \mathfrak{q})$ is weakly non-degenerate, then $\mathfrak{v} = \mathfrak{q}$ if and only if $(\mathfrak{g}, \mathfrak{q})$ is totally real. $\square$

By using Corollary 6.17, starting from a parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, we can construct a chain of parabolic CR algebras $(\mathfrak{g}, \mathfrak{q}_h)$, $(\mathfrak{g}, \mathfrak{v}_h)$ and $\mathfrak{g}$ -equivariant CR homomorphisms:

$$(6.20) \quad \begin{array}{ccc} & (\mathfrak{g}, \mathfrak{q}) & \\ & \downarrow & \\ (\mathfrak{g}, \mathfrak{q}_1) & \longleftarrow & (\mathfrak{g}, \mathfrak{v}_1) \\ & \downarrow & \\ & (\mathfrak{g}, \mathfrak{q}_2) \longleftarrow (\mathfrak{g}, \mathfrak{v}_2) & \\ & \downarrow & \\ & (\mathfrak{g}, \mathfrak{q}_3) \longleftarrow (\mathfrak{g}, \mathfrak{v}_3) & \\ & \vdots & \end{array}$$

where each vertical arrow is a weakly nondegenerate reduction and each horizontal arrow is a lifting to the weakest CR model.

If we denote by $\mathfrak{P}(\hat{\mathfrak{g}})$ the set of all parabolic complex Lie subalgebras of $\hat{\mathfrak{g}}$, and set $\mathfrak{v}_0 = \mathfrak{g}_{-1} = \mathfrak{g}$, we have, for all integers $h > 0$:

$$(6.21) \quad \begin{cases} \mathfrak{q}_h = \text{the largest } \mathfrak{a} \in \mathfrak{P}(\hat{\mathfrak{g}}) \text{ such that } \mathfrak{v}_{h-1} \subset \mathfrak{a} \subset \mathfrak{v}_{h-1} + \bar{\mathfrak{v}}_{h-1}, \\ \mathfrak{v}_h = \text{the smallest } \mathfrak{a} \in \mathfrak{P}(\hat{\mathfrak{g}}) \text{ such that } \mathfrak{q}_h \cap \bar{\mathfrak{q}}_h \subset \mathfrak{a} \subset \mathfrak{q}_h. \end{cases}$$

This characterization shows that the construction in (6.20) is uniquely determined and independent of the choices of the adapted Cartan pairs in $(\mathfrak{g}, \mathfrak{q}_h)$ and $(\mathfrak{g}, \mathfrak{v}_h)$.

We know that the $\mathbf{G}$ -equivariant maps $M(\mathfrak{g}, \mathfrak{v}_h) \rightarrow M(\mathfrak{g}, \mathfrak{q}_h)$ (for $h > 0$) are smooth diffeomorphisms, while the $\mathbf{G}$ -equivariant maps $M(\mathfrak{g}, \mathfrak{v}_h) \rightarrow M(\mathfrak{g}, \mathfrak{q}_{h+1})$ (for $h \geq 0$) are CR fibrations. In particular, there is a smallest integer $m \geq -1$ such that $\dim_{\mathbb{R}} M(\mathfrak{g}, \mathfrak{v}_h) > \dim_{\mathbb{R}} M(\mathfrak{g}, \mathfrak{v}_{h+1})$ for $h < m+1$, and $M(\mathfrak{g}, \mathfrak{v}_h) = M(\mathfrak{g}, \mathfrak{v}_{m+1}) = M(\mathfrak{g}, \mathfrak{q}_m)$ for all $h > m+1$. Hence, by the characterization of the fibers in Theorem 4.9, we have:

Proposition 6.18. *With the notation above: let $(\mathfrak{g}, \mathfrak{q}_h)$, $(\mathfrak{g}, \mathfrak{v}_h)$ be the sequence of weakly nondegenerate parabolic CR algebra and of their weakest CR models defined above. Then there exists a smallest integer $m \geq 0$ such that $\mathfrak{q}_h = \mathfrak{v}_h = \mathfrak{q}_m$ for all $h > m$. Moreover, $(\mathfrak{g}, \mathfrak{q}_m)$ is totally real and m is the smallest nonnegative integer for which $(\mathfrak{g}, \mathfrak{q}_m)$ is totally real.*

By composition, we obtain a $\mathbf{G}$ -equivariant fibration $M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}_m)$ where the basis $M(\mathfrak{g}, \mathfrak{q}_m)$ is a totally real parabolic CR manifold and each connected component of the fiber is a Cartesian product of Euclidean complex nilmanifolds and of simply connected totally complex parabolic CR manifolds. $\square$

Example 6.1. Let $\varepsilon_1, \dots, \varepsilon_6$ be the canonical basis of $\mathbb{R}^6 \subset \mathbb{C}^6$. Let $\mathbf{G} = \mathbf{SL}(6, \mathbb{R})$ consist of the matrices of $\hat{\mathbf{G}} = \mathbf{SL}(6, \mathbb{C})$ which have real entries in the canonical basis. We consider in $\mathbb{C}^6$ the basis:


$$e_1 = \varepsilon_1 + i\varepsilon_4, \quad e_2 = \varepsilon_2 + i\varepsilon_5, \quad e_3 = \varepsilon_3 + i\varepsilon_6, \quad e_4 = \varepsilon_1 - i\varepsilon_4, \quad e_5 = \varepsilon_2 - i\varepsilon_5, \quad e_6 = \varepsilon_3 - i\varepsilon_6$$

and we want to investigate the $\mathbf{G}$ -orbit M of the flag

$$\langle e_1 \rangle \subset \langle e_1, e_2 \rangle \subset \langle e_1, e_2, e_3 \rangle \subset \langle e_1, e_2, e_3, e_4 \rangle \subset \langle e_1, e_2, e_3, e_4, e_5 \rangle.$$


Let $(\mathfrak{g}, \mathfrak{q})$, with $\mathfrak{g} = \mathfrak{sl}(6, \mathbb{R})$, be the associated parabolic CR algebra. We consider the Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{sl}(6, \mathbb{R})$ that is represented, in the basis $e_1, \dots, e_6$, by the diagonal matrices $\text{diag}(\lambda_1, \lambda_2, \lambda_3, \bar{\lambda}_1, \bar{\lambda}_2, \bar{\lambda}_3)$ with $\text{Re}(\lambda_1 + \lambda_2 + \lambda_3) = 0$. Then $\mathfrak{h}_{\mathbb{R}}$ consists of the 6×6 traceless real diagonal matrices.

The cross-marked diagram associated to $(\mathfrak{g}, \mathfrak{q})$ in the adapted Weyl chamber C with simple roots $\mathcal{B}(C) = \{\alpha_i = e_1 - e_{i+1}, 1 \leq i \leq 5\}$, is the following:




Since $\mathfrak{q}$ is a complex Borel subalgebra of $\hat{\mathfrak{g}}$, the chamber C is S-fit and then $(\mathfrak{g}, \mathfrak{q})$ is fundamental by Theorem 5.5, because $\bar{\alpha}_3 = -(\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5)$. The chamber C is also V-fit, and we obtain the weakly non-degenerate basis $(\mathfrak{g}, \mathfrak{q}_1)$ by


dropping the cross under the simple root α_3 with $\bar{\alpha}_3 \prec_C 0$. The diagram associated to $(\mathfrak{g}, \mathfrak{q}_1)$ is:



The parabolic $\mathfrak{q}_1$ is defined by the element $A_1 = \text{diag}(2, 1, 0, 0, -1, -2) \in \mathfrak{h}_{\mathbb{R}}$. To compute its weakest CR model $(\mathfrak{g}, \mathfrak{v}_1)$, we observe that $\bar{A}_1 = \text{diag}(0, -1, -2, 2, 1, 0)$, so that $A_1 + \varepsilon \bar{A}_1 = \text{diag}(2, 1 - \varepsilon, -2\varepsilon, 2\varepsilon, -1 + \varepsilon, -2)$. To take a Weyl chamber adapted to $(\mathfrak{g}, \mathfrak{v}_1)$, it is convenient to consider the basis obtained from $e_1, \dots, e_6$ by reordering its elements according to the decreasing ordering of the diagonal entries of $A_1 + \varepsilon \bar{A}_1$. We obtain the new basis: $e_1, e_2, e_4, e_3, e_5, e_6$. With $\alpha'_1 = e_1 - e_2$, $\alpha'_2 = e_2 - e_4$, $\alpha'_3 = e_4 - e_3$, $\alpha'_4 = e_3 - e_5$, $\alpha'_5 = e_5 - e_6$ being the simple roots a Weyl chamber $C' \in \mathfrak{C}(\mathcal{R}, \mathcal{V}_1)$, we obtain the diagram:




The weakly nondegenerate reduction $(\mathfrak{g}, \mathfrak{q}_2)$ of $(\mathfrak{g}, \mathfrak{v}_1)$ has the diagram:




We have $\mathfrak{q}_2 = \mathfrak{q}_{A_2}$, with

$$A_2 = \text{diag}(2, 1, -1, 1, -1, -2), \quad \bar{A}_2 = \text{diag}(1, -1, -2, 2, 1, -1),$$

so that $A_2 + \varepsilon \bar{A}_2 = \text{diag}(2 + \varepsilon, 1 - \varepsilon, -1 - 2\varepsilon, 1 + 2\varepsilon, -1 + \varepsilon, -2 - \varepsilon)$. To describe $(\mathfrak{g}, \mathfrak{v}_2)$, for $\mathfrak{v}_2 = \mathfrak{q}_{A_2 + \varepsilon \bar{A}_2}$, it is convenient to consider the Weyl chamber $C'' \in \mathfrak{C}(\mathcal{R}, \mathcal{V}_2)$ that corresponds to the simple roots related to the ordered basis $e_1, e_4, e_2, e_5, e_3, e_6$ of $\mathbb{C}^6$, for which the entries of $A_2 + \varepsilon \bar{A}_2$ are decreasing: with $\alpha''_1 = e_1 - e_4$, $\alpha''_2 = e_4 - e_2$, $\alpha''_3 = e_2 - e_5$, $\alpha''_4 = e_5 - e_3$, $\alpha''_5 = e_3 - e_6$, we obtain for $(\mathfrak{g}, \mathfrak{v}_2)$ the diagram:



Since $\mathfrak{v}_2$ is Borel, C'' is V-fit for $(\mathfrak{g}, \mathfrak{v}_2)$ and the diagram of the weakly nondegenerate reduction $(\mathfrak{g}, \mathfrak{q}_3)$ of $(\mathfrak{g}, \mathfrak{v}_2)$ is obtained by dropping the crosses under the α''_i 's with $\bar{\alpha}''_i \prec_{C''} 0$:



The parabolic CR algebra $(\mathfrak{g}, \mathfrak{q}_3)$ is totally real, as $\mathfrak{q}_3 = \mathfrak{q}_{A_3}$ with

$$A_3 = \text{diag}(1, 0, -1, 1, 0, -1) = \bar{A}_3.$$

Hence $(\mathfrak{g}, \mathfrak{q}_2) = (\mathfrak{g}, \mathfrak{v}_h) = (\mathfrak{g}, \mathfrak{q}_h)$ for all $h \geq 3$. The map $M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}_3)$ is given by $(\ell_1, \ell_2, \ell_3, \ell_4, \ell_5) \rightarrow (\ell_1 + \bar{\ell}_1, \ell_2 + \bar{\ell}_2)$.

7. THE ISOTROPY SUBGROUPS

Let $\mathfrak{g}$ be a real semisimple Lie algebra, $\hat{\mathfrak{g}}$ its complexification, ϑ a Cartan involution of $\mathfrak{g}$. Let $\mathfrak{h}$ be a ϑ -invariant Cartan subalgebra of $\mathfrak{g}$ and $\mathcal{R} = \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}})$ the root system of $\hat{\mathfrak{g}}$ with respect to the complexification $\hat{\mathfrak{h}}$ of $\mathfrak{h}$. Denote by $\Lambda(\mathcal{R})$ the additive subgroup of $\mathfrak{h}_{\mathbb{R}}^*$ generated by $\mathcal{R}$ and by $\Pi(\mathcal{R})$ the *lattice of weights*, consisting of all $\eta \in \mathfrak{h}_{\mathbb{R}}^*$ for which $(\eta|\alpha^\vee) = 2(\eta|\alpha)/\|\alpha\|^2 \in \mathbb{Z}$. We have $\Lambda(\mathcal{R}) \subset \Pi(\mathcal{R})$.

Given a lattice (i.e. a free Abelian group) $\mathcal{L}$, a *character* of $\mathcal{L}$ is a homomorphism $\chi : \mathcal{L} \rightarrow \mathbb{C}^*$ of $\mathcal{L}$ into the multiplicative group $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ of non-zero complex numbers. If $b_1, \dots, b_\ell \in \mathcal{L}$ is a basis of $\mathcal{L}$ over $\mathbb{Z}$, a character $\chi \in \text{Hom}(\mathcal{L}, \mathbb{C}^*)$ is completely determined by its values $\lambda_i = \chi(b_i)$ (for $1 \leq i \leq \ell$) on the basis, so that $\text{Hom}(\mathcal{L}, \mathbb{C}^*) \simeq [\mathbb{C}^*]^\ell$.

We keep the notation of the previous sections, in particular $\hat{\mathbf{G}}$ is a connected and simply connected complex Lie group with Lie algebra $\hat{\mathfrak{g}}$ and $\mathbf{G}$ its analytic subgroup with Lie algebra $\mathfrak{g}$. It is a covering group of any linear group with Lie algebra $\mathfrak{g}$. Let $\hat{\mathbf{H}}$ be the Cartan subgroup of $\hat{\mathbf{G}}$ corresponding to $\hat{\mathfrak{h}}$:

$$(7.1) \quad \hat{\mathbf{H}} = \mathbf{Z}_{\hat{\mathbf{G}}}(\hat{\mathfrak{h}}) = \{z \in \hat{\mathbf{G}} \mid \text{Ad}_{\hat{\mathfrak{g}}}(z)(H) = H, \forall H \in \hat{\mathfrak{h}}\}.$$

All finite dimensional $\mathbb{C}$ -linear representations of the complex semisimple Lie algebra $\hat{\mathfrak{g}}$ are differentials of representations of the complex Lie group $\hat{\mathbf{G}}$. Each element h of $\hat{\mathbf{H}}$ defines a character $\chi_h \in \text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*)$, and vice versa, a character $\chi \in \text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*)$ defines the element $h_\chi \in \hat{\mathbf{H}}$. To explain this correspondence, take first a faithful representation $\rho : \hat{\mathbf{G}} \hookrightarrow \mathbf{SL}_{\mathbb{C}}(V)$, corresponding to $\rho : \hat{\mathfrak{g}} \hookrightarrow \mathfrak{sl}_{\mathbb{C}}(V)$, for a finite dimensional complex linear space V , and define $\rho(h_\chi)(v) = \chi(\omega)v$ for $v \in V^\omega = \{v \in V \mid \rho(H)(v) = \omega(H)v, \forall H \in \hat{\mathfrak{h}}\}$, for $\omega \in \Pi(\mathcal{R})$. The Cartan subgroup $\hat{\mathbf{H}}$ is analytic and $\exp : \hat{\mathfrak{h}} \rightarrow \hat{\mathbf{H}}$ is onto, so that the correspondence with $\text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*)$ can also be described by $\chi_{\exp(H)}(\omega) = \exp(\omega(H))$ for $H \in \hat{\mathfrak{h}}$. With $\ell = \dim_{\mathbb{C}}(\hat{\mathfrak{h}}) = \text{rank of } \hat{\mathfrak{g}}$ we have:

$$(7.2) \quad \hat{\mathbf{H}} = \mathbf{Z}_{\hat{\mathbf{G}}}(\hat{\mathfrak{h}}) = \{h_\chi \mid \chi \in \text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*)\} \simeq \text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*) \simeq [\mathbb{C}^*]^\ell.$$

The Cartan subgroup $\mathbf{H}$ of $\mathbf{G}$ corresponding to $\mathfrak{h}$ is the centralizer of $\mathfrak{h}$ in $\mathbf{G}$:

$$(7.3) \quad \mathbf{H} = \mathbf{Z}_{\mathbf{G}}(\mathfrak{h}) = \{h \in \mathbf{G} \mid \text{Ad}_{\mathfrak{g}}(h)(H) = H, \forall H \in \mathfrak{h}\}.$$

For a lattice $\mathcal{L} \subset \mathfrak{h}_{\mathbb{R}}^*$, with $\sigma(\mathcal{L}) = \mathcal{L}$, we set

$$\text{Hom}^\sigma(\mathcal{L}, \mathbb{C}^*) = \{\chi \in \text{Hom}(\mathcal{L}, \mathbb{C}^*) \mid \chi(\bar{\eta}) = \overline{\chi(\eta)}, \forall \eta \in \mathcal{L}\}.$$

We obtain:

Lemma 7.1. *Let $\mathfrak{g}$ be a real semisimple Lie algebra, $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$. Then:*

$$(7.4) \quad \mathbf{H} = \mathbf{Z}_{\mathbf{G}}(\mathfrak{h}) = \{h_\chi \mid \chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*)\}.$$

Proof. The action of σ on $\text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*)$, given by $\sigma(\chi)(\eta) = \overline{\chi(\bar{\eta})}$ coincides, under the correspondence (7.1), with the action of σ on $\hat{\mathbf{H}}$. Then the statement follows from the fact that $\mathbf{H} = \hat{\mathbf{H}} \cap \mathbf{G}$ and $\mathbf{H} = \hat{\mathbf{H}}^\sigma$ because of (4.2). $\square$

In view of the preceding Lemma, we now give a more explicit description of $\text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*)$. Fix a Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$ that is S-adapted to the conjugation σ . Set $\mathcal{B}(C) = \{\beta_1, \dots, \beta_a\} \cup \{\tau_1, \dots, \tau_b\}$, with $\{\beta_1, \dots, \beta_a\} = \mathcal{B}(C) \setminus \mathcal{R}_{\text{im}}$ and $\{\tau_1, \dots, \tau_b\} = \mathcal{B}(C) \cap \mathcal{R}_{\text{im}}$. By Theorem 5.1, the conjugation σ is described in $\mathcal{B}(C)$ by an involutive permutation $j : \beta_i \rightarrow j(\beta_i) = \beta_{i'}$ of $\{\beta_1, \dots, \beta_a\}$ and a matrix of nonnegative integers $(k_{i,q})$, $1 \leq i \leq a$, $1 \leq q \leq b$, such that :

$$(7.5) \quad \begin{cases} \bar{\beta}_i = \beta_{i'} + \sum_{q=1}^b k_{i,q} \tau_q, \\ j(\beta_i) = \beta_{i'}, \quad k_{i',p} = k_{i,p} \quad \text{for } 1 \leq i \leq a, \quad 1 \leq p \leq b. \end{cases}$$

In $\Pi(\mathcal{R})$ we consider the basis $\mathcal{B}^*(C) = \{\omega_1, \dots, \omega_a, \theta_1, \dots, \theta_b\}$, adjoint of $\mathcal{B}(C)$, defined by :

$$(7.6) \quad \begin{cases} (\omega_i | \beta_j^\vee) = \delta_{i,j} & (\omega_i | \tau_q^\vee) = 0 \\ (\theta_p | \beta_j^\vee) = 0 & (\theta_p | \tau_q^\vee) = \delta_{p,q} \end{cases} \quad \text{for } 1 \leq i, j \leq a, \quad 1 \leq p, q \leq b.$$

The conjugation σ is described in $\mathcal{B}^*(C)$ by :

$$(7.7) \quad \begin{cases} \bar{\omega}_i = \omega_{i'} & \text{for } 1 \leq i, i' \leq a, \quad j(\beta_i) = \beta_{i'} \\ \bar{\theta}_p = -\theta_p + \sum_{j=1}^a k'_{j,p} \omega_j & \text{for } 1 \leq p \leq b, \end{cases}$$

where $k'_{j,p} = k_{j,p} \|\tau_p\|^2 / \|\beta_j\|^2$.

The characters $\chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*)$ are those satisfying :

$$(7.8) \quad \begin{cases} \chi(\omega_i) \in \mathbb{C}^*, \quad \chi(\theta_p) \in \mathbb{C}^*, & \text{for } 1 \leq i \leq a, \quad 1 \leq p \leq b, \\ \overline{\chi(\omega_i)} = \chi(\omega_{i'}) & \text{for } 1 \leq i, i' \leq a, \quad j(\beta_i) = \beta_{i'} \\ |\chi(\theta_p)|^2 = \prod_{j=1}^a [\chi(\omega_j)]^{k'_{j,p}} & \text{for } 1 \leq p \leq b. \end{cases}$$

Each lattice $\mathcal{L}$ in $\mathfrak{h}_{\mathbb{R}}^*$, with $\Lambda(\mathcal{R}) \subset \mathcal{L} \subset \Pi(\mathcal{R})$, is the set of weights of all finite dimensional linear representations of an essentially unique connected complex semisimple Lie group $\hat{\mathbf{G}}_{\mathcal{L}}$. For instance, we have $\hat{\mathbf{G}}_{\Pi(\mathcal{R})} = \hat{\mathbf{G}}$ (the simply connected complex Lie group with Lie algebra $\hat{\mathfrak{g}}$) and $\hat{\mathbf{G}}_{\Lambda(\mathcal{R})} = \mathbf{Int}_{\mathbb{C}}(\hat{\mathfrak{g}})$ (the group of inner automorphisms of the complex semisimple Lie algebra $\hat{\mathfrak{g}}$) (see e.g. [OV, Ch.3, Theorem 2.11]). When moreover $\sigma(\mathcal{L}) = \mathcal{L}$, the analytic Lie subgroup of $\hat{\mathbf{G}}_{\mathcal{L}}$ with Lie algebra $\mathfrak{g}$ is a real form $\mathbf{G}_{\mathcal{L}}$ of $\hat{\mathbf{G}}_{\mathcal{L}}$. Vice versa, every connected linear semisimple Lie group with Lie algebra $\mathfrak{g}$ can be obtained in this way. The Cartan subgroup :

$$(7.9) \quad \mathbf{H}_{\mathcal{L}} = \mathbf{Z}_{\mathbf{G}_{\mathcal{L}}}(\mathfrak{h})$$

of $\mathbf{G}_{\mathcal{L}}$, relative to $\mathfrak{h}$, is a real Lie subgroup of the complex Cartan subgroup :

$$(7.10) \quad \hat{\mathbf{H}}_{\mathcal{L}} = \mathbf{Z}_{\hat{\mathbf{G}}_{\mathcal{L}}}(\hat{\mathfrak{h}}) = \{h_{\chi} \mid \chi \in \text{Hom}(\mathcal{L}, \mathbb{C}^*)\}$$

of $\hat{\mathbf{G}}_{\mathcal{L}}$, relative to $\hat{\mathfrak{h}}$. From Lemma 7.1 we obtain :

Corollary 7.2. *Let $\mathcal{L}$ be a lattice in $\mathfrak{h}_{\mathbb{R}}^*$ with $\Lambda(\mathcal{R}) \subset \mathcal{L} \subset \Pi(\mathcal{R})$ and $\sigma(\mathcal{L}) = \mathcal{L}$. Denote by $\chi \rightarrow \chi^b$ the restriction homomorphism $\text{Hom}(\Pi(\mathcal{R}), \mathbb{C}^*) \rightarrow \text{Hom}(\mathcal{L}, \mathbb{C}^*)$. Then :*

$$(7.11) \quad \mathbf{H}_{\mathcal{L}} = \{h_{\chi^b} \mid \chi \in \text{Hom}^{\sigma}(\Pi(\mathcal{R}), \mathbb{C}^*)\}.$$

Proof. The covering map $\mathbf{G} \rightarrow \mathbf{G}_{\mathcal{L}}$ transforms the Cartan subgroup $\mathbf{H}$ of $\mathbf{G}$ into the Cartan subgroup $\mathbf{H}_{\mathcal{L}}$ of $\mathbf{G}_{\mathcal{L}}$. $\square$

From Corollary 7.2 we obtain the exact sequence :

$$(7.12) \quad \mathbf{1} \rightarrow \text{Hom}^{\sigma}(\Pi(\mathcal{R})/\mathcal{L}, \mathbb{C}^*) \rightarrow \text{Hom}^{\sigma}(\Pi(\mathcal{R}), \mathbb{C}^*) \rightarrow \mathbf{H}_{\mathcal{L}} \rightarrow \mathbf{1}.$$

We can utilize (7.12) to compute the group $\pi_0(\mathbf{H}_{\mathcal{L}})$ of the connected components of the Cartan subgroup $\mathbf{H}_{\mathcal{L}}$. We have indeed :

Theorem 7.3. *Keep the previous notation. Let $C \in \mathfrak{C}(\mathcal{R})$ be S -adapted to the conjugation σ . Denote by $\omega_1, \dots, \omega_a$ the weights in $\mathcal{B}^*(C)$ that vanish on $\mathcal{B}(C) \cap \mathcal{R}_{\text{im}}$, and by $\{\theta_1, \dots, \theta_b\}$ those vanishing on $\mathcal{B}(C) \setminus \mathcal{R}_{\text{im}}$. By reordering, we assume that $\{\omega_1, \dots, \omega_c\}$ is the set of weights in $\mathcal{B}^*(C)$ with $\bar{\omega}_i = \omega_i$. Define the non negative integers $k'_{i,p}$, for $1 \leq i \leq a$, $1 \leq p \leq b$, by $\bar{\theta}_p = -\theta_p + \sum_{i=1}^a k'_{i,p} \omega_i$. Consider the subgroups of the free Abelian group $\mathbb{Z}_2^c$:*

$$(7.13) \quad \begin{aligned} \mathbf{A} &= \{(\eta_1, \dots, \eta_c) \in \mathbb{Z}_2^c \mid \sum_{i=1}^c k'_{i,p} \eta_i \equiv 0 \pmod{2}, \forall 1 \leq p \leq b\}, \\ \mathbf{A}_{\mathcal{L}} &= \{(\eta_1, \dots, \eta_c) \in \mathbf{A} \mid \sum_{i=1}^c k_i \eta_i \equiv 0 \pmod{2}, \text{ if } \sum_{i=1}^c k_i \omega_i \in \mathcal{L}\}. \end{aligned}$$

Then :

$$(7.14) \quad \pi_0(\mathbf{H}_{\mathcal{L}}) \cong \mathbf{A}/\mathbf{A}_{\mathcal{L}}.$$

Proof. The exact sequence (7.12) yields the exact sequence for the groups of the connected components :

$$\pi_0(\text{Hom}^{\sigma}(\Pi(\mathcal{R})/\mathcal{L}, \mathbb{C}^*)) \rightarrow \pi_0(\text{Hom}^{\sigma}(\Pi(\mathcal{R}), \mathbb{C}^*)) \rightarrow \pi_0(\mathbf{H}_{\mathcal{L}}) \rightarrow \mathbf{1}.$$

The statement follows because $\pi_0(\text{Hom}^{\sigma}(\Pi(\mathcal{R}), \mathbb{C}^*)) \cong \mathbf{A}$, and, in this isomorphism, the image of $\pi_0(\text{Hom}^{\sigma}(\Pi(\mathcal{R})/\mathcal{L}, \mathbb{C}^*))$ in $\pi_0(\text{Hom}^{\sigma}(\Pi(\mathcal{R}), \mathbb{C}^*))$ is identified with $\mathbf{A}_{\mathcal{L}}$. $\square$

Example 7.1. Consider the B_2 system of roots $\mathcal{R} = \{\pm e_1 \pm e_2\} \cup \{\pm e_1, \pm e_2\} \subset \mathbb{R}^2$, with σ defined by $\sigma(e_1) = e_2$ and $\sigma(e_2) = e_1$. In the Weyl chamber C with simple roots $\{e_1 - e_2, e_2\}$, that is S -adapted to σ , it can be represented by the diagram :

$$\begin{array}{ccc} \circledast & \xrightarrow{\quad\quad\quad} & \circ \\ \tau & & \beta \end{array}$$

We have, for the adjoint basis, $\theta = e_1$ and $\omega = (e_1 + e_2)/2$. We obtain $\bar{\theta} = -\theta + 2\omega$ and $\bar{\omega} = \omega$. Then $k' = 2$, $m = 1$, and $\mathbf{A} = \mathbb{Z}_2$. The quotient $\Pi(\mathcal{R})/\Lambda(\mathcal{R})$ is a group of order 2, with the generator $[\omega]$ and $2[\omega] \equiv 0$. Hence $\pi_0(\mathbf{H}) \simeq \mathbb{Z}_2$, while $\pi_0(\mathbf{H}_{\Lambda(\mathcal{R})}) \cong \mathbb{Z}_2/\mathbb{Z}_2 \cong \mathbf{1}$.

Let $\mathbf{G}_{\mathcal{L}}$ be any connected real linear Lie group with Lie algebra $\mathfrak{g}$, and $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$. The Cartan subgroup $\mathbf{H}_{\mathcal{L}}$ is a normal subgroup of the normalizer

$$(7.15) \quad \mathbf{N}_{\mathcal{L}} = \mathbf{N}_{\mathbf{G}_{\mathcal{L}}}(\mathfrak{h}) = \{g \in \mathbf{G}_{\mathcal{L}} \mid \text{Ad}(g)(\mathfrak{h}) = \mathfrak{h}\}$$

of $\mathfrak{h}$ in $\mathbf{G}_{\mathcal{L}}$. The quotient

$$(7.16) \quad \mathbf{W}_{\mathcal{L}} = \mathbf{N}_{\mathcal{L}}/\mathbf{H}_{\mathcal{L}}$$

is the *analytic Weyl group* corresponding to $\mathfrak{h}$ and $\mathcal{L}$.

We know (see e.g. [Wa, Ch.1,§4,p.115]) that actually the *analytic Weyl group* only depends, modulo natural isomorphisms, upon the real Lie algebra $\mathfrak{g}$ and its Cartan subalgebra $\mathfrak{h}$; to stress this fact, we shall write $\mathbf{W}(\mathfrak{g}, \mathfrak{h})$ instead of $\mathbf{W}_{\mathcal{L}}$.

Let us consider now an effective parabolic *CR* algebra $(\mathfrak{g}, \mathfrak{q})$.

We keep the notation introduced in previous sections, in particular we fix a Cartan pair $(\vartheta, \mathfrak{h})$ adapted to $(\mathfrak{g}, \mathfrak{q})$.

We have the decomposition

$$(7.17) \quad \mathfrak{g}_+ = \mathfrak{q} \cap \mathfrak{g} = \mathfrak{n} \oplus \mathfrak{g}_0 = \mathfrak{r} \oplus \mathfrak{s}_0 = \mathfrak{n} \oplus \mathfrak{z}_0 \oplus \mathfrak{s}_0$$

where :

$\mathfrak{r} = \mathfrak{n} \oplus \mathfrak{z}_0$ is the radical of $\mathfrak{g}_+$,

$\mathfrak{n}$ is the ideal of the $\text{ad}_{\mathfrak{g}}$ -nilpotent elements of the radical of $\mathfrak{g}_+$,

$\mathfrak{z}_0$ is a maximal Abelian subalgebra of ad-semisimple elements of $\mathfrak{r}$,

$\mathfrak{g}_0 = \mathfrak{z}_0 \oplus \mathfrak{s}_0$ is the ϑ -invariant reductive complement of $\mathfrak{n}$ in $\mathfrak{g}_+$,

$\mathfrak{s}_0 = [\mathfrak{g}_0, \mathfrak{g}_0]$ is the semisimple ideal and $\mathfrak{z}_0$ is the center of $\mathfrak{g}_0$.

The Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$, contained in $\mathfrak{g}_+$, decomposes into the direct sum :

$$(7.18) \quad \mathfrak{h} = \mathfrak{z}_0 \oplus \mathfrak{h}_0$$

where $\mathfrak{h}_0 = \mathfrak{h} \cap \mathfrak{s}_0$ is a Cartan subalgebra of $\mathfrak{s}_0$. The subalgebra $\mathfrak{z}_0$ is characterized by :

$$(7.19) \quad \mathfrak{z}_0 = \{H \in \mathfrak{h} \mid \alpha(H) = 0, \forall \alpha \in \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r\}.$$

Indeed its complexification $\hat{\mathfrak{z}}_0$ is the center of the reductive complex Lie algebra $\mathfrak{q}^r \cap \bar{\mathfrak{q}}^r$.

The isotropy subgroup

$$(7.20) \quad \mathbf{G}_+ = \mathbf{N}_{\mathbf{G}}(\mathfrak{q}) = \{g \in \mathbf{G} \mid \text{Ad}(g)(\mathfrak{q}) = \mathfrak{q}\}$$

is a closed real-algebraic subgroup of $\mathbf{G}$, with Lie algebra $\mathfrak{g}_+$. Thus we have the Chevalley decomposition :

$$(7.21) \quad \mathbf{G}_+ = \mathbf{G}_0 \rtimes \mathbf{N}$$

where $\mathbf{G}_0$ is a closed, real-algebraic, reductive subgroup of $\mathbf{G}$, with Lie algebra $\mathfrak{g}_0$, and $\mathbf{N}$ is a unipotent closed connected and simply connected subgroup of $\mathbf{G}$, diffeomorphic to its Lie algebra $\mathfrak{n}$.

We also define $\mathbf{S}_0$ to be the analytic semisimple Lie subgroup of $\mathbf{G}$ with Lie algebra $\mathfrak{s}_0$.

Proposition 7.4. *The group $\mathbf{G}_0$ is the normalizer in $\mathbf{G}$ of $\mathfrak{g}_0$ and the centralizer in $\mathbf{G}$ of $\mathfrak{z}_0$:*

$$(7.22) \quad \begin{aligned} \mathbf{G}_0 &= \mathbf{N}_{\mathbf{G}}(\mathfrak{g}_0) = \{g \in \mathbf{G} \mid \text{Ad}(g)(\mathfrak{g}_0) = \mathfrak{g}_0\} \\ &= \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) = \{g \in \mathbf{G} \mid \text{Ad}(g)(H) = H, \forall H \in \mathfrak{z}_0\}. \end{aligned}$$

Proof. While proving this Proposition we can assume that $\mathfrak{h}$ is maximally non-compact among the Cartan subalgebras of $\mathfrak{g}$ that are contained in $\mathfrak{g}_+$. Indeed, all admissible Cartan subalgebras of $(\mathfrak{g}, \mathfrak{q})$ are conjugate to the direct sum of $\mathfrak{z}_0$ and a Cartan subalgebra $\mathfrak{h}_0$ of $\mathfrak{s}_0$. Thus our assumption means that we have chosen a maximally noncompact Cartan subalgebra $\mathfrak{h}_0$ of $\mathfrak{s}_0$.

We first prove that

$$(7.23) \quad \mathbf{G}_0 = \{g \in \mathbf{G}_+ \mid \text{Ad}(g)(\mathfrak{g}_0) = \mathfrak{g}_0\}.$$

If $g \in \mathbf{G}_+$, then $\text{Ad}(g)(\mathfrak{g}_0)$ is a maximal reductive subalgebra of $\mathfrak{g}_+$. By [OV, Ch.I, §6.5] there is an element $n \in \mathbf{N}$ such that $\mathfrak{g}_0 = \text{Ad}(g n^{-1})(\mathfrak{g}_0)$. This yields a decomposition $g = g_0 n$, with $g_0 \in \mathbf{N}_{\mathbf{G}_+}(\mathfrak{g}_0)$ and $n \in \mathbf{N}$. Since $\mathfrak{z}_0 \subset \mathfrak{g}_0$, and $[\mathfrak{z}_0, \mathfrak{n}] = \mathfrak{n}$, we have $\mathbf{N} \cap \mathbf{N}_{\mathbf{G}_+}(\mathfrak{g}_0) = \{1\}$. This yields the decomposition $\mathbf{G}_+ = \mathbf{N}_{\mathbf{G}_+}(\mathfrak{g}_0) \rtimes \mathbf{N}$. Since we also have (7.21) and $\mathfrak{g}_0$ is the Lie algebra of both $\mathbf{G}_0$ and $\mathbf{N}_{\mathbf{G}_+}(\mathfrak{g}_0)$, we obtain that $\mathbf{G}_0 = \mathbf{N}_{\mathbf{G}_+}(\mathfrak{g}_0)$.

Next we show that $\mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) = \mathbf{N}_{\mathbf{G}}(\mathfrak{g}_0)$. If $g_0 \in \mathbf{N}_{\mathbf{G}}(\mathfrak{g}_0)$, then $\text{Ad}(g_0)(\mathfrak{z}_0) = \mathfrak{z}_0$. Moreover, $\text{Ad}(g_0)(\mathfrak{h}_0)$ is another maximally noncompact Cartan subalgebra of $\mathfrak{s}_0$. Therefore there is an inner automorphism of $\mathfrak{s}_0$, and thus also an element $g_1 \in \mathbf{S}_0$ such that $g_1 \circ g_0(\mathfrak{h}) = \mathfrak{h}$ (see e.g. [S1], [S2], [OV, Ch.4, §4.7]). Let $g = g_1 \circ g_0$. Since $\mathbf{S}_0 \subset \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$, it suffices to show that $g \in \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$. We have $\text{Ad}(g)(\mathfrak{z}_0) = \mathfrak{z}_0$ and $\text{Ad}(g)(\mathfrak{h}_0) = \mathfrak{h}_0$. In particular, $g \in \mathbf{N}_{\mathbf{G}}(\mathfrak{h})$. Since $\text{Ad}(g)$ commutes with the conjugation σ in $\hat{\mathfrak{g}}$, and $\text{Ad}(g)(\mathfrak{q}^n) = \mathfrak{q}^n$, $\text{Ad}(g)(\mathfrak{q}^r) = \mathfrak{q}^r$, we also have:

$$(7.24) \quad \text{Ad}(g)(\mathfrak{q}^r \cap \bar{\mathfrak{q}}^r) = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r \quad \text{and} \quad \text{Ad}(g)(\mathfrak{q}^n + [\bar{\mathfrak{q}} \cap \mathfrak{q}^r]) = \mathfrak{q}^n + [\bar{\mathfrak{q}} \cap \mathfrak{q}^r].$$

The analytic Weyl group $\mathbf{W}(\mathfrak{g}, \mathfrak{h}) = \mathbf{N}_{\mathbf{G}}(\mathfrak{h})/\mathbf{H}$ can be identified to a subgroup of the Weyl group $\mathbf{W}(\mathcal{R}) = \mathbf{N}_{\hat{\mathbf{G}}}(\hat{\mathfrak{h}})/\hat{\mathbf{H}}$. The element $s_g \in \mathbf{W}(\mathcal{R})$ defined by g satisfies: $\text{Ad}(g)(\hat{\mathfrak{g}}^\alpha) \subset \hat{\mathfrak{g}}^{s_g(\alpha)}$ for all $\alpha \in \mathcal{R}$. Because of (7.24), g normalizes the complex parabolic subalgebra $\mathfrak{v} = \mathfrak{q}^n + (\mathfrak{q}^r \cap \bar{\mathfrak{q}})$, with $\mathfrak{v}^r = \mathfrak{q}^r \cap \bar{\mathfrak{q}}^r$. Hence s_g is the composition of symmetries s_α with $\alpha \in \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r$. If $\alpha \in \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r$ and $H \in \mathfrak{z}_0$ we obtain for $X_\beta \in \hat{\mathfrak{g}}^\beta$:

$$\begin{aligned} [\text{Ad}(g)(H), \text{Ad}(g)(X_\beta)] &= \text{Ad}(g)([H, X_\beta]) \\ &= \beta(H) \text{Ad}(g)(X_\beta) = s_\alpha(\beta)(H) \text{Ad}(g)(X_\beta) \end{aligned}$$

because $s_\alpha(\beta)(H) = (\beta - (\beta|\alpha^\vee)\alpha)(H) = \beta(H)$, since $\alpha(H) = 0$ by (7.19). This shows that $[\text{Ad}(g)(H), X_\beta] = \beta(H)X_\beta$ for all $\beta \in \mathcal{R}$ and hence that $\text{Ad}(g)(H) = H$. Moreover $\mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) \subset \mathbf{N}_{\mathbf{G}}(\mathfrak{g}_0)$ because $\mathfrak{g}_0$ is the centralizer of $\mathfrak{z}_0$ in $\mathfrak{g}$, and hence the equality follows.

Finally we observe that $\mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) \subset \mathbf{Q} = \mathbf{N}_{\hat{\mathbf{G}}}(\mathfrak{q})$. Indeed, if $g_0 \in \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$, we can find $g_1 \in \mathbf{S}_0$ such that $g = g_1 g_0$ normalizes our Cartan subalgebra $\mathfrak{h}$, still centralizing $\mathfrak{z}_0$. Being $\mathcal{Q} = \{\alpha \in \mathcal{R} \mid \alpha(A) \geq 0\}$ for an element $A \in \mathfrak{h}_{\mathbb{R}} \cap \hat{\mathfrak{z}}_0$, clearly $\text{Ad}(g)(\mathfrak{q}) = \mathfrak{q}$. Hence also $\text{Ad}(g_0)(\mathfrak{q}) = \mathfrak{q}$. This shows that $\mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) \subset \mathbf{G}_+$, completing the proof of the Theorem. $\square$

Next we prove that our choice of $\mathbf{G} = \mathbf{G}_{\Pi(\mathcal{R})}$ brings that also $\mathbf{S}_0$ is the semisimple real group associated to the full weights lattice of $\hat{\mathfrak{s}}_0$. Indeed we have:

Theorem 7.5. *The complexification $\hat{\mathbf{S}}_0$ of the linear Lie group $\mathbf{S}_0$ is simply connected.*

First we prove the following :

Lemma 7.6. *Let E be a linear hyperplane of $\mathfrak{h}_{\mathbb{R}}^*$ and let $\mathcal{R}' = E \cap \mathcal{R}$. Then every basis of simple roots $\{\alpha_1, \dots, \alpha_m\}$ of $\mathcal{R}'$ is a subset of a basis of simple roots $\{\alpha_1, \dots, \alpha_m, \alpha_{m+1}, \dots, \alpha_\ell\}$ of $\mathcal{R}$.*

Proof. Let $A \in \mathfrak{h}_{\mathbb{R}}$ be such that $E = \{\eta \in \mathfrak{h}_{\mathbb{R}}^* \mid \eta(A) = 0\}$. Next we take a regular element $B \in \mathfrak{h}_{\mathbb{R}}$ such that $\{\alpha_1, \dots, \alpha_m\}$ is the system of simple roots in $\mathcal{R}'^+ = \{\alpha \in \mathcal{R}' \mid \alpha(B) > 0\}$. For small $\varepsilon > 0$ the element $A + \varepsilon B$ is regular and $\alpha_1, \dots, \alpha_m$ are simple in $\mathcal{R}^+ = \{\alpha \in \mathcal{R} \mid \alpha(A + \varepsilon B) > 0\}$. Indeed, we take $\varepsilon > 0$ such that $\varepsilon |\alpha(B)| < |\alpha(A)|$ when $\alpha(A) \neq 0$. Assume by contradiction that, for some $1 \leq i \leq m$, the root α_i is not simple in $\mathcal{R}^+$, i.e. that we have $\alpha_i = \beta + \gamma$, with $\beta(A + \varepsilon B) > 0$, $\gamma(A + \varepsilon B) > 0$, $\beta(B) \geq \gamma(B)$. Since α_i is simple in $\mathcal{R}'^+$, we have $\gamma(B) < 0 < \alpha_i(B) < \beta(B)$. Hence $\gamma(A) > 0$. Moreover $\beta(A) \geq 0$, because otherwise $\beta(A + \varepsilon B) < 0$ by our choice of ε . Thus we obtain :

$$\alpha_i(A) = \beta(A) + \gamma(A) \geq \gamma(A) > 0,$$

contradicting $\alpha_i \in E$. $\square$

Proof of Theorem 7.5. We keep the notation introduced in the previous discussion. While applying Lemma 7.6 to our situation, we observe that, if our parabolic $\mathcal{Q}$ is $\{\alpha \in \mathcal{R} \mid \alpha(A) \geq 0\}$ for some $A \in \mathfrak{h}_{\mathbb{R}}$, then the set of roots α with $\hat{\mathfrak{g}}^\alpha \subset \hat{\mathfrak{s}}_0$ is $\mathcal{R}' = \mathcal{Q}^r \cap \bar{\mathcal{Q}}^r = \{\alpha \in \mathcal{R} \mid \alpha(A + \varepsilon \bar{A}) = 0\}$, where ε is any positive real number with $\varepsilon |\bar{\alpha}(A)| < \alpha(A)$ for $\alpha \in \mathcal{Q}^n$. This set $\mathcal{R}'$ is naturally isomorphic to the root system $\mathcal{R}(\hat{\mathfrak{s}}_0, \hat{\mathfrak{h}}_0)$ of the semisimple complex Lie algebra $\hat{\mathfrak{s}}_0$ with respect to its Cartan subalgebra $\hat{\mathfrak{h}}_0$. We identify $\Pi(\mathcal{R}')$ to the set of elements ω in $\langle \mathcal{R}' \rangle_{\mathbb{R}} \subset \mathfrak{h}_{\mathbb{R}}^*$ for which $(\omega | \alpha^\vee) \in \mathbb{Z}$ for all $\alpha \in \mathcal{R}'$. We have a natural projection $\varpi : \Pi(\mathcal{R}) \rightarrow \Pi(\mathcal{R}')$, that is defined by $(\varpi(\eta) | \alpha^\vee) = (\eta | \alpha^\vee)$ for all $\alpha \in \mathcal{R}'$, and coincides with the orthogonal projection onto $\langle \mathcal{R}' \rangle_{\mathbb{R}}$.

The lattice $\mathcal{L} = \varpi(\Pi(\mathcal{R}))$ satisfies $\Lambda(\mathcal{R}') \subset \mathcal{L} \subset \Pi(\mathcal{R}')$ and is the set of weights of the finite dimensional linear representations of $\mathbf{S}_0$.

According to Lemma 7.6 we can fix $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, with $\mathcal{B}(C) = \{\alpha_1, \dots, \alpha_\ell\}$, in such a way that $\mathcal{B}' = \{\alpha_1, \dots, \alpha_m\}$, with $m \leq \ell$, is a basis of $\mathcal{R}'$. Let $\mathcal{B}^*(C) = \{\mu_1, \dots, \mu_\ell\} \subset \Pi(\mathcal{R})$ and $\mathcal{B}'^* = \{\nu_1, \dots, \nu_m\} \subset \Pi(\mathcal{R}') \subset \langle \mathcal{R}' \rangle_{\mathbb{R}}$ be the corresponding adjoint basis. Then $\varpi(\mu_i) = \nu_i$ for $i = 1, \dots, m$. This shows that actually $\mathcal{L} = \Pi(\mathcal{R}')$, proving our statement. $\square$

Now we give a more accurate description of the group $\pi_0(\mathbf{G}_+)$ of the connected components of $\mathbf{G}_+$.

Theorem 7.7. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra. Keep the notation of (7.17), (7.18), (7.20) and (7.21). Then $\mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}_0)$ is a closed normal Lie subgroup of $\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$. The natural inclusion $\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) \hookrightarrow \mathbf{G}_+$ passes to the quotient to define a one-to-one correspondence of connected components :*

$$(7.25) \quad \pi_0(\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0) / \mathbf{N}_{\mathbf{S}_0}(\mathfrak{h})) \longleftrightarrow \pi_0(\mathbf{G}_+).$$

Proof. The inclusion $\mathbf{G}_0 \hookrightarrow \mathbf{G}_+$ defines an isomorphism $\pi_0(\mathbf{G}_0) \xrightarrow{\sim} \pi_0(\mathbf{G}_+)$.

We showed in the proof of Proposition 7.4 that every connected component of $\mathbf{G}_0$ contains an element of $\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{G}_0$ and that $\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{G}_0 = \mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$.

Since $\mathfrak{s}_0$ is an ideal of $\mathfrak{g}_0 = \mathfrak{s}_0 \oplus \mathfrak{z}_0$, the analytic subgroup $\mathbf{S}_0$ is normal in $\mathbf{G}_0$ and we already noticed that is contained in $\mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$. Moreover, $\mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}) = \mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{S}_0$.

Let $\mathbf{T}_0$ be the analytic Lie subgroup of $\mathbf{G}_0$ with Lie algebra $\mathfrak{z}_0$. Then $\mathbf{S}_0 \bowtie \mathbf{T}_0$ is the connected component of the identity in $\mathbf{G}_0$. We have:

$$\mathbf{T}_0 \subset \mathbf{H} \subset \mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0).$$

Hence we obtain isomorphisms:

$$(7.26) \quad \frac{\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)}{\mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}) \bowtie \mathbf{T}_0} \xrightarrow{\cong} \frac{\mathbf{G}_0}{\mathbf{S}_0 \bowtie \mathbf{T}_0} \xrightarrow{\cong} \pi_0(\mathbf{G}_+),$$

yielding the isomorphism in (7.25). $\square$

We denote by $\mathbf{H}_0$ the Cartan subalgebra of $\mathbf{S}_0$ corresponding to $\mathfrak{h}_0$. Since $\mathbf{S}_0 \subset \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$, we have:

$$(7.27) \quad \mathbf{H}_0 = \{g \in \mathbf{S}_0 \mid \text{Ad}(g)(H) = H, \forall H \in \mathfrak{h}_0\} = \mathbf{H} \cap \mathbf{S}_0.$$

By (7.27), we obtain for the analytic Weyl group $\mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0) = \mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}_0)/\mathbf{H}_0$:

$$(7.28) \quad \mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0) \cong \mathbf{N}_{\mathbf{S}_0}(\mathfrak{h})/\mathbf{Z}_{\mathbf{S}_0}(\mathfrak{h}) = \mathbf{N}_{\mathbf{S}_0}(\mathfrak{h})/(\mathbf{H} \cap \mathbf{S}_0).$$

Thus we can identify $\mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)$ with a subgroup of the centralizer $\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)$ of $\mathfrak{z}_0$ in $\mathbf{W}(\mathfrak{g}, \mathfrak{h})$. Since $\mathbf{S}_0$ is normal in $\mathbf{G}_0$, it turns out that $\mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)$ is normal in $\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)$. We have:

Theorem 7.8. *Keep the notation and assumptions of Theorem 7.7. We have a short exact sequence of groups and homomorphisms:*

$$(7.29) \quad 1 \rightarrow \frac{\mathbf{H}}{\mathbf{H}_0} \rightarrow \frac{\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)}{\mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}_0)} \rightarrow \frac{\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)}{\mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)} \rightarrow 1.$$

We obtain an exact sequence of finite groups:

$$(7.30) \quad 1 \rightarrow \pi_0 \left(\frac{\mathbf{H}}{\mathbf{H}_0} \right) \rightarrow \pi_0(\mathbf{G}_+) \rightarrow \frac{\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)}{\mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)} \rightarrow 1.$$

When $\mathfrak{h}$ is maximally compact in $\mathfrak{g}_+$, then $\mathbf{H}_0$ is connected and (7.30) yields the exact sequence:

$$(7.31) \quad 1 \rightarrow \pi_0(\mathbf{H}) \rightarrow \pi_0(\mathbf{G}_+) \rightarrow \frac{\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)}{\mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)} \rightarrow 1.$$

When $\mathfrak{h}$ is maximally noncompact in $\mathfrak{g}_+$, then $\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0) \cong \mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)$ and (7.30) yields the isomorphism:

$$(7.32) \quad \pi_0 \left(\frac{\mathbf{H}}{\mathbf{H}_0} \right) \cong \pi_0(\mathbf{G}_+).$$

Proof. Since $\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)$ is a finite group, (7.30) is a consequence of the exact homotopy sequence of a fiber bundle, of (7.29) and of Theorem 7.7. Thus a proof of (7.29) also provides a proof of (7.30).

We observe that $\mathbf{H}$ is a normal subgroup of $\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$ and that the quotient $(\mathbf{N}_{\mathbf{G}}(\mathfrak{h}) \cap \mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)) / \mathbf{H}$ is naturally isomorphic to the centralizer $\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0)$ of $\mathfrak{z}_0$ in the analytic Weyl group $\mathbf{W}(\mathfrak{g}, \mathfrak{h})$. Moreover, $\mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}_0)$ is contained in $\mathbf{Z}_{\mathbf{G}}(\mathfrak{z}_0)$ and $\mathbf{N}_{\mathbf{S}_0}(\mathfrak{h}_0) \cap \mathbf{H} = \mathbf{H}_0$. Thus (7.29) follows from the general homomorphism theorems of groups.

If $\mathfrak{h}$ is maximally compact, then $\mathfrak{h}_0$ is a maximally compact Lie subalgebra of $\mathfrak{s}_0$. Then (see e.g. [Kn, Proposition 7.90, p.488]) the Cartan subalgebra $\mathbf{H}_0$ is connected and (7.31) follows from (7.30).

If $\mathfrak{h}$ is maximally noncompact, then, by [Kn, Proposition 7.90, p.488] the Cartan subalgebra $\mathbf{H}$ intersects every connected component of $\mathbf{G}_+$. This implies that the map $\pi_0(\mathbf{H}/\mathbf{H}_0) \rightarrow \pi_0(\mathbf{G}_+)$ in (7.30) is onto and hence (7.32) holds true and $\mathbf{Z}_{\mathbf{W}(\mathfrak{g}, \mathfrak{h})}(\mathfrak{z}_0) \cong \mathbf{W}(\mathfrak{s}_0, \mathfrak{h}_0)$. $\square$

We describe now $\pi_0(\mathbf{G}_+)$ in terms of characters :

Proposition 7.9. *Let $\mathfrak{h}$ be a maximally noncompact Cartan subalgebra of $\mathfrak{g}_+$. Fix a Weyl chamber $C \in \mathfrak{C}(\mathcal{R})$ such that $\mathcal{B}(C) = \{\alpha_1, \dots, \alpha_\ell\}$ contains a basis $\mathcal{B}_0(C) = \{\alpha_1, \dots, \alpha_m\}$ of simple roots of $\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r$. Let $E = \langle \alpha_{m+1}, \dots, \alpha_\ell \rangle^\perp$. Then :*

$$(7.33) \quad \pi_0(\mathbf{G}_+) \simeq \pi_0(\{\chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*) \mid \chi(\mu) = 1, \forall \mu \in \Pi(\mathcal{R}) \cap E\}).$$

Proof. We denote by ϖ_0 the orthogonal projection onto $\langle \alpha_{m+1}, \dots, \alpha_\ell \rangle^\perp$. Let $\mathcal{B}^*(C) = \{\omega_1, \dots, \omega_\ell\}$ be the adjoint basis of $\mathcal{B}(C)$. Then we have $\varpi_0(\eta) = \sum_{i=1}^m (\eta | \alpha_i^\vee) \omega_i$. We can identify $\mathcal{R}' = \mathcal{R}(\hat{\mathfrak{s}}_0, \hat{\mathfrak{h}}_0)$ with $\varpi_0(\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r)$. The advantage of this identification is that the corresponding weight space $\Pi(\mathcal{R}')$ becomes a subspace of $\Pi(\mathcal{R})$, and moreover :

$$(7.34) \quad \Pi(\mathcal{R}') = \varpi_0(\Pi(\mathcal{R})) = \Pi(\mathcal{R}) \cap \langle \alpha_{m+1}, \dots, \alpha_\ell \rangle^\perp.$$

In this way, for each character $\chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*)$, the composition $\chi \circ \varpi_0$ is still a character of $\text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*)$, whose restriction to $\Pi(\mathcal{R}')$ is a character in $\text{Hom}^\sigma(\Pi(\mathcal{R}'), \mathbb{C}^*)$, and we have :

$$(7.35) \quad \mathbf{H}_0 = \{h_\chi \mid \chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}'), \mathbb{C}^*)\} = \{h_{\chi \circ \varpi} \mid \chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*)\}.$$

Thus, setting :

$$(7.36) \quad \mathbf{H}_1 = \{h_\chi \mid \chi \in \text{Hom}^\sigma(\Pi(\mathcal{R}), \mathbb{C}^*), \chi(\eta) = 1 \forall \eta \in \Pi(\mathcal{R}')\},$$

the map :

$$(7.37) \quad \mathbf{H} \ni h_\chi \rightarrow h_{\chi \circ \varpi}^{-1} \circ h_\chi \in \mathbf{H}_1$$

yields, by passing to the quotient, the isomorphism $\mathbf{H}/\mathbf{H}_0 \xrightarrow{\sim} \mathbf{H}_1$, which implies (7.33). $\square$

Corollary 7.10. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic totally real CR algebra. Let $\mathfrak{h}$ be a maximally noncompact Cartan subalgebra of $\mathfrak{g}$ contained in $\mathfrak{g}_+$ and let $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ be S-fit for $(\mathfrak{g}, \mathfrak{q})$ and hence S-adapted to the conjugation σ . Then $\pi_0(\mathbf{G}_+)$ is isomorphic to $\mathbb{Z}_2^e$, where e is the number of roots in $\Phi_C \cap \mathcal{R}_{\text{re}}$.*

Proof. By inspecting the Satake diagrams of the simple real Lie algebras, we obtain that $\chi(\omega_i)$ is a positive real number when ω_i is a real weight corresponding to a complex root $\alpha_i \in \mathcal{B}(C)$. Then the statement follows from Proposition 7.9.

Another independent proof can be given, using [DKV] and [Wi]. Because $\mathbf{G}$ is connected, we have an exact sequence :

$$(7.38) \quad \cdots \longrightarrow \pi_1(\mathbf{G}/\mathbf{G}_+) \longrightarrow \pi_0(\mathbf{G}_+) \longrightarrow \mathbf{1}.$$

Take $\Phi'_C = \Phi_C \setminus \mathcal{R}_{\text{re}}$. Then also $\mathfrak{q}' = \mathfrak{q}_{\Phi'_C}$ is the complexification of a real parabolic subalgebra $\mathfrak{g}'_+$ of $\mathfrak{g}$. By [Wi] and (7.38), the stabilizer $\mathbf{G}'_+$ of $\mathfrak{q}'$ in $\mathbf{G}$ is connected. By Proposition 7.9, this implies that $\chi(\omega_i) > 0$ on the real $\omega_i \in \mathcal{B}^*(C)$ corresponding to a complex root α_i in $\mathcal{B}(C)$. Again, the statement follows from Proposition 7.9. $\square$

For each $\omega \in \Pi(\mathcal{R})$ there is, modulo isomorphisms, a unique irreducible finite dimensional complex linear representation $\rho_\omega : \hat{\mathfrak{g}} \rightarrow \mathfrak{gl}_{\mathbb{C}}(V_\omega)$, for which ω is an extremal weight. For each weight η we denote by :

$$(7.39) \quad V_\omega^\eta = \{v \in V_\omega \mid \rho_\omega(H)(v) = \eta(H)v, \forall H \in \hat{\mathfrak{h}}\}$$

the eigenspace of V_ω corresponding to the weight η . Let :

$$(7.40) \quad \hat{\mathfrak{g}}_\omega = \{Z \in \hat{\mathfrak{g}} \mid \rho_\omega(V_\omega^\omega) \subset V_\omega^\omega\}.$$

Since the eigenspace V_ω^ω of ω is one-dimensional, for each $Z \in \hat{\mathfrak{g}}_\omega$ there is a unique complex number, that we shall denote by $\omega(Z)$, such that :

$$(7.41) \quad \rho_\omega(Z)(v) = \omega(Z)v, \quad \forall Z \in \hat{\mathfrak{g}}_\omega, \forall v \in V_\omega^\omega.$$

This agrees with the natural definition of $\omega(Z)$ by the duality pairing when $Z \in \hat{\mathfrak{h}}$.

Proposition 7.11. *Let $\omega_1, \omega_2 \in \Pi(\mathcal{R})$ and $\omega = \omega_1 + \omega_2$. Then :*

$$(7.42) \quad \hat{\mathfrak{g}}_{\omega_1} \cap \hat{\mathfrak{g}}_{\omega_2} \subset \hat{\mathfrak{g}}_\omega$$

and :

$$(7.43) \quad \omega(Z) = \omega_1(Z) + \omega_2(Z), \quad \forall Z \in \hat{\mathfrak{g}}_{\omega_1} \cap \hat{\mathfrak{g}}_{\omega_2}.$$

Proof. There is an injective homomorphism of $\hat{\mathfrak{g}}$ -modules $V_\omega \rightarrow V_{\omega_1} \otimes V_{\omega_2}$ that maps V_ω^ω into $V_{\omega_1}^{\omega_1} \otimes V_{\omega_2}^{\omega_2}$. For $Z \in \hat{\mathfrak{g}}$, we have $(\rho_{\omega_1} \otimes \rho_{\omega_2})(v_1 \otimes v_2) = \rho_{\omega_1}(Z)(v_1) \otimes v_2 + v_1 \otimes \rho_{\omega_2}(Z)(v_2)$ for all $v_1 \in V_{\omega_1}, v_2 \in V_{\omega_2}$. This implies (7.42) and (7.43). $\square$

We also have :

Lemma 7.12. *Let $\omega \in \Pi(\mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}}))$ and $\hat{\mathfrak{h}}'$ another Cartan subalgebra of $\hat{\mathfrak{g}}$ contained in $\hat{\mathfrak{g}}_\omega$. Let V_ω be an irreducible $\hat{\mathfrak{g}}$ -module with extremal weight ω . Then :*

- (i) *There is a weight $\omega' \in \Pi(\mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}}'))$ that is an extremal weight for V_ω ;*
- (ii) *we can choose $\omega' \in \Pi(\mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}}'))$ in such a way that $V_\omega^\omega = V_\omega^{\omega'}$;*
- (iii) *$\hat{\mathfrak{g}}_\omega = \hat{\mathfrak{g}}_{\omega'}$ and $\omega(Z) = \omega'(Z) \quad \forall Z \in \hat{\mathfrak{g}}_\omega$. $\square$*

Since we took a simply connected $\hat{\mathbf{G}}$, for each $\omega \in \Pi(\mathcal{R})$, the representation $\rho_\omega : \hat{\mathfrak{g}} \rightarrow \mathfrak{gl}_\mathbb{C}(V_\omega)$ lifts to a representation $\rho_\omega : \hat{\mathbf{G}} \rightarrow \mathbf{GL}_\mathbb{C}(V_\omega)$. The stabilizer of the line V_ω^ω in $\hat{\mathbf{G}}$ contains the Cartan subgroup $\hat{\mathbf{H}}$ of $\hat{\mathbf{G}}$ and therefore is the closed connected subgroup $\hat{\mathbf{G}}_\omega$ of $\hat{\mathbf{G}}$ with Lie algebra $\hat{\mathfrak{g}}_\omega$. The map $\omega : \hat{\mathfrak{g}}_\omega \rightarrow \mathbb{C}$ also lifts to a character $\varphi_\omega : \hat{\mathbf{G}}_\omega \rightarrow \mathbb{C}^*$, with $\varphi_\omega(\exp(Z)) = \exp(\omega(Z))$ for all $Z \in \hat{\mathfrak{g}}_\omega$.

We observe that every parabolic $\mathfrak{q}$ containing $\hat{\mathfrak{h}}$ is equal to some $\hat{\mathfrak{g}}_\omega$, for a suitable choice of $\omega \in \Pi(\mathcal{R})$. This choice is not unique. It can be done in the following way. First we choose a Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$. Let $\mathfrak{q} = \mathfrak{q}_{\Phi_C}$ for $\Phi_C \subset \mathcal{B}(C)$. Let $\mathcal{B}^*(C)$ be the adjoint basis in $\Pi(\mathcal{R})$ and $\Phi^*(C) = \{\omega_1, \dots, \omega_k\}$ the set of elements of $\mathcal{B}^*(C)$ that vanish on $\mathcal{B}(C) \setminus \Phi_C$. Then for any k -tuple of strictly positive integers $(t_1, \dots, t_k)$ we have :

$$(7.44) \quad \mathfrak{q} = \hat{\mathfrak{g}}_{t_1\omega_1 + \dots + t_k\omega_k} = \bigcap_{1 \leq i \leq k} \hat{\mathfrak{g}}_{\omega_i}.$$

Proposition 7.13. *Let $\mathbf{Q}$ be the parabolic subgroup of $\hat{\mathbf{G}}$ of the parabolic complex Lie subalgebra $\mathfrak{q}$. Then, with the notation above :*

$$(7.45) \quad \mathbf{Q} = \bigcap_{1 \leq i \leq k} \hat{\mathbf{G}}_{\omega_i} \quad \text{and} \quad \text{Hom}_\mathbb{C}(\mathbf{Q}, \mathbb{C}^*) = \{\varphi_\omega \mid \omega \in \Pi(\mathcal{R}) \cap \langle \mathcal{Q}^r \rangle_\mathbb{R}^\perp\}$$

is the free Abelian group generated by $\varphi_{\omega_1}, \dots, \varphi_{\omega_k}$.

Proof. The inclusion $\{\varphi_\omega \mid \omega \in \Pi(\mathcal{R}) \cap \langle \mathcal{Q}^r \rangle_\mathbb{R}^\perp\} \subset \text{Hom}_\mathbb{C}(\mathbf{Q}, \mathbb{C}^*)$ is a consequence of the previous discussion.

To prove the opposite inclusion, we first observe that a character in $\text{Hom}_\mathbb{C}(\mathbf{Q}, \mathbb{C}^*)$ restricts to a character in $\text{Hom}_\mathbb{C}(\hat{\mathbf{H}}, \mathbb{C}^*)$, and hence is of the form φ_ω for some $\omega \in \Pi(\mathcal{R})$. Moreover, being connected, $\mathbf{Q}$ admits a Levi decomposition $\mathbf{Q} = \mathbf{Q}_{\text{ss}} \mathbf{Q}_{\text{rad}}$ and $\varphi_\omega(g) = 1$ for $g \in \mathbf{Q}_{\text{ss}}$ (= the Levi subgroup of $\mathbf{Q}$). This yields $(\omega|\alpha^\vee) = 0$ for $\alpha \in \mathcal{Q}^r$. $\square$

Lemma 7.14. *If $\omega \in \Pi(\mathcal{R})$ is a real weight, then φ_ω is real valued on $\hat{\mathbf{G}}_\omega \cap \mathbf{G}$.*

Proof. From $\varphi_\omega(\exp(Z)) = \exp(\omega(Z))$ for $Z \in \hat{\mathfrak{g}}_\omega$ we obtain that $\varphi_\omega(g)$ is real when g belongs to a neighborhood of the identity in $\hat{\mathbf{G}}_\omega \cap \mathbf{G}$. Since $\varphi_\omega(g)$ is an algebraic function of g , the statement follows because any neighborhood of the identity is Zariski dense in $\hat{\mathbf{G}}_\omega \cap \mathbf{G}$. $\square$

Let $(\vartheta, \mathfrak{h})$ be a Cartan pair adapted to the effective parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$, choose $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ and set :

$$\Phi_C^* = \{\omega_1, \dots, \omega_k\} = \mathcal{B}^*(C) \cap (\mathcal{Q}^r \cap \bar{\mathcal{Q}}^r)^\perp.$$

Then $\mathbf{G}_+ \subset \bigcap_{1 \leq i \leq k} \hat{\mathbf{G}}_{\omega_i}$, and we can define the map :

$$(7.46) \quad \varphi : \mathbf{G}_+ \ni g \rightarrow (\varphi_{\omega_1}(g), \dots, \varphi_{\omega_k}(g)) \in (\mathbb{C}^*)^k.$$

We have :

Theorem 7.15. *The map (7.46) yields, by passing to the quotients, a group isomorphism: $\pi_0(\mathbf{G}_+) \xrightarrow{\varphi_*} \pi_0(\varphi(\mathbf{G}_+))$.*

Proof. By using Lemma 7.12, we can restrain to proving the Theorem in the case $\mathfrak{h}$ is maximally noncompact among the admissible Cartan subalgebras of $\mathfrak{g}$ contained in $\mathfrak{g}_+$. Moreover, by passing to the weakest CR model, we can as well assume that $\mathcal{Q}^r = \bar{\mathcal{Q}}^r$. By Proposition 7.9, we obtain a commutative diagram:

$$(7.47) \quad \begin{array}{ccc} \pi_0(\mathbf{H}/\mathbf{H}_0) & \xrightarrow{\sim} & \pi_0(\mathbf{G}_+) \\ \downarrow \wr & & \downarrow \varphi_* \\ \pi_0(\varphi(\mathbf{H})) & \xrightarrow{\sim} & \pi_0(\varphi(\mathbf{G}_+)) \end{array}$$

showing that also φ_* is an isomorphism. $\square$

In particular, when $(\mathfrak{g}, \mathfrak{q})$ is totally real, i.e. $\mathfrak{q} = \bar{\mathfrak{q}}$, we obtain, by using Corollary 7.10:

Theorem 7.16. *Let $(\mathfrak{g}, \mathfrak{q})$ be a totally real parabolic CR algebra, $\mathfrak{h}$ a maximally noncompact Cartan subalgebra of $\mathfrak{g}$ contained in $\mathfrak{g}_+$, $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ an S-fit (and S-adapted) Weyl chamber. Set:*

$$\mathcal{B}(C) = \{\alpha_1, \dots, \alpha_\ell\}, \quad \{\alpha_1, \dots, \alpha_\mu\} = \Phi_C \cap \mathcal{R}_{\text{re}}, \quad \mathcal{B}^*(C) = \{\omega_1, \dots, \omega_\ell\}.$$

Then $\varphi_{\omega_i}(g) \in \mathbb{R}^*$ for $1 \leq i \leq \mu$. The map:

$$(7.48) \quad \varphi^b: \mathbf{G}_+ \ni g \rightarrow \left(\frac{\varphi_{\omega_1}(g)}{|\varphi_{\omega_1}(g)|}, \dots, \frac{\varphi_{\omega_\mu}(g)}{|\varphi_{\omega_\mu}(g)|} \right) \in \{-1, 1\}^\mu = \mathbb{Z}_2^\mu$$

defines, by passing to the quotient, an isomorphism:

$$(7.49) \quad \varphi_*^b: \pi_0(\mathbf{G}_+) \ni [g] \rightarrow \varphi^b(g) \in \mathbb{Z}_2^\mu. \quad \square$$

Theorem 7.17. *We keep the notation and assumptions of Theorem 7.16. For each $\alpha \in \Phi_C \cap \mathcal{R}_{\text{re}}$, let $\mathbf{S}_\alpha$ be the simple analytic real subgroup of $\mathbf{G}$, with Lie algebra $\mathfrak{s}_\alpha = \mathfrak{g} \cap (\hat{\mathfrak{g}}^\alpha \oplus \hat{\mathfrak{g}}^{-\alpha} \oplus [\hat{\mathfrak{g}}^\alpha, \hat{\mathfrak{g}}^{-\alpha}])$. Let t_α^* be a generator of $\pi_1(\mathbf{S}_\alpha) \simeq \mathbb{Z}$, and let t_α be its image in $\pi_1(\mathbf{G}/\mathbf{G}_+)$. Then $\{t_\alpha \mid \alpha \in \Phi_C \cap \mathcal{R}_{\text{re}}\}$ is a set of generators of $\pi_1(\mathbf{G}/\mathbf{G}_+)$, and the homomorphism $\delta: \pi_1(\mathbf{G}/\mathbf{G}_+) \rightarrow \pi_0(\mathbf{G}_+)$ of the exact homotopy sequence (7.38) of the principal bundle $\mathbf{G} \rightarrow \mathbf{G}/\mathbf{G}_+$ is given by:*

$$(7.50) \quad \delta: \pi_1(\mathbf{G}/\mathbf{G}_+) \ni t_{\alpha_i} \rightarrow (1, \dots, \underbrace{-1}_i, \dots, 1) \in \mathbb{Z}_2^\mu \simeq \pi_0(\mathbf{G}_+).$$

Proof. By using results from [DKV], in [Wi, §2] it is shown that the fundamental group of $\mathbf{G}/\mathbf{G}_+$ is generated by the images of the generators of the $\pi_1(\mathbf{S}_\alpha)$'s, for simple roots $\alpha \in \Phi_C \cap \mathcal{R}_{\text{re}}$. In fact the generalization of the Bruhat decomposition in [DKV] yields a cell decomposition of $\mathbf{G}/\mathbf{G}_+$ where the open 1-cells correspond to roots $\alpha \in \Phi_C \cap \mathcal{R}_{\text{re}}$. Let $\rho_\alpha: \mathfrak{sl}(2, \mathbb{R}) \rightarrow \mathfrak{s}_\alpha$ be the representation with:

$$\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \xrightarrow{\rho_\alpha} X_{-\alpha}, \quad \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} \xrightarrow{\rho_\alpha} X_\alpha, \quad \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \xrightarrow{\rho_\alpha} H_\alpha.$$

We still denote by ρ_α the corresponding group representation $\mathbf{SL}(2, \mathbb{R}) \rightarrow \mathbf{S}_\alpha$. The open 1-cell corresponding to α can be parametrized by :

$$\left\{ g(t) = \rho_\alpha \left[\begin{pmatrix} \tan t & 1 \\ -1 & 0 \end{pmatrix} \right] \mid -\pi/2 < t < \pi/2 \right\}.$$

Since $g_+(t) = \rho_\alpha \left[\begin{pmatrix} \cos t & -\sin t \\ 0 & 1/\cos t \end{pmatrix} \right] \in \mathbf{G}_+$ for all $|t| < \pi/2$, the closure in $\mathbf{G}/\mathbf{G}_+$ of this 1-cell is the loop that is the image in $\mathbf{G}/\mathbf{G}_+$ of:

$$[-\pi/2, \pi/2] \ni t \rightarrow g_0(t) = g(t)g_+(t) = \rho_\alpha \left[\begin{pmatrix} \sin t & \cos t \\ -\cos t & \sin t \end{pmatrix} \right],$$

that coincides with a maximal compact torus of $\mathbf{S}_\alpha$. We note that $g_0(\pi/2) = e$, while $g_0(-\pi/2) = \exp(i\pi H_\alpha)$. Hence $\varphi_{\omega_j}(g_0(\pi/2)) = 1$, and :

$$\varphi_{\omega_j}(g_0(-\pi/2)) = \exp(i\pi\omega_j(H_\alpha)) = \exp(i\pi(\omega_j|\alpha^\vee)) = (-1)^{(\omega_j|\alpha^\vee)}.$$

This proves (7.50). $\square$

8. THE FUNDAMENTAL GROUP OF PARABOLIC CR MANIFOLDS

In this section we give an explicit combinatorial description of the fundamental group of parabolic CR manifolds. We keep the notation of the previous sections. By using Proposition 6.18 and Theorem 7.16, we obtain :

Theorem 8.1. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and let $M = M(\mathfrak{g}, \mathfrak{q})$ be the corresponding homogeneous CR manifold. Then there exists a totally real parabolic CR algebra $(\mathfrak{g}, \mathfrak{q}')$ such that*

$$\mathbf{G}_+ = \{g \in \mathbf{G} \mid \text{Ad}_{\mathfrak{g}}(g)(\mathfrak{q}) \subset \mathfrak{q}\} \subset \mathbf{G}'_+ = \{g \in \mathbf{G} \mid \text{Ad}_{\mathfrak{g}}(g)(\mathfrak{q}') \subset \mathfrak{q}'\}$$

and the $\mathbf{G}$ -equivariant map

$$f : M = \mathbf{G}/\mathbf{G}_+ \rightarrow M' = M(\mathfrak{g}, \mathfrak{q}') = \mathbf{G}/\mathbf{G}'_+$$

has simply connected complex fibers. With $F = \mathbf{G}'_+/\mathbf{G}_+$, we obtain exact sequences :

$$(8.1) \quad \mathbf{1} \longrightarrow \pi_0(\mathbf{G}_+) \longrightarrow \pi_0(\mathbf{G}'_+) \longrightarrow \pi_0(F) \longrightarrow \mathbf{1},$$

$$(8.2) \quad \mathbf{1} \longrightarrow \pi_1(M) \longrightarrow \pi_1(M') \longrightarrow \pi_0(F) \longrightarrow \mathbf{1}.$$

The induced map in homotopy $f_* : \pi_1(M) \rightarrow \pi_1(M')$ is injective and $f_*(\pi_1(M))$ is a normal subgroup with finite index in $\pi_1(M')$.

Proof. Let $\mathfrak{q}'$ be the parabolic $\mathfrak{q}_m$ of Proposition 6.18. We proved that the typical fiber $F = \mathbf{G}'_+/\mathbf{G}_+$ is simply connected. Thus (8.1) and (8.2) are consequences of Serre's long exact sequence for the homotopy groups of a fiber bundle. By Theorem 7.16, $\pi_0(\mathbf{G}'_+)$ is a finite Abelian group. Hence $\pi_0(\mathbf{G}_+)$ is normal in $\pi_0(\mathbf{G}'_+)$ and $\pi_0(F) \simeq \pi_0(\mathbf{G}'_+)/\pi_0(\mathbf{G}_+)$ has a natural structure of Abelian group. Then also the maps in (8.2) are group homomorphisms and therefore $f_*(\pi_1(M))$ is a normal subgroup of $\pi_1(M')$. $\square$

Corollary 8.2. *If $\mathfrak{g}$ is a simple real Lie algebra of the complex type or of real type A II, A IIIa, A IV, B II, C II, D II, D IIIb, E III, E IV, F II, then all orbits $M = M(\mathfrak{g}, \mathfrak{q})$ are simply connected.*

Proof. The multiplicities of the simple real roots of $\hat{\mathfrak{g}}$ are always different from one, hence $M' = M(\mathfrak{g}, \mathfrak{q}')$ is simply connected, forcing M to be simply connected. $\square$

Note that the $M(\mathfrak{g}, \mathfrak{q}')$ for a totally real parabolic CR algebra $(\mathfrak{g}, \mathfrak{q}')$ are the *real flag manifolds*, and that there is a precise formula to compute their fundamental groups (see e.g. [DKV], [Wi]). Let $\mathfrak{h}'$ be a maximally noncompact Cartan subalgebra of $\mathfrak{g}$ contained in $\mathfrak{g}'_+$, fix an S-fit (and S-adapted) $C \in \mathfrak{C}(\mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}}'), \mathcal{Q}')$ and let $\{\alpha_1, \dots, \alpha_m\} = \mathcal{B}(C) \cap \mathcal{R}_{\text{re}}$. Then $\pi_1(M')$ is described by generators $t_{\alpha_1}, \dots, t_{\alpha_m}$ that satisfy the relations:

$$(8.3) \quad t_{\alpha_i} t_{\alpha_j} = t_{\alpha_j} t_{\alpha_i}^{(-1)^{(\alpha_i | \alpha_j^\vee)}} \quad \text{for } 1 \leq i, j \leq m, \quad t_{\alpha_i} = e \quad \text{if } \alpha_i \notin \Phi'_C.$$

By using Theorem 7.17 we can now give a description of $\pi_1(M)$:

Theorem 8.3. *Let $(\mathfrak{g}, \mathfrak{q})$ be an effective parabolic CR algebra and $(\vartheta, \mathfrak{h})$ an adapted Cartan pair for $(\mathfrak{g}, \mathfrak{q})$ with $\mathfrak{h}$ maximally noncompact in $\mathfrak{g}_+$. Let $\mathbf{H}$ be the corresponding Cartan subgroup of $\mathbf{G}$.*

Let $(\mathfrak{g}, \mathfrak{q}')$ be the totally real parabolic CR algebra of Theorem 8.1, $(\vartheta, \mathfrak{h}')$ an adapted Cartan pair for $(\mathfrak{g}, \mathfrak{q}')$ with $\mathfrak{h}'$ maximally noncompact in $\mathfrak{g}$.

Fix a Weyl chamber $C' \in \mathfrak{C}(\mathcal{R}', \mathcal{Q}')$, where $\mathcal{R}' = \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}}')$ and $\mathcal{Q}'$ is the parabolic set of $\mathfrak{q}'$ in $\mathcal{R}'$, that is S-fit to $(\mathfrak{g}, \mathfrak{q}')$ and S-adapted. With $\mathcal{B}(C') = \{\alpha_1, \dots, \alpha_\ell\}$ and $\mathcal{B}^(C') = \{\omega_1, \dots, \omega_\ell\}$ (defined by $(\omega_i | \alpha_j^\vee) = \delta_{i,j}$), if $\{\omega_1, \dots, \omega_\mu\} = \mathcal{B}^*(C') \cap [\mathcal{Q}'^r]^\perp$, we have $\{\alpha_1, \dots, \alpha_\mu\} = \Phi_{C'}(\mathcal{Q}') \cap \mathcal{R}'_{\text{re}}$. Let us consider the maps*

$$\varphi^b: \mathbf{G}'_+ \rightarrow \mathbb{Z}_2^\mu \text{ of (7.48)} \quad \text{and} \quad \delta: \pi_1(M') \rightarrow \mathbb{Z}_2^\mu \text{ of (7.50)}.$$

Then we have:

$$(8.4) \quad \pi_1(M) = \delta^{-1} \left(\phi^b(\mathbf{H}) \right). \quad \square$$

Corollary 8.4. *Let $(\mathfrak{g}, \mathfrak{q})$, be an effective parabolic CR algebra, $M = M(\mathfrak{g}, \mathfrak{q})$ the corresponding parabolic CR manifold. Let $M' = M(\mathfrak{g}, \mathfrak{q}')$ and $f: M \rightarrow M'$ be defined as in Theorem 8.1. If there is a Cartan subalgebra $\mathfrak{h}$ adapted to both $(\mathfrak{g}, \mathfrak{q})$ and $(\mathfrak{g}, \mathfrak{q}')$ and maximally noncompact in $\mathfrak{g}'_+$, such that moreover $\mathbf{H}/\mathbf{Z}(\mathbf{G})$ is connected, then $f_*: \pi_1(M) \rightarrow \pi_1(M')$ is an isomorphism.*

Proof. For the isotropy subgroup $\mathbf{G}'_+$ of M' we have, by (7.32), that $\pi_0(\mathbf{G}'_+) \simeq \pi_0(\mathbf{H}/(\mathbf{H} \cap \mathbf{S}'_0))$, where $\mathbf{S}'_0$ is an analytic semisimple subgroup of $\mathbf{G}'_+$. Since $\mathbf{Z}(\mathbf{G}) \subset \mathbf{G}_+$, the inclusion $\mathbf{H} \hookrightarrow \mathbf{G}'_+$ defines, passing to the quotients, a surjective map $\pi_0(\mathbf{H}/\mathbf{Z}(\mathbf{G})) \rightarrow \pi_0(\mathbf{G}'_+/\mathbf{G}_+)$. Thus the fiber of $f: M \rightarrow M'$ is connected and, by Theorem 8.1, $f_*: \pi_1(M) \rightarrow \pi_1(M')$ is an isomorphism. $\square$

Corollary 8.5. *With the notation of Corollary 8.4, if $\mathfrak{g}$ is a simple Lie algebra of real type A IIIb, D IIIa, then the map $f_*: \pi_1(M) \rightarrow \pi_1(M')$ is an isomorphism.*

Proof. In fact in these cases $\mathbf{H}/\mathbf{Z}(\mathbf{G})$ is connected for every choice of $\mathfrak{h}$, and we can apply Corollary 8.4 to obtain an isomorphism of the fundamental group of $M(\mathfrak{g}, \mathfrak{q})$ with the fundamental group of a totally real $M(\mathfrak{g}, \mathfrak{q}')$. $\square$

9. EXAMPLES

In this section we apply our results to several examples.

Example 9.1. Consider the semisimple real Lie algebra $\mathfrak{g} = \mathfrak{sl}(3, \mathbb{R})$ of real 3×3 matrices with zero trace. Let e_1, e_2, e_3 be the canonical basis of $\mathbb{R}^3 \subset \mathbb{C}^3$ and $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$ the basis of $\mathbb{C}^3$ given by:

$$\varepsilon_1 = e_1 + ie_2, \quad \varepsilon_2 = e_1 - ie_2, \quad \varepsilon_3 = e_3.$$

Let $\mathfrak{q}$ be the complex Borel subalgebra of complex matrices $Z \in \mathfrak{sl}(3, \mathbb{C})$ such that $Z(\langle \varepsilon_1 \rangle) \subset \langle \varepsilon_1 \rangle$ and $Z(\langle \varepsilon_1, \varepsilon_2 \rangle) \subset \langle \varepsilon_1, \varepsilon_2 \rangle$. Let $\mathfrak{h} \subset \mathfrak{q}$ be the Cartan subalgebra of traceless matrices that are diagonal in the basis $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$. The corresponding Cartan subgroup:

$$\mathbf{H} = \{\text{diag}(\lambda, \bar{\lambda}, |\lambda|^{-2}) \in \mathbf{SL}(3, \mathbb{R}) \mid \lambda \neq 0\}$$

is connected, hence also $\mathbf{G}_+$ is connected. There exists a unique Weyl chamber $C \in \mathfrak{C}(\mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}}))$ adapted to $(\mathfrak{g}, \mathfrak{q})$, which is both S-fit and V-fit. The corresponding diagram is:

$$\begin{array}{cc} \circledast & \text{---} & \oplus \\ \alpha_1 & & \alpha_2 \\ \times & & \times \end{array}$$

and we see that $M = M(\mathfrak{g}, \mathfrak{q})$ is weakly degenerate. The basis of the weakly nondegenerate reduction is the totally real parabolic CR manifold $M' = M(\mathfrak{g}, \mathfrak{q}')$, where $\mathfrak{q}' = \{Z \in \mathfrak{sl}(3, \mathbb{C}) \mid Z(\langle \varepsilon_1, \varepsilon_2 \rangle) \subset \langle \varepsilon_1, \varepsilon_2 \rangle\}$. Its diagrams, with respect to the Cartan subalgebras $\mathfrak{h}$ and $\mathfrak{h}'$, where $\mathfrak{h}'$ is the maximally noncompact Cartan subalgebra of traceless diagonal matrices in the basis (e_1, e_2, e_3) , are:

$$\begin{array}{cc} \circledast & \text{---} & \oplus \\ \alpha_1 & & \alpha_2 \\ \times & & \times \end{array} \simeq \begin{array}{cc} \circ & \text{---} & \circ \\ \alpha_1 & & \alpha_2 \\ \times & & \times \end{array}$$

By [Wi] the fundamental group of M' is $\pi_1(M') = \mathbb{Z}_2$. On the other hand, by Corollary 7.10 the fiber of the weakly nondegenerate reduction has two connected components. Hence the exact sequence (8.1) implies that M is simply connected.

Example 9.2. Let us compute the fundamental group of $M(\mathfrak{g}, \mathfrak{q})$ in the case of Example 6.1. Since in this case $\mathfrak{q}$ is Borel and the Cartan subalgebra $\mathfrak{h}$ is maximally compact, we know that the isotropy $\mathbf{G}_+ = \{X \in \mathfrak{g} \mid \text{ad}_{\mathfrak{g}}(X)(\mathfrak{q}) = \mathfrak{q}\}$ is connected (see [Kn, Prop.7.90]). Consider the fiber F over the point $(\langle e_1, e_4 \rangle, \langle e_1, e_2, e_4, e_5 \rangle)$. We can verify that F has 4 connected components and $\pi_0(F) \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$. The fundamental group of $M(\mathfrak{g}, \mathfrak{q}_3)$ can be computed using [Wi]. We have $\pi_1(M(\mathfrak{g}, \mathfrak{q}_3)) \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$ and thus, from the exact sequence:

$$\mathbf{1} \rightarrow \pi_1(M(\mathfrak{g}, \mathfrak{q})) \rightarrow \pi_1(M(\mathfrak{g}, \mathfrak{q}_3)) \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \pi_0(F) \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbf{1}$$

we obtain that $M(\mathfrak{g}, \mathfrak{q})$ is simply connected.

Example 9.3. Consider the complex flag manifold $\mathfrak{M}$ of $\mathbf{SO}(5, \mathbb{C})$, consisting of the complex projective lines contained in the quadric $\{z_2^2 + 2z_0z_4 + 2z_1z_3 = 0\} \subset \mathbb{CP}^4$. We identify the Lie algebra $\mathfrak{so}(5, \mathbb{C})$ to the matrix algebra:

$$\mathfrak{so}(5, \mathbb{C}) \simeq \hat{\mathfrak{g}} = \{Z \in \mathfrak{sl}(5, \mathbb{C}) \mid {}^t Z S_5 + S_5 Z = 0\} \quad \text{for} \quad S_5 = \begin{pmatrix} & & & & 1 \\ & & & 1 & \\ & & 1 & & \\ & 1 & & & \\ 1 & & & & \end{pmatrix}.$$

We consider the real form $\mathfrak{g} \simeq \mathfrak{so}(2, 3)$ of $\mathfrak{so}(5, \mathbb{C})$ defined by:

$$\mathfrak{so}(2, 3) \simeq \mathfrak{g} = \{Z \in \hat{\mathfrak{g}} \mid Z^* K + KZ = 0\} \quad \text{for} \quad K = \begin{pmatrix} & & & 1 \\ 1 & 0 & 0 & \\ 0 & -1 & 0 & \\ 0 & 0 & 1 & \end{pmatrix}.$$

Let M be the orbit of the projective line corresponding to the plane $\ell_2 = \langle e_1, e_2 \rangle$ of $\mathbb{C}^5$ by the action of the analytic subgroup $\mathbf{G}$ with Lie algebra $\mathfrak{g}$: M is the submanifold of the Grassmannian of the complex 2-planes in $\mathbb{C}^5$, consisting of those that are totally isotropic for the symmetric form S_5 and degenerate, with signature $(+, 0)$, with respect to the Hermitian symmetric form K . Denoting by $\mathfrak{q}$ the stabilizer of $\langle e_1, e_2 \rangle$ in $\mathbf{SO}(5, \mathbb{C})$, we have $M = M(\mathfrak{g}, \mathfrak{q})$. Take the Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ consisting of the diagonal matrices. With e_1, e_2 also denoting the value of the first and the second diagonal entry, we note that the conjugation σ is defined in $\mathfrak{h}_{\mathbb{R}}^*$ by $\sigma(e_i) = -(-1)^i e_i$. Take the Weyl chambers $C, C' \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ associated to the basis $\mathcal{B}(C) = \{e_1 - e_2, e_2\}$ and $\mathcal{B}(C') = \{e_2 - e_1, e_1\}$. Then C is S-fit and C' V-fit for $(\mathfrak{g}, \mathfrak{q})$. We can describe M by the cross-marked diagrams:

$$\begin{array}{ccc} \oplus & \xrightarrow{\quad} & \otimes \\ \alpha_1 & & \alpha_2 \\ & \times & \end{array} \quad \begin{array}{ccc} \ominus & \xrightarrow{\quad} & \circ \\ \alpha'_1 & & \alpha'_2 \\ & \times & \end{array}$$

From the first diagram we see that $(\mathfrak{g}, \mathfrak{q})$ is fundamental, since $\Phi_C = \{\alpha_2\}$ and $\bar{\alpha}_1 = \alpha_1 + 2\alpha_2 \succ_C \alpha_2$; from the second we see that $(\mathfrak{g}, \mathfrak{q})$ is weakly non-degenerate, because $\Phi_{C'} = \{\alpha'_2\}$ and $\bar{\alpha}'_2 = \alpha'_2 \succ_{C'} 0$. The weakest CR model of $(\mathfrak{g}, \mathfrak{q})$ is the parabolic CR algebra $(\mathfrak{g}, \mathfrak{v}) = (\mathfrak{g}, \mathfrak{q}_{\Phi_C^\#})$, with $\Phi_C^\# = \{\alpha_1, \alpha_2\}$. The weakly non-degenerate reduction of $(\mathfrak{g}, \mathfrak{v})$ is the totally real parabolic CR algebra $(\mathfrak{g}, \mathfrak{q}_{\Psi_C})$, with $\Psi_C = \{\alpha_1\}$:

$$\begin{array}{ccc} \oplus & \xrightarrow{\quad} & \otimes \\ \alpha_1 & & \alpha_2 \\ & \times & \end{array} \longrightarrow \begin{array}{ccc} \oplus & \xrightarrow{\quad} & \otimes \\ \alpha_1 & & \alpha_2 \\ & \times & \end{array}$$

By composition we obtain the $\mathbf{G}$ -equivariant projection:

$$M(\mathfrak{g}, \mathfrak{q}) \xrightarrow{\sim} M(\mathfrak{g}, \mathfrak{v}) \rightarrow M' = M(\mathfrak{g}, \mathfrak{q}_{\Psi_C = \{\alpha_1\}})$$

of M onto a totally real parabolic CR manifold M' . This projection associates to each $\ell_2 \in M$ the isotropic line $\ell_2 \cap \ell_2^\perp$, where $\perp$ is taken with respect to the Hermitian symmetric form K . The fiber over $\ell_1 = \langle e_1 \rangle$ consists of the planes generated by the columns of the matrices

$$\begin{pmatrix} 1 & 0 \\ 0 & z_0 \\ 0 & z_1 \\ 0 & z_2 \\ 0 & 0 \end{pmatrix} \quad \text{with} \quad \begin{cases} (z_0 : z_1 : z_2) \in \mathbb{CP}^2 \\ 2z_0 z_2 + z_1^2 = 0 \\ z_0 \bar{z}_0 + z_2 \bar{z}_2 > z_1 \bar{z}_1. \end{cases}$$

Then we see that the fiber is biholomorphic to $\mathbb{CP}^1 \setminus \mathbb{RP}^1$, that is the disjoint union of two disks in $\mathbb{C}$. Thus the fiber F of the projection $M \rightarrow M'$ has two connected components and $\pi_0(F) \simeq \mathbb{Z}_2$. Note that, by [Wi], $\pi_1(M') \simeq \mathbb{Z}$. Thus the Serre's exact sequence:


$$1 \longrightarrow \pi_1(M) \longrightarrow \pi(M') \simeq \mathbb{Z} \longrightarrow \pi_0(F) \simeq \mathbb{Z}_2 \longrightarrow 1$$

shows that $\pi_1(M) \simeq 2\mathbb{Z} \simeq \mathbb{Z}$.


Example 9.4. Let $(\varepsilon_i)_{1 \leq i \leq 4}$ be the canonical basis of $\mathbb{C}^4$, and $\mathfrak{g} \simeq \mathfrak{sl}(4, \mathbb{R})$ consist of the elements of $\mathfrak{sl}(4, \mathbb{C})$ that have real entries in the basis $(\varepsilon_i)_{1 \leq i \leq 4}$. We introduce the basis

$$e_1 = \varepsilon_1 + i\varepsilon_2, \quad e_2 = \varepsilon_1 - i\varepsilon_2, \quad e_3 = \varepsilon_3 + i\varepsilon_4, \quad e_4 = \varepsilon_3 - i\varepsilon_4.$$

We take the complex flag manifold $\mathfrak{M}$ whose points are the pairs (ℓ_1, ℓ_3) of a complex line ℓ_1 of $\mathbb{C}^4$ and a complex 3-plane ℓ_3 with $\ell_1 \subset \ell_3 \subset \mathbb{C}^4$, and consider the $\mathbf{G}$ -orbit M that contains the point $\mathfrak{o} = \langle e_1 \rangle \subset \langle e_1, e_2, e_3 \rangle$: we have $M = M(\mathfrak{g}, \mathfrak{q})$ where $\mathfrak{q}$ is the stabilizer in $\hat{\mathfrak{g}} \simeq \mathfrak{sl}(4, \mathbb{C})$ of $\mathfrak{o}$. Consider the Cartan subalgebra $\mathfrak{h}$ of the elements of $\mathfrak{g}$ that are diagonal matrices in the basis $(e_i)_{1 \leq i \leq 4}$. With $e_i(H)$ also denoting the value of the i -th entry of $H \in \mathfrak{h}_{\mathbb{R}}$, we note that $\mathcal{B}(C) = \{\alpha_i = e_i - e_{i+1} \mid 1 \leq i \leq 3\}$ is the system of simple roots for an S-fit Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$, and $\mathcal{B}(C) = \{\alpha'_1 = e_1 - e_3, \alpha'_2 = e_3 - e_2, \alpha'_3 = e_2 - e_4\}$ is the system of simple roots for a V-fit Weyl chamber. The corresponding cross-marked diagrams are:



Since $\bar{\alpha}_2 = \alpha_1 + \alpha_2 + \alpha_3$, from the first we see that $(\mathfrak{g}, \mathfrak{q})$ is fundamental, while the second shows that $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate. Its weakest CR model $(\mathfrak{g}, \mathfrak{v})$ has a totally real weakly nondegenerate reduction $(\mathfrak{g}, \mathfrak{q}')$, corresponding to diagrams:



where the last diagram is obtained by utilizing the Cartan subalgebra of real diagonal matrices of $\mathfrak{g}$ with respect to the canonical basis $(\varepsilon_i)_{1 \leq i \leq 4}$. Using [Wi], we obtain that $\pi_1(M(\mathfrak{g}, \mathfrak{q}')) \simeq \mathbb{Z}_2$. The isotropy subgroup $\mathbf{G}_+$ of $M(\mathfrak{g}, \mathfrak{q}')$ is isomorphic to the group of matrices of the form $\begin{pmatrix} A & B \\ 0 & C \end{pmatrix}$ with A, B, C real 2×2 matrices with $\det(A) \cdot \det(C) = 1$, and hence has two connected components. The isotropy subgroup $\mathbf{G}_+$ of $M(\mathfrak{g}, \mathfrak{q})$ is connected: indeed $\mathbf{G}_+ = \mathbf{N} \ltimes \mathbf{H}$ for an Euclidean $\mathbf{N} = \exp(\mathfrak{n})$ and a Cartan subgroup $\mathbf{H} = \mathbf{Z}_{\mathbf{G}}(\mathfrak{h})$ that is connected because $\mathfrak{h}$ is maximally noncompact (cf. [Kn, Proposition 7.90, p.488]). Thus the fiber $\mathbf{G}'_+/\mathbf{G}_+$ has two connected components. Thus, from the exact sequence:

$$1 \longrightarrow \pi_1(M(\mathfrak{g}, \mathfrak{q})) \longrightarrow \underbrace{\pi_1(M(\mathfrak{g}, \mathfrak{q}'))}_{\simeq \mathbb{Z}_2} \longrightarrow \underbrace{\pi_0(\mathbf{G}'_+/\mathbf{G}_+)}_{\simeq \mathbb{Z}_2} \longrightarrow 1$$


we obtain that $M(\mathfrak{g}, \mathfrak{q})$ is simply connected.

Example 9.5. Let $(\varepsilon_i)_{1 \leq i \leq 6}$ be the canonical basis of $\mathbb{C}^6$. Let $\mathbf{G} \simeq \mathbf{SL}(6, \mathbb{R})$, with Lie algebra $\mathfrak{g}$, be the subgroup of $\mathbf{SL}(6, \mathbb{C})$ consisting of the matrices with real entries. Consider the basis

$$e'_1 = \varepsilon_1 + i\varepsilon_4, \quad e'_2 = \varepsilon_2, \quad e'_3 = \varepsilon_3 + i\varepsilon_6, \quad e'_4 = \varepsilon_1 - i\varepsilon_4, \quad e'_5 = \varepsilon_5, \quad e'_6 = \varepsilon_3 + i\varepsilon_6.$$

Let $\mathfrak{q}$ be the stabilizer of $\langle e'_1, e'_2 \rangle \subset \langle e'_1, e'_2, e'_3, e'_4 \rangle$ in $\mathfrak{sl}(6, \mathbb{C})$ and consider the parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$. The matrices in $\mathfrak{g}$ that are diagonal in the basis $e'_1, \dots, e'_6$


form a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$. Then $\mathfrak{h}_{\mathbb{R}}$ consists of the matrices of $\hat{\mathfrak{g}} = \mathfrak{sl}(6, \mathbb{C})$ that are real and diagonal in the basis $e'_1, \dots, e'_6$. We identify e'_i to the evaluation function of the i -th diagonal term of $H \in \mathfrak{h}_{\mathbb{R}}$. Then the $\alpha'_i = e'_i - e'_{i+1}$ are the simple root of a $C' \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ that is V-fit for $(\mathfrak{g}, \mathfrak{q})$. We have the cross-marked diagram for $(\mathfrak{g}, \mathfrak{q})$:




Since $\Phi_{C'} = \{\alpha'_2, \alpha'_4\}$ and $\bar{\alpha}'_2 = \alpha'_2 + \alpha'_3 + \alpha'_4 + \alpha'_5 \succ_{C'} 0$, $\bar{\alpha}'_4 = \alpha'_1 + \alpha'_2 + \alpha'_3 + \alpha'_4 \succ_{C'} 0$, the CR algebra $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate. We obtain an S-fit Weyl chamber for $(\mathfrak{g}, \mathfrak{q})$ by reordering the basis $e'_1, \dots, e'_6$. Set:

$$e_1 = e'_2, e_2 = e'_1, e_3 = e'_4, e_4 = e'_3, e_5 = e'_6, e_6 = e'_5.$$

Then $\alpha_i = e_i - e_{i+1}$ ($1 \leq i \leq 5$) is the basis $\mathcal{B}(C)$ of an S-fit Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ yielding the cross-marked diagram:



Since $\Phi_C = \{\alpha_2, \alpha_4\}$ and $\bar{\alpha}_3 = \alpha_2 + \alpha_3 + \alpha_4 \succ_C \alpha_2, \alpha_4$, the CR algebra $(\mathfrak{g}, \mathfrak{q})$ is also fundamental. The weakest CR model $(\mathfrak{g}, \mathfrak{v})$ of $(\mathfrak{g}, \mathfrak{q})$ is obtained by taking the complex Borel subalgebra of $\hat{\mathfrak{g}}$ associated to the chamber C . We have $\mathfrak{v} = \mathfrak{q}_{\Phi_C^\#}$ with $\Phi_C^\# = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5\}$. The weakly nondegenerate reduction of $(\mathfrak{g}, \mathfrak{q})$ is the totally real CR algebra $(\mathfrak{g}, \mathfrak{q}_{\Psi_C})$ with $\Psi_C = \{\alpha_1, \alpha_3, \alpha_5\}$:



By composition we obtain a $\mathbf{G}$ -equivariant fibration:

$$M = M(\mathfrak{g}, \mathfrak{q}) \rightarrow M' = M(\mathfrak{g}, \mathfrak{q}_{\Psi_C}), \text{ with } (\ell_2 \subset \ell_4) \rightarrow (\ell_2 \cap \bar{\ell}_2 \subset \ell_4 \cap \bar{\ell}_4 \subset \ell_4 + \bar{\ell}_4).$$

The fiber F over $\langle e_1 \rangle \subset \langle e_1, e_2, e_3 \rangle \subset \langle e_1, e_2, e_3, e_4, e_5 \rangle$ consists of the pairs $(\ell_2 \subset \ell_4)$ where ℓ_2 is a complex 2-plane with $\langle e_1 \rangle \subset \ell_2 \subset \langle e_1, e_2, e_3 \rangle$ and ℓ_4 is a complex 4-plane with $\langle e_1, e_2, e_3 \rangle \subset \ell_4 \subset \langle e_1, e_2, e_3, e_4, e_5 \rangle$, with $\ell_2 \neq \bar{\ell}_2$ and $\ell_4 \neq \bar{\ell}_4$. Thus $\pi_0(F) \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$. On the other hand, by [Wi], we have $\pi_1(M') \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$. From the exact sequence

$$\mathbf{1} \longrightarrow \pi_1(M) \longrightarrow \pi_1(M') \simeq \mathbb{Z}_2^3 \longrightarrow \pi_0(F) \simeq \mathbb{Z}_2^2 \longrightarrow \mathbf{1}$$

we obtain that $\pi_1(M) \simeq \mathbb{Z}_2$.


Example 9.6. Let $\hat{\mathfrak{g}} \simeq \mathfrak{so}(5, \mathbb{C})$ be as in Example 9.3. We take now $\mathfrak{g} \simeq \mathfrak{so}(2, 3)$, defined by:

$$\mathfrak{g} = \{Z \in \hat{\mathfrak{g}} \mid Z^* K + K Z = 0\} \quad \text{with} \quad K = \begin{pmatrix} & & & & 1 \\ & & & & \\ & & & & \\ & & & & \\ 1 & & & & \end{pmatrix}.$$


Let $\mathfrak{q}$ be the stabilizer of the line $\langle e_1 \rangle \subset \mathbb{C}^5$ and consider the orbit $M = M(\mathfrak{g}, \mathfrak{q})$. Our M is one of the two connected components of the manifold M^+ the nonreal

lines ℓ_1 , contained in the quadric cone $\{2z_0z_4 + 2z_1z_3 + z_2^2 = 0\} \subset \mathbb{C}^5$, for which $(\ell_1 + \bar{\ell}_1)$ is a totally isotropic complex 2-plane for the Hermitian symmetric form K . The involution $\ell_1 \rightarrow \bar{\ell}_1$ interchanges the two connected components of M^+ .

The diagonal matrices in $\mathfrak{g}$ define a Cartan subalgebra of $\mathfrak{g}$ adapted to $(\mathfrak{g}, \mathfrak{q})$. As usual we denote again by e_i the value of the i -th entry in the diagonal of $\mathfrak{h}_{\mathbb{R}}$. Then we define a V-fit C' and an S-fit C by taking $\mathcal{B}(C') = \{\alpha'_1 = e_1 + e_2, \alpha'_2 = -e_2\}$ and $\mathcal{B}(C) = \{\alpha_1 = e_1 - e_2, \alpha_2 = e_2\}$. The associated cross-marked diagrams are:




From the first, as $\mathfrak{q} = \mathfrak{q}_{\Phi_{C'}}$ with $\Phi_{C'} = \{\alpha'_1\}$ and $\bar{\alpha}'_1 = \alpha'_1 \succ_{C'} 0$, we see that $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate; from the second, as $\mathfrak{q} = \mathfrak{q}_{\Phi_C}$ with $\Phi_C = \{\alpha_1\}$ and $\bar{\alpha}_2 = \alpha_1 + \alpha_2 \succ_C \alpha_1$, we see that $(\mathfrak{g}, \mathfrak{q})$ is also fundamental. The weakest CR model of $(\mathfrak{g}, \mathfrak{q})$ is $(\mathfrak{g}, \mathfrak{q}_{\Psi_C^\#})$ with $\Psi_C^\# = \{\alpha_1, \alpha_2\}$. The basis of its weakly nondegenerate reduction is the totally real CR algebra $(\mathfrak{g}, \mathfrak{q}_{\Psi_C})$ with $\Psi_C = \{\alpha_2\}$. We can represent these maps by the diagram:




where the last is the cross-marked Satake diagram of $(\mathfrak{g}, \mathfrak{q}_{\Psi_C})$, i.e. the cross-marked diagram for an S-fit and S-adapted C and a maximally noncompact $\mathfrak{h}$. We have, by [Wi, Theorem 1.1], $\pi_1(M(\mathfrak{g}, \mathfrak{q}_{\Psi_C})) \simeq \mathbb{Z}_2$. We observe that the stabilizer in the connected component of the identity of $\mathbf{SO}(2, 3)$ of a totally isotropic 2-plane $\ell_2 \subset \mathbb{R}^5$ keeps its orientation and is connected. Thus the fiber F of the projection $M(\mathfrak{g}, \mathfrak{q}) \rightarrow M(\mathfrak{g}, \mathfrak{q}_{\Psi_C})$ is connected because the isotropy subgroup of $M(\mathfrak{g}, \mathfrak{q}_{\Psi_C})$ is connected. Therefore $\pi_1(M(\mathfrak{g}, \mathfrak{q})) \simeq \pi_1(M(\mathfrak{g}, \mathfrak{q}_{\Psi_C})) \simeq \mathbb{Z}_2$.

Example 9.7. Let $\mathfrak{g}$ be a simple real Lie algebra of type FI (split real form). We fix a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$, such that the conjugation σ defined in $\hat{\mathfrak{g}}$ by the real form $\mathfrak{g}$ restricts in $\mathfrak{h}_{\mathbb{R}}^* = \mathbb{R}^4 = \langle e_1, e_2, e_3, e_4 \rangle_{\mathbb{R}}$ to the linear involution that is defined on the canonical basis by: $\sigma(e_1) = -e_3$, $\sigma(e_2) = e_4$, $\sigma(e_3) = -e_1$, $\sigma(e_4) = e_2$. The vectors $\alpha_1 = e_2 - e_3$, $\alpha_2 = e_3 - e_4$, $\alpha_3 = e_4$ and $\alpha_4 = \frac{1}{2}(e_1 - e_2 - e_3 - e_4)$ are a basis of simple roots of $\mathcal{R} = \mathcal{R}(\hat{\mathfrak{g}}, \hat{\mathfrak{h}})$. We consider the parabolic CR manifold $M(\mathfrak{g}, \mathfrak{q})$ that corresponds to the cross-marked diagram:



This is a representation of $(\mathfrak{g}, \mathfrak{q})$ in a V-fit Weyl chamber $C \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$. Since $\bar{\alpha}_1 \succ_C 0$, $\bar{\alpha}_3 \succ_C 0$, by Theorem 5.9 the CR algebra $(\mathfrak{g}, \mathfrak{q})$ is weakly nondegenerate.


Consider the Weyl chamber C_1 obtained from C by the symmetry s_{α_2} . We have $\mathcal{B}(C_1) = \{e_2 - e_4, e_4 - e_3, e_3, \frac{1}{2}(e_1 - e_2 - e_3 - e_4)\} = \{\alpha'_1, \alpha'_2, \alpha'_3, \alpha'_4\}$ and the cross-marked diagram for $(\mathfrak{g}, \mathfrak{q})$ for the chamber $C_1 \in \mathfrak{C}(\mathcal{R}, \mathcal{Q})$ is:




This diagram is S-fit. Since $\bar{\alpha}'_2 = 2\alpha'_1 + 3\alpha'_2 + 4\alpha'_3 + 2\alpha'_4$, by Theorem 5.5 the parabolic CR algebra $(\mathfrak{g}, \mathfrak{q})$ is also fundamental.

Our next aim is to construct the fibration of Proposition 6.18. First we observe that the weakest CR model of $(\mathfrak{g}, \mathfrak{q})$ is $(\mathfrak{g}, \mathfrak{v}_1)$ with $\mathfrak{v}_1 = \mathfrak{q}_{\{\alpha'_1, \alpha'_2, \alpha'_3\}}$. Its weakly nondegenerate reduction is $(\mathfrak{g}, \mathfrak{q}_2)$ with $\mathfrak{q}_2 = \mathfrak{q}_{\{\alpha'_2\}}$.

To compute its weakest CR model, we need to find an S-fit Weyl chamber C_2 for $(\mathfrak{g}, \mathfrak{q}_2)$. This is provided by the basis of simple roots $\mathcal{B}(C) = \{\beta_1, \beta_2, \beta_3, \beta_4\}$ with $\beta_1 = e_2 - e_4$, $\beta_2 = e_1 - e_2$, $\beta_3 = \frac{1}{2}(e_1 - e_2 - e_3 - e_4)$, $\beta_4 = -\frac{1}{2}(e_1 - e_2 + e_3 - e_4)$, that is obtained from $\mathcal{B}(C_1)$ by the rotation $s_{\frac{1}{2}(e_1 - e_2 + e_3 - e_4)} \circ s_{e_3}$ of $\mathbf{W}(\mathcal{R})$. The corresponding cross-marked diagram for $(\mathfrak{g}, \mathfrak{q}_2)$ is:



Since $\bar{\beta}_4 = \beta_1 + 2\beta_2 + 2\beta_3 + \beta_4$, the weakest CR model of $(\mathfrak{g}, \mathfrak{q}_2)$ is $(\mathfrak{g}, \mathfrak{v}_2)$ with $\mathfrak{v}_2 = \mathfrak{q}_{\{\beta_2, \beta_4\}}$. The CR algebra $(\mathfrak{g}, \mathfrak{v}_2)$ has the weakly nondegenerate reduction $(\mathfrak{g}, \mathfrak{q}_{\{\beta_4\}})$. The element $A = (-1, -1, +1, -1)$ of $\mathfrak{h}_{\mathbb{R}}$ defines the parabolic set of $\mathfrak{v}_2$ and $\bar{A} = A$ shows then that $(\mathfrak{g}, \mathfrak{v}_2)$ is totally real. Thus, by choosing a maximally noncompact Cartan subalgebra $\mathfrak{h}'$ adapted to $(\mathfrak{g}, \mathfrak{v}_2)$, we can associate to $(\mathfrak{g}, \mathfrak{v}_2)$ its cross-marked Satake diagram as a totally real parabolic minimal CR algebra:



Then the isotropy subgroup $\mathbf{G}'_+$ of $M(\mathfrak{g}, \mathfrak{v}_2)$ has two connected components, and the fundamental group $\pi_1(M(\mathfrak{g}, \mathfrak{v}_2))$ is isomorphic to $\mathbb{Z}_2$ and is generated by any simple path joining the two connected components of $\mathbf{G}'_+$. By using Lemma 7.1 and (7.32) of Theorem 7.8, we find that $\mathbf{H}$ has four connected components and $\varphi^b(\mathbf{H})$ has two connected components. Hence Theorem 8.3 yields $\pi_1(M) \simeq \mathbb{Z}_2$.

REFERENCES

- [AF] A.Andreotti, G.Fredricks, *Embeddability of real analytic Cauchy-Riemann manifolds*, Ann. Scuola Norm. Sup. Pisa (4) **6** (1979), 285-304.
- [AFN] A.Andreotti, G.Fredricks, M.Nacinovich, *On the absence of Poincaré lemma in Tangential Cauchy-Riemann complexes*, Ann. Scuola Norm. Sup. Pisa **8** (1981), 365-404.
- [AMN] A.Altomani, C.Medori, M.Nacinovich, *The CR structure of minimal orbits in complex flag manifolds*, J. Lie Theory **16** (2006), 483-530.
- [Ar] S.Araki, *On root systems and an infinitesimal classification of irreducible symmetric spaces*, J. Math. Osaka City Univ. **A 13** (1962), 1-34.
- [BER] M.S.Baouendi, P.Ebenfelt, L.Preiss Rothschild, *Real submanifolds in complex space and their mappings*, Princeton University Press, Princeton, NJ, 1999.
- [BIG] T.Bloom, I.Graham, *A geometric characterization of points of type m on real submanifolds of $\mathbb{C}^n$* , J. Differential Geometry **12** (1977), 171-182.
- [BL] R.Bremigan, J.Lorch, *Orbit duality for flag manifolds*, Manuscripta Math. **109** (2002), 233-261.
- [BT1] A.Borel, J.Tits, *Groupes réductifs*, Publ. Math. IHES **27** (1965), 55-151.
- [BT2] ———, *Compléments à l'article "Groupes réductifs"*, Publ. Math. IHES **41** (1972), 253-276.
- [B1] N.Bourbaki, *Groupes et algèbres de Lie, Ch. 4,5 et 6*, Masson, Paris, 1981.
- [B2] ———, *Groupes et algèbres de Lie, Ch. 7 et 8*, Masson, Paris, 1990.
- [CS] L.Casian, R.J.Stanton, *Schubert cells and representation theory*, Invent. Math. **137** (1999), 461-539.

- [Ch] C.Chevalley, *Théorie des groupes de Lie*, Hermann, Paris, 1955.
- [DKV] J.J.Duistermaat, J.A.C.Kolk, V.S.Varadarajan, *Functions, flows and oscillatory integrals on flag manifolds and conjugacy classes in real semisimple Lie groups*, *Compositio Math.* **49** (1983), 309-398.
- [Fe] G.Fels, *Locally homogeneous finitely nondegenerate CR manifolds*; preprint, arXiv:math.CV/0606032, Jun 2006, 1-21.
- [FHW] G.Fels, A.T.Huckleberry, J.A.Wolf, *Cycle spaces of flag domains. A complex geometric viewpoint*, *Progress in Mathematics*, vol. 245, Birkhäuser, Boston, 2006.
- [FeK] G.Fels, W.Kaup, *Classification of Levi degenerate homogeneous CR manifolds of dimension 5*; arXiv.org:math/0610375 2006-10-11, 1-49.
- [GM] S.Gindikin, T.Matsuki, *Stein extensions of Riemannian symmetric spaces and dualities of orbits on flag manifolds*, *Transform. Groups* **8** (2003), 333-376.
- [HN] C.D.Hill, M.Nacinovich, *Pseudoconcave CR manifolds*, *Complex analysis and geometry*, *Lect. Notes in Pure and Appl. Math.*, vol. 173, Marcel and Dekker, New York, 1996, pp. 275-297.
- [HW] A.T.Huckleberry, J.A.Wolf, *Schubert varieties and cycle spaces*, *Duke Math. J.* **120** (2003), 229-249.
- [Ka] M.Kashiwara, *D-modules and representation theory of Lie groups*, *Ann. Inst. Fourier* **43** (1993), 1597-1618.
- [KZ] W.Kaup, D.Zaitsev, *On the CR-structure of compact group orbits associated with bounded symmetric domains*, *Invent. Math.* **153** (2003), 45-104.
- [Kn] A.Knapp, *Lie groups beyond an introduction (2nd ed.)*, Birkhäuser, Boston, 2002.
- [LN] A.Lotta, M.Nacinovich, *On a class of symmetric CR manifolds*, *Adv. in Math.* **191** (2005), 114-146.
- [MN1] C.Medori, M.Nacinovich, *Levi-Tanaka algebras and homogeneous CR manifolds*, *Compositio Math.* **109** (1997), 195-250.
- [MN2] ———, *Classification of semisimple Levi-Tanaka algebras*, *Ann. Mat. Pura Appl.* **174** (1998), 285-349.
- [MN3] ———, *Standard CR manifolds of codimension 2*, *Transformation groups* **6** (2001), 53-78.
- [MN4] ———, *Algebras of infinitesimal CR automorphisms*, *Journal of Algebra* **287** (2005), 234-274.
- [Mo] G.D.Mostow, *On maximal subgroups of real Lie groups*, *Ann. Math.* **74** (1961), 503-517.
- [Po] M.M.Postnikov, *Lie groups and Lie algebras*, Nauka, Moskow, 1982; French translation MIR 1985.
- [OV] A.L. Onishchik, E.B. Vinberg (eds.), *Lie groups and Lie algebras III*, *Encyclopaedia of Mathematical Sciences*, vol. 43, Springer, Berlin, 1994.
- [S1] M.Sugiura, *Conjugate classes of Cartan subalgebras in real semisimple Lie algebras*, *J. Math. Soc. Japan* **11** (1959), 374-434.
- [S2] ———, *Correction to my paper: Conjugate classes of Cartan subalgebras in real semisimple Lie algebras*, *J. Math. Soc. Japan* **23** (1971), 379-383.
- [Wa] G.Warner, *Harmonic analysis on semi-simple Lie groups I*, Springer, Berlin, 1972.
- [Wi] M.Wiggerman, *The fundamental group of a real flag manifold*, *Indag.Matem.,N.S.* **9** (1998), 141-153.
- [W] J.A.Wolf, *The action of real semisimple Lie groups on a complex manifold I: Orbit structure and holomorphic arc components*, *Bull. Amer. Math. Soc.* **75** (1969), 1121-1137.

A. ALTOMANI: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA (ITALY)

E-mail address: altomani@mat.uniroma2.it

C. MEDORI: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PARMA, PARCO AREA DELLE SCIENZE 53/A, 43100 PARMA (ITALY)

E-mail address: medori@prmat.math.unipr.it

M. NACINOVICH: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA (ITALY)

E-mail address: nacinovi@mat.uniroma2.it